

GP OOHSS

11 October 2020

GP Out-of-Hours Surveillance System: England

Data to:

14 October 2020 Year: 2020 Week: 41

In This Issue:

Key Messages.

Weekly summary.

Total contacts.

Syndromic indicators.

Notes and caveats.

Further information.

Acknowledgements.

Key messages

During week 41, GP out of hours respiratory indicators either remained stable or decreased slightly (figures 2-6). Contacts for difficulty breathing/ wheeze/asthma were stable but remain elevated in children aged 1-4 years (figures 5 & 5a).

Syndromic indicators at a glance:


Number of contacts and percentage of Read coded contacts.

	No. of	%	%	
Key indicator	contacts	Week 41	Week 40	Trend*
All OOH contacts, all causes	165,481			
Acute respiratory infection	4,088	6.95	7.45	$lack \Psi$
Influenza-like illness	142	0.24	0.23	←→
Bronchitis/bronchiolitis	40	0.07	0.10	←→
Difficulty breathing/wheeze/asthma	1,280	2.18	2.14	←→
Pharyngitis	15	0.03	0.04	$lack \Psi$
Gastroenteritis	1,538	2.61	2.51	^
Diarrhoea	454	0.77	0.81	←→
Vomiting	676	1.15	1.03	^
Chest pain/myocardial infarction	862	1.46	1.40	^

^{*}Trend: reports on the trend seen over previous weeks in the percentage of Read coded contacts.

1: Total out-of-hours contacts:

Daily total number of out-of-hours and unscheduled contacts and 7-day moving average (adjusted for bank holidays).







2: Acute Respiratory Infection daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7-day moving average*.

2a: Acute Respiratory Infection by age group.

As a percentage of total contacts within each age group, shown as a 7-day moving average*.






Intentionally left blank

*7-day moving average adjusted for bank holidays.




3: Influenza-like illness daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7-day moving average*.



3a: Influenza-like illness by age group.

As a percentage of total contacts within each age group, shown as a 7-day moving average*.



Intentionally left blank

*7-day moving average adjusted for bank holidays.



4: Bronchitis/ bronchiolitis daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7day moving average*.

4a: Bronchitis/ bronchiolitis daily contacts by age group*.

As a percentage of total contacts within each age group, shown as a 7-day moving average*.


5: Difficulty breathing/ wheeze/asthma daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7day moving average*.

*7-day moving average adjusted for bank holidays.

Bank Holiday

weekend



Diff.BreathingWheezeAsthma -

■7 day moving average* ■■■ Baseline



5a: Difficulty breathing/wheeze/ asthma daily contacts by age group*.

As a percentage of total contacts within each age group, shown as a 7-day moving average *.

6: Acute pharyngitis and persistent sore

Shown as a percentage of the total contacts with a Read code and as a 7day moving average*.

throat.

7: Gastroenteritis daily contacts

Shown as a percentage of the total contacts with a Read code and as a 7day moving average*.


*7-day moving average adjusted for bank holidays.

0 13/10/19 10/11/19 08/12/19 05/01/20 02/02/20 01/03/20

weekend

Gastroenteritis -

Bank Holiday



29/03/20 26/04/20 24/05/20 21/06/20 19/07/20 16/08/20 13/09/20 11/10/20 08/11/20

7 day moving average*

--- Baseline



14 October 2020

8: Diarrhoea daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7 -day moving average*.

8a: Diarrhoea daily contacts by age group*.

As a percentage of total contacts within each age group, shown as a 7-day moving average*.

9: Vomiting daily contacts.


Shown as a percentage of the total contacts with a Read code and as a 7 -day moving average*.

*7-day moving average adjusted for bank holidays.

Bank Holiday

weekend

Vomiting -




--- Baseline

7 day moving average*



10: Chest pain/ myocardial infarction daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7 -day moving average*.



Intentionally left blank

Intentionally left blank

*7-day moving average adjusted for bank holidays.



Notes and caveats:

- This bulletin presents data from the Public Health England (PHE) GP Out
 -of-hours\Unscheduled Care Surveillance System (GP OOHSS).
- Fully anonymised data from GP out-of-hours (OOH) and unscheduled care service providers in England are being transferred to PHE for analysis and interpretation by the PHE Real-time Syndromic Surveillance Team (ReSST).
- This system supplements existing PHE syndromic surveillance systems by monitoring data on GP consultations outside of routine surgery opening times (evenings, weekends and bank holidays) and unplanned contacts within NHS primary care.
- The key indicators presented within this bulletin are derived by grouping selected Read coded consultations.
- GP OOH consultation data are analysed on a daily basis to identify
 national and regional trends. A statistical algorithm underpins each
 system, routinely identifying activity that has increased significantly or is
 statistically significantly high for the time of year. Results from these daily
 analyses are assessed by the ReSST, along with analysis by age group,
 and anything deemed of public health importance is alerted by the team.
- Baselines represent seasonally expected levels of activity and are constructed from historical data since Nov 2009. They take into account any known substantial changes in data collection, population coverage or reporting practices. Gastroenteritis, diarrhoea and vomiting baselines also account for changes since the introduction of rotavirus vaccine in July 2013. Baselines are refreshed using the latest data on a regular basis however they currently exclude data from 2020 due to the COVID-19 pandemic affecting GP services and patient health care seeking behaviour.

Moving Epidemic Method (MEM):

- During winter we present Moving Epidemic Method (MEM) influenza thresholds on selected indicators. MEM is a standard methodology used for setting influenza thresholds across many European nations.¹
- MEM is used for GP OOH ILI thresholds at a national level.
- MEM thresholds should be interpreted using 7 day moving averages rather than daily data.
- MEM thresholds currently use six years of historic data (2013-2019). The thresholds are re-calculated every year.
- Baseline ('Pre-epidemic') thresholds are used alongside other surveillance systems to identify the start of influenza circulating in the community
- 40%, 95% and 97.5% intensity thresholds are used to identify when influenza activity moves from low to medium, high or very high.
 1Vega T et al. Influenza Other Respir Viruses. 2013;7(4):546-58.

Further information:

The GP Out-of-Hours Surveillance System Bulletin can also be downloaded from the PHE Real-time Syndromic Surveillance website which also contains more information about syndromic surveillance:

https://www.gov.uk/government/collections/syndromic-surveillance-systems-and-analyses

Acknowledgements:

We are grateful to Advanced and the GP OOH and unscheduled care service providers who have kindly agreed to participate in this system.

Contact ReSST:

<u>syndromic.surveillance</u> <u>@phe.gov.uk</u>

PHE Out-of-Hours/Unscheduled Care Surveillance

Produced by: PHE Real-time Syndromic Surveillance Team
1st Floor, 5 St Philips Place, Birmingham, B3 2PW

Tel: 0344 225 3560 > Option 4 > Option 2 Fax: 0121 236 2215

Web: https://www.gov.uk/government/collections/syndromic-surveillance-systems-and-analyses