

GP In Hours

Syndromic Surveillance System: England

20 January 2021

Year: 2021 Week: 2

In This Issue:

Key messages.

Diagnostic indicators at a glance.

GP practices and denominator population.

National syndromic indicators.

Notes and further information.

Key messages

Data to:

17 January 2021

During week 2 COVID-19-like, influenza-like illness (ILI) and asthma consultations levelled off and decreased slightly (figures 1, 3 & 11). ILI consultations remain highest in adults aged 45-64 years and London, South East and North West, but are now decreasing (figures 3a & 3b).

Please note: COVID-19-like GP consultations presented in this report are currently based on reporting by one GPIH data provider and therefore a smaller population denominator. All other indicators remain unaffected.

Note: during the COVID-19 pandemic, patients with COVID-19 symptoms are generally advised to initially access a COVID-19 test through the national COVID-19 testing programme. This is likely to result in lower numbers of patients accessing health advice as monitored through syndromic surveillance systems. Syndromic data should therefore be interpreted with some caution and in the context of other COVID-19 monitoring data sources. Please see 'notes and caveats' for information about the COVID-19-like GPIH syndromic indicator including important caveats around the interpretation of this indicator.

A Cold Watch System operates in England from 1 November to 31 March each year. As part of the Public Health England Cold Weather Plan for England the PHE Real-time Syndromic Surveillance team will be monitoring the impact of cold weather on syndromic surveillance data during this period.

Cold weather alert level (current reporting week): Level 1/3 - Winter preparedness/Cold weather action http://www.metoffice.gov.uk/weather/uk/coldweatheralert/

Diagnostic indicators at a glance:

Indicator	Trend	Level
COVID-19-like	no trend	-
Upper respiratory tract infection	no trend	below baseline levels
Influenza-like illness	no trend	below baseline levels
Pharyngitis	no trend	below baseline levels
Scarlet fever	no trend	below baseline levels
Lower respiratory tract infection	no trend	below baseline levels
Pneumonia	no trend	below baseline levels
Gastroenteritis	no trend	below baseline levels
Vomiting	no trend	below baseline levels
Diarrhoea	no trend	below baseline levels
Asthma	no trend	below baseline levels
Conjunctivitis	no trend	below baseline levels
Mumps	no trend	below baseline levels
Measles	no trend	below baseline levels
Rubella	no trend	below baseline levels
Pertussis	no trend	below baseline levels
Chickenpox	no trend	below baseline levels
Herpes zoster	no trend	below baseline levels
Cellulitis	no trend	below baseline levels
Impetigo	decreasing	below baseline levels

GP practices and denominator population:

Year	Week	GP Practices Reporting**	Population size**
2021	2	4,319	39.5 million

^{**}based on the average number of practices and denominator population in the reporting working week.



1. COVID-19-like consultations

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, based on a denominator population of approximately 5.5 million patients).

Please note: based on a single data provider only.

1a: COVID-19-like consultations by age group

Daily incidence rate by age group per 100,000 population all England, based on a denominator population of approximately 5.5 million patients).


Please note: based on a single data provider only


1b: COVID-19-like consultations by PHE Centre

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, based on a denominator population of approximately 5.5 million patients).


Please note: based on a single data provider only

* 7-day moving average adjusted for bank holidays.






covid-19 by PHE Centre 19/01/2020 - 17/01/2021







2: Upper respiratory tract infection (URTI)

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).



Average daily incidence rate by week per 100,000 population (all England).





^{* 7-}day moving average adjusted for bank holidays.



3: Influenza-like illness (ILI)


Daily incidence rates (and 7-day moving average*) per 100,000 population (all England, all ages).


3a: Influenza-like illness by age


Average daily incidence rate by week per 100,000 population (all England).


3b: Influenza-like illness by PHE Centre

Average daily incidence rate by week per 100,000 population (all ages).











4: Pharyngitis or scarlet fever


Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).


5: Scarlet fever

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, based on a denominator population of approximately 5.5 million patients)

5a: Scarlet fever by age

Average daily incidence rate by week per 100,000 population (all England, based on a denominator population of approximately 5.5 million patients).







20 January 2021 Year


6: Lower respiratory tract infection (LRTI)

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).

6a: Lower respiratory tract infection (LRTI) by age

Average daily incidence rate by week per 100,000 population (all England).





Intentionally left blank

^{* 7-}day moving average adjusted for bank holidays.



20 January 2021


Year: 2021 Week: 2


7: Pneumonia

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).

7a: Pneumonia by age

Average daily incidence rate by week per 100,000 population (all England).





Intentionally left blank

^{* 7-}day moving average adjusted for bank holidays.




20 January 2021

Year: 2021 Week: 2


8: Gastroenteritis

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).




8a: Gastroenteritis by age

Average daily incidence rate by week per 100,000 population (all England).



9: Vomiting

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).



* 7-day moving average adjusted for bank holidays.



9a: Vomiting by age


Average daily incidence rate by week per 100,000 population (all England).


10: Diarrhoea


Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).


10a. Diarrhoea by age

Average daily incidence rate by week per 100,000 population (all England).













11: Asthma

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).




11a: Asthma by age

Average daily incidence rate by week per 100,000 population (all England).



12: Conjunctivitis

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).



^{* 7-}day moving average adjusted for bank holidays.



13: Mumps


Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).


14: Measles

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).

15: Rubella

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).







16: Pertussis

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, based on a denominator population of approximately 5.5 million patients)


17: Chickenpox


Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, based on a denominator population of approximately 5.5 million patients)


18: Herpes zoster

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).

* 7-day moving average adjusted for bank holidays.











19 Cellulitis

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).

20: Impetigo

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).





Intentionally left blank

^{* 7-}day moving average adjusted for bank holidays.



Notes and further information

- The PHE GP in hours surveillance system is a syndromic surveillance system monitoring community-based morbidity recorded by GP practices.
- GP consultation data are analysed on a daily basis to identify national and regional trends. A statistical algorithm underpins each system, routinely identifying activity that has increased significantly or is statistically significantly high for the time of year. Results from these daily analyses are assessed along with analysis by age group, and anything deemed of public health importance is alerted.
- This system captures anonymised GP morbidity data from two GP clinical software systems, EMIS, from version 1 of the QSurveillance® database, and TPP SystmOne.
- Baselines represent seasonally expected levels of activity and are constructed from historical data since April 2012. They take into account any known substantial changes in data collection, population coverage or reporting practices.
 Gastroenteritis, diarrhoea and vomiting baselines also account for changes since the introduction of rotavirus vaccine in July 2013. Baselines are refreshed using the latest data on a regular basis however they currently exclude data from 2020 due to the COVID-19 pandemic affecting GP services and patient health care seeking behaviour.

COVID-19 consultations

- A collection of new COVID-19 Snomed codes were released in March 2020 to facilitate the recording of patients presenting to primary care services with symptoms of COVID-19.
- The GPIH surveillance system monitors the use of these codes in a selection of TPP and EMIS practices across England.
- However, patients presenting with COVID-19 symptoms may be diagnosed using other clinical codes used by the GP.
- Therefore, the COVID-19-like indicator presented in this report is primarily for monitoring trends in GP consultations, and it must be interpreted in context with the other respiratory syndromic indicators presented in this report. The number/ rate of COVID-19-like consultations should therefore not be used as an absolute count of those patients with COVID-19.
- During April 2020 a new COVID-19 Care Pathway template was introduced into GP systems that has affected recording of influenza-like illness (ILI), resulting in an increase in the consultation rate for ILI (figures 2a-c).
- All indicator trends should be interpreted with caution due to current national advice and guidance regarding access to GP surgeries and changes in clinical coding for COVID-19.
- Centre level COVID-19 consultation data should be interpreted with some caution.
 Different GP clinical system providers have different coding for COVID-19 and
 therefore rates can differ between Centres depending on the relative contribution
 of individual GP system providers in GPIH. Centre-specific data should not be
 compared across Centres: trends should only be interpreted for each individual
 Centre.

Acknowledgements:

We thank and acknowledge the University of Oxford, ClinRisk[®] and the contribution of EMIS and EMIS practices. Data source: version 1 of the QSurveillance® database.

We thank TPP, ResearchOne and the SystmOne GP practices contributing to this surveillance system.

Contact ReSST: syndromic.surveillance @phe.gov.uk

GP In Hours Syndromic Surveillance System Bulletin.

Produced by: PHE Real-time Syndromic Surveillance Team 1st Floor, 5 St Philips Place, Birmingham, B3 2PW

Tel: 0344 225 3560 > Option 4 > Option 2 Fax: 0121 236 2215

Web: https://www.gov.uk/government/collections/syndromic-surveillance-systems-and-analyses