



Trident Reference No: RB292404  
ZEN Ref: TR1899592

**CUSTOMER INFO:**  
Minimum Point Size = 7.00pt

N521

Reckitt Benckiser Healthcare  
UK Ltd., Slough SL1 3UH  
PL 00063/0649



**CONTAINS PARACETAMOL.** Do not take anything else containing paracetamol while taking this medicine. Talk to a doctor at once if you take too much of this medicine, even if you feel well. This is because too much paracetamol can cause delayed, serious liver damage.

**DO NOT TAKE IF YOU:**

- Have (or have had) two or more episodes of a stomach ulcer, perforation or bleeding
- Are allergic to ibuprofen, paracetamol or any other ingredient of the product, aspirin or other related
- Headache
- Dental pain
- Cold and flu symptoms
- Period pain
- Rheumatic and musculoskeletal pain
- Back ache

**FOR RELIEF FROM:**

- Cold and flu symptoms
- Period pain
- Rheumatic and musculoskeletal pain
- Back ache

**DO NOT TAKE IF YOU:**

- Have (or have had) two or more episodes of a stomach ulcer, perforation or bleeding
- Are allergic to ibuprofen, paracetamol or any other ingredient of the product, aspirin or other related

**WARNING:** Do not take more medicine than the label tells you to. You do not get better, talk to your doctor. Do not use this product if you have not first tried paracetamol or ibuprofen to relieve your pain. Keep out of the sight and reach of children.

**DO NOT TAKE IF YOU:**

- Have (or have had) two or more episodes of a stomach ulcer, perforation or bleeding
- Are allergic to ibuprofen, paracetamol or any other ingredient of the product, aspirin or other related

**FOR RELIEF FROM:**

- Cold and flu symptoms
- Period pain
- Rheumatic and musculoskeletal pain
- Back ache

**DO NOT TAKE IF YOU:**

- Have (or have had) two or more episodes of a stomach ulcer, perforation or bleeding
- Are allergic to ibuprofen, paracetamol or any other ingredient of the product, aspirin or other related

**WARNING:** Do not take more medicine than the label tells you to. You do not get better, talk to your doctor. Do not use this product if you have not first tried paracetamol or ibuprofen to relieve your pain. Keep out of the sight and reach of children.

**DO NOT TAKE IF YOU:**

- Have (or have had) two or more episodes of a stomach ulcer, perforation or bleeding
- Are allergic to ibuprofen, paracetamol or any other ingredient of the product, aspirin or other related

**FOR RELIEF FROM:**

- Cold and flu symptoms
- Period pain
- Rheumatic and musculoskeletal pain
- Back ache

**DO NOT TAKE IF YOU:**

- Have (or have had) two or more episodes of a stomach ulcer, perforation or bleeding
- Are allergic to ibuprofen, paracetamol or any other ingredient of the product, aspirin or other related

**WARNING:** Do not take more medicine than the label tells you to. You do not get better, talk to your doctor. Do not use this product if you have not first tried paracetamol or ibuprofen to relieve your pain. Keep out of the sight and reach of children.

**DO NOT TAKE IF YOU:**

- Have (or have had) two or more episodes of a stomach ulcer, perforation or bleeding
- Are allergic to ibuprofen, paracetamol or any other ingredient of the product, aspirin or other related

**FOR RELIEF FROM:**

- Cold and flu symptoms
- Period pain
- Rheumatic and musculoskeletal pain
- Back ache

**WHEN TO TAKE THIS PRODUCT:**

For the first day of treatment, try either ibuprofen or paracetamol individually in accordance with the product instructions. If the pain has not been relieved then the next day you can start taking this product.

**FOR ORAL ADMINISTRATION AND SHORT TERM USE ONLY:** Adults over 18 years old and elderly: take 1 tablet (or 2 if required), up to 3 times daily with food. Do not give to children under 18 years. If your symptoms persist for more than 3 days or worsen, consult your healthcare professional.

**DO NOT TAKE IF YOU:**

- Have (or have had) two or more episodes of a stomach ulcer, perforation or bleeding
- Are allergic to ibuprofen, paracetamol or any other ingredient of the product, aspirin or other related

**WARNING:** Do not take more medicine than the label tells you to. You do not get better, talk to your doctor. Do not use this product if you have not first tried paracetamol or ibuprofen to relieve your pain. Keep out of the sight and reach of children.

**DO NOT TAKE IF YOU:**

- Have (or have had) two or more episodes of a stomach ulcer, perforation or bleeding
- Are allergic to ibuprofen, paracetamol or any other ingredient of the product, aspirin or other related

**FOR RELIEF FROM:**

- Cold and flu symptoms
- Period pain
- Rheumatic and musculoskeletal pain
- Back ache

**SPÉAK TO A PHARMACIST/DOCTOR BEFORE TAKING IF YOU:**

- Have or have had asthma, diabetes, high cholesterol, high/low blood pressure, a stroke, heart, liver, kidney or bowel problems
- Are a smoker
- Are pregnant

**DO NOT TAKE IF YOU:**

- Have (or have had) two or more episodes of a stomach ulcer, perforation or bleeding
- Are allergic to ibuprofen, paracetamol or any other ingredient of the product, aspirin or other related

**WARNING:** Do not take more medicine than the label tells you to. You do not get better, talk to your doctor. Do not use this product if you have not first tried paracetamol or ibuprofen to relieve your pain. Keep out of the sight and reach of children.

**DO NOT TAKE IF YOU:**

- Have (or have had) two or more episodes of a stomach ulcer, perforation or bleeding
- Are allergic to ibuprofen, paracetamol or any other ingredient of the product, aspirin or other related

**FOR RELIEF FROM:**

- Cold and flu symptoms
- Period pain
- Rheumatic and musculoskeletal pain
- Back ache

**NUROMOL**  
DUAL ACTION PAIN RELIEF  
200mg/500mg Tablets

**For use only after trying ibuprofen or paracetamol first**



**NUROMOL**  
DUAL ACTION PAIN RELIEF  
200mg/500mg Tablets  
Ibuprofen & Paracetamol  
6 Tablets

Ink & varnish free area

**RE00000**

LOT:  
EXP:

Pharmacode Reads This Way  
First Bar  
Last Bar



TBC - English

Nuromol  
Dual Action  
Pain Relief  
Number 200 mg / Sign 500 mg  
Tablets

Trident Reference No: RB292404  
ZEN Ref: TR1307922

**CUSTOMER INFO:**  
Minimum Point Size = 7.00pt

N521

Reckitt Benckiser Healthcare  
UK Ltd., Slough SL1 3UH  
PL 00063/0649



**NUROMOL**  
DUAL ACTION PAIN RELIEF  
200mg/500mg Tablets

For use only after trying ibuprofen or paracetamol first



NUROMOL

DUAL ACTION PAIN RELIEF

200mg/500mg Tablets

Ibuprofen & Paracetamol

12 Tablets

LOT:      EXP:

RE00000

Ink & varnish free area

**CONTAINS PARACETAMOL.** Do not take anything else containing paracetamol while taking this medicine. Talk to a doctor at once if you take too much of this medicine, even if you feel well. This is because too much paracetamol can cause delayed, serious liver damage.

**DO NOT TAKE IF YOU:**

- Have (or have had) two or more episodes of a stomach ulcer, perforation or bleeding
- Are allergic to ibuprofen, paracetamol or any other ingredient of the product, aspirin or other related
- Headache
- Dental pain
- Cold and flu symptoms
- Rheumatic and
- Sore throat and fever
- Back ache
- Muscular pain
- Migraine

**FOR RELIEF FROM:**

- Cold and flu symptoms
- Rheumatic and
- Sore throat and fever
- Back ache
- Muscular pain
- Migraine

**DO NOT TAKE IF YOU:**

- Have (or have had) two or more episodes of a stomach ulcer, perforation or bleeding
- Are allergic to ibuprofen, paracetamol or any other ingredient of the product, aspirin or other related

**WARNING:** Do not take more medicine than the label tells you to. If you do not get better, talk to your doctor. Do not use this product if you have not first tried paracetamol or ibuprofen to relieve your pain. Keep out of the sight and reach of children.

**DO NOT TAKE IF YOU:**

- Have (or have had) two or more episodes of a stomach ulcer, perforation or bleeding
- Are allergic to ibuprofen, paracetamol or any other ingredient of the product, aspirin or other related

**DO NOT TAKE IF YOU:**

- Have (or have had) two or more episodes of a stomach ulcer, perforation or bleeding
- Are allergic to ibuprofen, paracetamol or any other ingredient of the product, aspirin or other related

**DO NOT TAKE IF YOU:**

- Have (or have had) two or more episodes of a stomach ulcer, perforation or bleeding
- Are allergic to ibuprofen, paracetamol or any other ingredient of the product, aspirin or other related

**DO NOT TAKE IF YOU:**

- Have (or have had) two or more episodes of a stomach ulcer, perforation or bleeding
- Are allergic to ibuprofen, paracetamol or any other ingredient of the product, aspirin or other related

Pharmacode Reads This Way

First Bar

Last Bar





