



10 May 2021

Year: 2021 Week: 18

In This Issue:

- Key Messages.
- Weekly summary.
- Total contacts.
- Syndromic indicators.
- Notes and caveats.
- Further information.
- Acknowledgements.

Syndromic indicators at a glance:

Number of contacts and percentage of Read coded contacts.

1: Total out-of-hours contacts:

Daily total number of out-of-hours and unscheduled contacts and 7-day moving average (adjusted for bank holidays).

Key messages


Data to: 09 May 2021

During week 18 GP out of hours contacts for acute respiratory infection continued to increase across all age groups but noticeably in those aged 5-14 years (figures 2 and 2a). Contacts for difficulty breathing/wheeze/asthma increased in children aged 5-14 years (figure 5a).

Note: during the COVID-19 pandemic, patients with COVID-19 symptoms are generally advised to initially access a COVID-19 test through the national COVID-19 testing programme. This is likely to result in lower numbers of patients accessing health advice as monitored through syndromic surveillance systems. Syndromic data should therefore be interpreted with some caution and in the context of other COVID-19 monitoring data sources.

Key indicator	No. of contacts	% Week 18	% Week 17	Trend*
All OOH contacts, all causes	214,033			
Acute respiratory infection	6,320	7.99	6.55	↑
Influenza-like illness	79	0.10	0.08	↔
Bronchitis/bronchiolitis	69	0.09	0.06	↑
Difficulty breathing/wheeze/asthma	1,334	1.69	1.63	↔
Pharyngitis	41	0.05	0.04	↔
Gastroenteritis	1,999	2.53	2.32	↔
Diarrhoea	664	0.84	0.78	↔
Vomiting	784	0.99	0.99	↔
Chest pain/myocardial infarction	870	1.10	1.08	↔

*Trend: reports on the trend seen over previous weeks in the percentage of Read coded contacts.




10 May 2021

Year: 2021 Week: 18


2: Acute Respiratory Infection daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7-day moving average*.



2a: Acute Respiratory Infection by age group.

As a percentage of total contacts within each age group, shown as a 7-day moving average*.



Intentionally left blank


*7-day moving average adjusted for bank holidays.

10 May 2021

Year: 2021 Week: 18


3: Influenza-like illness daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7-day moving average*.



3a: Influenza-like illness by age group.

As a percentage of total contacts within each age group, shown as a 7-day moving average*.



Intentionally left blank


*7-day moving average adjusted for bank holidays.

10 May 2021

Year: 2021 Week: 18


**4: Bronchitis/
bronchiolitis daily
contacts.**

Shown as a percentage of the total contacts with a Read code and as a 7-day moving average*.




**4a: Bronchitis/
bronchiolitis daily
contacts by age
group*.**

As a percentage of total contacts within each age group, shown as a 7-day moving average*.



**5: Difficulty breathing/
wheeze/asthma daily
contacts.**

Shown as a percentage of the total contacts with a Read code and as a 7-day moving average*.




*7-day moving average
adjusted for bank
holidays.

10 May 2021

Year: 2021 Week: 18


5a: Difficulty breathing/wheeze/asthma daily contacts by age group*.

As a percentage of total contacts within each age group, shown as a 7-day moving average*.




6: Acute pharyngitis and persistent sore throat.

Shown as a percentage of the total contacts with a Read code and as a 7-day moving average*.



7: Gastroenteritis daily contacts

Shown as a percentage of the total contacts with a Read code and as a 7-day moving average*.




*7-day moving average adjusted for bank holidays.

10 May 2021

Year: 2021 Week: 18


8: Diarrhoea daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7-day moving average*.




8a: Diarrhoea daily contacts by age group*.

As a percentage of total contacts within each age group, shown as a 7-day moving average*.



9: Vomiting daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7-day moving average*.




*7-day moving average adjusted for bank holidays.

10 May 2021

Year: 2021 Week: 18


9a: Vomiting daily contacts by age group*.

As a percentage of total contacts within each age group, shown as a 7-day moving average*.



10: Chest pain/myocardial infarction daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7-day moving average*.



Intentionally left blank

*7-day moving average adjusted for bank holidays.

10 May 2021

Year: 2021 Week: 18

Notes and caveats:

- This bulletin presents data from the Public Health England (PHE) GP Out-of-hours\Unscheduled Care Surveillance System (GP OOHSS).
- Fully anonymised data from GP out-of-hours (OOH) and unscheduled care service providers in England are being transferred to PHE for analysis and interpretation by the PHE Real-time Syndromic Surveillance Team (ReSST).
- This system supplements existing PHE syndromic surveillance systems by monitoring data on GP consultations outside of routine surgery opening times (evenings, weekends and bank holidays) and unplanned contacts within NHS primary care.
- The key indicators presented within this bulletin are derived by grouping selected Read coded consultations.
- GP OOH consultation data are analysed on a daily basis to identify national and regional trends. A statistical algorithm underpins each system, routinely identifying activity that has increased significantly or is statistically significantly high for the time of year. Results from these daily analyses are assessed by the ReSST, along with analysis by age group, and anything deemed of public health importance is alerted by the team.
- Baselines represent seasonally expected levels of activity and are constructed from historical data since Nov 2009. They take into account any known substantial changes in data collection, population coverage or reporting practices. Gastroenteritis, diarrhoea and vomiting baselines also account for changes since the introduction of rotavirus vaccine in July 2013. Baselines are refreshed using the latest data on a regular basis however they currently exclude data from 2020 due to the COVID-19 pandemic affecting GP services and patient health care seeking behaviour.

Moving Epidemic Method (MEM):

- During winter we present Moving Epidemic Method (MEM) influenza thresholds on selected indicators. MEM is a standard methodology used for setting influenza thresholds across many European nations.¹
- MEM is used for GP OOH ILI thresholds at a national level.
- **MEM thresholds should be interpreted using 7 day moving averages rather than daily data.**
- MEM thresholds currently use six years of historic data (2013-2019). The thresholds are re-calculated every year.
- Baseline ('Pre-epidemic') thresholds are used alongside other surveillance systems to identify the start of influenza circulating in the community
- 40%, 95% and 97.5% intensity thresholds are used to identify when influenza activity moves from low to medium, high or very high.

¹Vega T et al. Influenza Other Respir Viruses. 2013;7(4):546-58.

Further information:

The GP Out-of-Hours Surveillance System Bulletin can also be downloaded from the PHE Real-time Syndromic Surveillance website which also contains more information about syndromic surveillance:

<https://www.gov.uk/government/collections/syndromic-surveillance-systems-and-analyses>

Acknowledgements:

We are grateful to Advanced and the GP OOH and unscheduled care service providers who have kindly agreed to participate in this system.

Contact ReSST:

syndromic_surveillance@phe.gov.uk

PHE Out-of-Hours/Unscheduled Care Surveillance

Produced by: PHE Real-time Syndromic Surveillance Team
1st Floor, 5 St Philips Place, Birmingham, B3 2PW

Tel: 0344 225 3560 > Option 4 > Option 2

Fax: 0121 236 2215

Web: <https://www.gov.uk/government/collections/syndromic-surveillance-systems-and-analyses>