

SPI-M-O

Medium-Term Projections

12th October 2021

SPI-M-O Medium-term Projections

- **These projections are not forecasts or predictions. They represent a scenario in which the trajectory of the epidemic continues to follow the trends that were seen in the data up to 11th October.**
- The delay between infection, developing symptoms, the need for hospital care, and death means they cannot fully reflect the impact of policy and behavioural changes made in the two to three weeks prior to 11th October.
- Predicting the peak of the epidemic in a particular nation or region is difficult and prone to large levels of uncertainty. Any changes in behaviour will impact transmission and alter the trends shown in the projections.
- **The projections do not include the effects of any future policy or behavioural changes.** The effect of school opening and closing has been included.
- **The projections include the impact of vaccines given over the next three weeks. This has been based on a rollout scenario provided by Cabinet Office for modelling purposes.** The rollout scenario assumes booster doses are administered according to [JCVI's advice](#). The scenario also includes the vaccination programme for 12- to 15-year olds. The continued rollout of doses will have a limited impact over the next three weeks, given lags between vaccination and protection, and between infection and hospital admission.
- Modelling groups have used their expert judgement and evidence from [Public Health England](#), [Scottish Universities & Public Health Scotland](#), and other published efficacy studies when making assumptions about vaccine effectiveness. A table summarising these assumptions is available in the annex.
- Not all modelling groups produce projections for both hospitalisations and deaths, so there will be some differences between the models included in the combined projections for each metric.

Metrics:

- **New hospitalisations per day:** Number of individuals admitted with COVID-19 and inpatients newly diagnosed with COVID-19. Data definitions differ slightly across all four nations.
- **New deaths per day (by date of death):** The number of COVID-19 deaths within 28 days of a positive test. Data definitions differ slightly across all four nations.

Modelled projections based on trends to 11th October 2021


New hospital admissions per day

These projections are based on current trends and will not fully reflect the impact of policy or behavioural changes over the past two to three weeks. They are not forecasts or predictions.


Key

- Real data
- Expected to Increase
- Projection Midpoint
- High and low estimates 5th to 95th percentile
- High and low estimates 25th to 75th percentile
- Models


ENGLAND




WALES



SCOTLAND



NORTHERN IRELAND



The fan charts show the **interquartile range** of the combined projections based on current trends for England and Scotland. For Wales and Northern Ireland, both the **90% confidence interval and interquartile range** is shown. Please note, not all the y axes start from zero on these plots.

The delay between infection, developing symptoms, the need for hospital care, and death means they cannot fully reflect the impact of policy or behavioural changes in the two to three weeks prior to 11th October. Predicting the peak of the epidemic in a particular nation or region is difficult and prone to large levels of uncertainty. Changes in behaviour will impact transmission and alter the trends shown in the projections. **The projections do not include the effects of any future policy or behavioural changes.**

These projections include the potential impact of vaccines to be given over the next three weeks. This has been based on a rollout scenario provided by Cabinet Office for modelling purposes. The rollout scenario assumes booster doses are administered according to [JCVI's advice](#). The scenario also includes the recently announced vaccination programme for 12- to 15-year olds. The continued rollout of doses will have a limited impact over the next three weeks, given lags between vaccination and protection, and between infection and hospital admission.

Data notes:

England: Number of patients admitted with confirmed COVID-19 and the number of inpatients diagnosed with COVID-19 in the past 24 hours. Taken from NHSE COVID-19 Situation reports.

Wales: Number of patients admitted with confirmed COVID-19 and inpatients diagnosed with COVID-19. Provided by Public Health Wales.


Scotland: Number of patients who tested positive for COVID-19 in the 14 days prior to admission, on the day of admission, or during their stay in hospital. Readmissions within 14 days of a positive test are excluded. Provided by Public Health Scotland.

Northern Ireland: Number of patients admitted with confirmed COVID-19 and inpatients diagnosed with COVID-19. Provided by Health and Social Care Northern Ireland.


Modelled projections based on trends to 11th October 2021

New hospital admissions per day


These projections are based on current trends and will not fully reflect the impact of policy or behavioural changes over the past two to three weeks. They are not forecasts or predictions.




EAST OF ENGLAND




LONDON




MIDLANDS




NORTH EAST AND YORKSHIRE




NORTH WEST



SOUTH EAST



SOUTH WEST



The fan charts show the **interquartile range** of the combined projections based on current trends. Please note, not all the y axes start from zero on these plots.

The delay between infection, developing symptoms, the need for hospital care, and death means they cannot fully reflect the impact of policy or behavioural changes in the two to three weeks prior to 11th October. Predicting the peak of the epidemic in a particular nation or region is difficult and prone to large levels of uncertainty. Changes in behaviour will impact transmission and alter the trends shown in the projections. **The projections do not include the effects of any future policy or behavioural changes.**

These projections include the potential impact of vaccines to be given over the next three weeks. This has been based on a rollout scenario provided by Cabinet Office for modelling purposes. The rollout scenario assumes booster doses are administered according to [JCVI's advice](#). The scenario also includes the recently announced vaccination programme for 12- to 15-year olds. The continued rollout of doses will have a limited impact over the next three weeks, given lags between vaccination and protection, and between infection and hospital admission.

Data notes:

England: Number of patients admitted with confirmed COVID-19 and the number of inpatients diagnosed with COVID-19 in the past 24 hours. The past data is taken from the NHS England COVID-19 Sitrep's.



Modelled projections based on trends to 11th October 2021

New deaths per day

These projections are based on current trends and will not fully reflect the impact of policy or behavioural changes over the past two to three weeks. They are not forecasts or predictions.

Key

- • • Real data
- • • Expected to Increase
- Projection Midpoint
- High and low estimates 25th to 75th percentile
- Models



The fan charts show the **interquartile range** of the combined projections for England and Scotland based on current trends.

The delay between infection, developing symptoms, the need for hospital care, and death means they cannot fully reflect the impact of policy or behavioural changes in the two to three weeks prior to 11th October. Predicting the peak of the epidemic in a particular nation or region is difficult and prone to large levels of uncertainty. Changes in behaviour will impact transmission and alter the trends shown in the projections. **The projections do not include the effects of any future policy or behavioural changes.**

These projections include the potential impact of vaccines to be given over the next three weeks. This has been based on a rollout scenario provided by Cabinet Office for modelling purposes. The rollout scenario assumes booster doses are administered according to [JCVI's advice](#). The scenario also includes the recently announced vaccination programme for 12- to 15-year olds. The continued rollout of doses will have a limited impact over the next three weeks, given lags between vaccination and protection, and between infection and hospital admission.

Due to an insufficient number of projections, it has not been possible to produce a reliable combination for the number of deaths in Wales and Northern Ireland. SPI-M-O's consensus view is that the number of deaths will remain broadly flat in these regions.

Data Notes:

The number of COVID-19 deaths (by date of death) within 28 days of a positive test.

The past data for England is taken from the PHE line list of deaths. The past data for Scotland, Wales, and Northern Ireland is taken from the Coronavirus (COVID-19) in the UK dashboard on Gov.uk.

Modelled projections based on trends to 11th October 2021


New deaths per day

These projections are based on current trends and will not fully reflect the impact of policy or behavioural changes over the past two to three weeks. They are not forecasts or predictions.


Key

- Real data
- Expected to Increase
- Projection Midpoint
- High and low estimates 5th to 95th percentile
- High and low estimates 25th to 75th percentile
- Models


EAST OF ENGLAND




LONDON




MIDLANDS




NORTH EAST AND YORKSHIRE




NORTH WEST



SOUTH EAST



SOUTH WEST



The fan charts show the **interquartile range** of the combined projections based on current trends for the East of England and South West. For all other regions, both the **90% confidence interval** and **interquartile range** is shown. Please note, not all the y axes start from zero on these plots.

The delay between infection, developing symptoms, the need for hospital care, and death means they cannot fully reflect the impact of policy or behavioural changes in the two to three weeks prior to 11th October. Predicting the peak of the epidemic in a particular nation or region is difficult and prone to large levels of uncertainty. Changes in behaviour will impact transmission and alter the trends shown in the projections. **The projections do not include the effects of any future policy or behavioural changes.**

These projections include the potential impact of vaccines to be given over the next three weeks. This has been based on a rollout scenario provided by Cabinet Office for modelling purposes. The rollout scenario assumes booster doses are administered according to [JCVI's advice](#). The scenario also includes the recently announced vaccination programme for 12- to 15-year olds. The continued rollout of doses will have a limited impact over the next three weeks, given lags between vaccination and protection, and between infection and hospital admission.

Data Notes:

The number of COVID-19 deaths (by date of death) within 28 days of a positive test. The past data for England is taken from the PHE line list of deaths.

Annex: SPI-M-O Vaccine Effectiveness Assumptions

The LSHTM EpiNow model has also been included in this week's combined projections. This model projects forward based on the recent trends in the data and doesn't explicitly include the impact of vaccination. However, the protection provided by vaccinations given to date will be reflected in the data and therefore implicitly included in the projections produced by the model.

		Manchester [3]	Warwick [2,4] (Death)	Warwick [2,4] (Hospitalisation)	PHE/ Cambridge [2]	Scottish Government [2]
Pfizer-BioNTech	1 Dose	75%	80%	80%	78%	80%
	2 Doses	75%	98%	95%	97%	95%
Oxford-AstraZeneca	1 Dose	75%	80%	80%	78%	80%
	2 Doses	75%	98%	95%	97%	95%
Moderna	1 Dose	75%	80%	80%	78%	80%
	2 Doses	75%	98%	95%	97%	95%

		Manchester [3]	Warwick [2,4]	PHE/ Cambridge [2]	Scottish Government [2]
Pfizer-BioNTech	1 Dose	75%	55%	31%	55%
	2 Doses	75%	85%	80%	75%
Oxford-AstraZeneca	1 Dose	75%	45%	31%	40%
	2 Doses	75%	70%	80%	65%
Moderna	1 Dose	75%	55%	31%	75%
	2 Doses	75%	85%	80%	85%

		Manchester [3,5]	Warwick [2,4]	PHE/ Cambridge [2,5]	Scottish Government [2]
Pfizer-BioNTech	1 Dose	-	30%	-	29%
	2 Doses	-	30%	-	40%
Oxford-AstraZeneca	1 Dose	-	30%	-	37%
	2 Doses	-	30%	-	44%
Moderna	1 Dose	-	30%	-	29%
	2 Doses	-	30%	-	40%

[1] The assumed delay between vaccination and protection varies between 10 and 21 days for dose 1 and between 7 and 21 days for dose 2 across the modelling groups.

[2] Warwick, PHE/Cambridge & Scottish Government's vaccine effectiveness assumptions are for the B.1.617.2 (delta) variant.

[3] Manchester's model does not split vaccine effectiveness by vaccine type or different doses.

[4] Warwick's model considers a range of scenarios for the partial waning of vaccine effectiveness. The results from these scenarios are then combined to form their projection.

[5] The Manchester and PHE/ Cambridge models do not include a reduction in the risk of onwards transmission after receiving either vaccine.