


Department for Environment, Food and Rural Affairs

The Expert Committee on Pesticide Residues in Food (PRiF)

# Report on the pesticide residues monitoring programme: Results of Quarter 4 2021





© Crown copyright 2021

You may re-use this information (excluding logos) free of charge in any format or medium, under the terms of the Open Government Licence v.3. To view this licence, visit [The National Archives](#) or email [PSI@nationalarchives.gov.uk](mailto:PSI@nationalarchives.gov.uk)

This publication is available at [Expert Committee on Pesticide Residues in Food](#)

For further details on information contained in this report, previous surveys or information concerning pesticide residues in food, please contact:

Expert Committee on Pesticide Residues in Food  
HSE's Chemicals Regulation Division  
Mallard House  
Kings Pool  
Peasholme Green  
York  
YO1 7PX

[prif@hse.gov.uk](mailto:prif@hse.gov.uk)

## Contents

Summary: Quarter 4 2021 findings	5
Table 1: Headlines with links to detailed information	7
Introduction to the work of the Expert Committee on Pesticide Residues in Food (PRiF)	11
UK National Monitoring Programmes	11
HSE assessment of risk	12
Table 2: 2021 Survey Design	14
Table 3: Summary of Results	16
Table 4: Summary of samples sent to FSA for follow up	20
Table 5: Summary of MRL Exceedances	22
Section 1: findings by food	27
Asparagus (GB)	27
Asparagus (NI)	28
Aubergine (GB)	29
Aubergine (NI)	30
Banana (GB)	31
Banana (NI)	34
Beans with pods (GB)	36
Beans with pods (NI)	39
Beef (GB)	40
Beef (NI)	41
Berries and small fruits (GB)	42
Berries and small fruits (NI)	44
Bread (GB)	46
Bread (NI)	48
Broccoli (GB)	50
Broccoli (NI)	52
Cheese (soft) (GB)	53
Cheese (soft) (NI)	55
Eggs (GB)	57
Eggs (NI)	58
Fish (white) (NI)	59
Grapefruit (GB)	61
Grapefruit (NI)	63
Grapes (GB)	65
Grapes (NI)	67
Infant food (cereal based) (GB)	68
Infant food (cereal based) (NI)	71
Melon (GB)	72
Melon (NI)	74
Milk (GB)	76
Milk (NI)	77
Mushroom (GB)	78
Mushroom (NI)	80
Nuts (GB)	82
Olive oil (GB)	84
Olive oil (NI)	86
Peppers (GB)	87
Peppers (NI)	88
Peppers (processed) (GB)	89
Potatoes (GB)	91
Potatoes (NI)	92
Raspberry (GB)	93

Raspberry (NI)	95
Spring greens and kale (GB)	97
Spring greens and kale (NI)	99
Wheat (GB)	101
Wheat (NI)	103
Section 2: Sample details and supplier responses	105
Sample details	105
Supplier responses	107
Section 3: HSE assessment of risk	108
Risk Assessment - dietary intake assessments	116
Section 4: issues arising in this report and updates on previous reports	138
Issues arising in this report	138
Follow-up from Previous Reports	143
Brand name details of samples where follow-up action is now complete	144
In our next report:	146
Section 5: background and reference	147
Glossary	155

# Summary: Quarter 4 2021 findings

## Chair's comments

During this year's surveillance programme, we are measuring up to 397 different pesticides in each of the foods we survey. The Quarter 4 programmes surveyed 1,085 samples of 25 different foods (see [contents page](#) for a full list). The majority of samples were collected between the beginning of October and the end of December 2021.

Of 1,085 samples, we found residues in 581 of them and of these 41 samples contained residues over the Maximum Residue Level (MRL). Most of the residues detected did not cause health concern.

HSE undertakes screening and detailed risk assessments, as required, for the pesticide residues found. This is to determine whether the residues present could lead to someone eating an amount above a level that is considered safe. HSE also produces [detailed risk assessments](#) for every case where the actual residue level found could lead to an intake above the safety levels.

We needed to consider the potential short-term health effects of some of the residues found in more detail. In most of these cases, we concluded that effects on health were either unlikely or not expected. Full details are presented in [section 3](#).

We also needed to consider the potential genotoxic health effects of some of the residues of pesticides not authorised in the UK but found in some imported food. We also have considered the potential genotoxic effects of residues arising from a pesticide which is currently authorised in the UK. We concluded that at the levels present, a risk of an adverse effect on health due to genotoxicity would be low.

These detailed considerations as well as links to underlying information are covered in our reports for banana/plantain, beans with pods, broccoli, grapefruit, grapes, infant food, melon, olive oil, processed peppers, spring greens and kale.

None of the individual commodity long-term exposure screening assessments performed in this quarter (for each of the pesticides found in this report) indicated any potential for adverse long-term health effects. This was based on the assessment of dietary intakes as below the ADI or other established long-term health-based reference values.

Full details of suppliers and retailers of the food sampled, and full analytical results, are available on [data.gov.uk](#) as ODS (Open Document Spreadsheet) files. We hope this data format is useful for people wanting to look at the individual results in more detail.

These samples were collected after the UK had left the EU, so this is the fourth report where we have reported the results for samples collected in Great Britain (GB) separately from those collected in Northern Ireland (NI). Surveys have been titled throughout the report as either GB or NI to make clear where the samples were collected. Samples collected in GB are subject to GB MRLs. GB MRLs are set by inclusion in a new statutory Register, implemented and updated by means of a database<sup>1</sup>. For samples collected in NI, certain aspects of EU food law, including compliance with EU set MRLs, continue to apply

---

<sup>1</sup> [GB MRL Register \(pesticides.gov.uk\)](#)

under the terms of the Northern Ireland Protocol. In the detailed data files HSE is, for 2021 results, still separating out EU from non-EU origin foods in the results.

We asked suppliers and the authorities of the exporting countries for an explanation of our findings. Any responses we have received specifically for publication are available in [Section 2](#) sample details and supplier responses.

If you have any feedback or comments on the monitoring programme or the reports produced please send them to our secretariat at [prif@hse.gov.uk](mailto:prif@hse.gov.uk).

Ann Davison  
Chair of the Expert Committee on Pesticide Residues in Food

## Consumer risk summary

HSE screens each residue detected for any consumer health issues to identify which need to be considered in more detail. We comment on any risks HSE considered in detail in our full report, and [HSE's risk assessments](#) are also published.

HSE liaises with the Food Standards Agency (FSA) on consumer risk assessment and the FSA also takes part in our meetings.

### Table 1: Headlines with links to detailed information

Survey title (where samples collected)	Number of pesticides sought	Samples tested	Detailed risk assessment presented?	MRL exceedances (samples)
<a href="#">All Foods tested</a>				
<a href="#">Asparagus (GB)</a>	390	24		0
<a href="#">Asparagus (NI)</a>	389	6		0
<a href="#">Aubergine (GB)</a>	373	25		2
<a href="#">Aubergine (NI)</a>	373	11		1
<a href="#">Banana (GB)</a>	393	13	Yes	2
<a href="#">Banana (NI)</a>	392	13	Yes	0
<a href="#">Beans with pods (GB)</a>	390	32	Yes	5
<a href="#">Beans with pods (NI)</a>	389	8		1
<a href="#">Beef (GB)</a>	116	24		0
<a href="#">Beef (NI)</a>	38	16		1
<a href="#">Berries and small fruits (GB)</a>	368	31		4
<a href="#">Berries and small fruits (NI)</a>	368	10		0
<a href="#">Bread (GB)</a>	393	54		1
<a href="#">Bread (NI)</a>	392	20		0
<a href="#">Broccoli (GB)</a>	370	31	Yes	0
<a href="#">Broccoli (NI)</a>	370	10		0
<a href="#">Cheese (soft) (GB)</a>	113	48		2
<a href="#">Cheese (soft) (NI)</a>	38	15		0
<a href="#">Eggs (GB)</a>	116	24		0
<a href="#">Eggs (NI)</a>	117	14		0
<a href="#">Fish (white) (NI)</a>	38	16		0
<a href="#">Grapefruit (GB)</a>	391	24	Yes	0

Survey title (where samples collected)	Number of pesticides sought	Samples tested	Detailed risk assessment presented?	MRL exceedances (samples)
<a href="#">Grapefruit (NI)</a>	390	14	Yes	1
<a href="#">Grapes (GB)</a>	397	18	Yes	0
<a href="#">Grapes (NI)</a>	396	12		0
<a href="#">Infant food (cereal based) (GB)</a>	397	36	Yes	13
<a href="#">Infant food (cereal based) (NI)</a>	395	10		0
<a href="#">Melon (GB)</a>	372	30	Yes	0
<a href="#">Melon (NI)</a>	372	11		0
<a href="#">Milk (GB)</a>	113	78		0
<a href="#">Milk (NI)</a>	114	16		0
<a href="#">Mushroom (GB)</a>	396	18		0
<a href="#">Mushroom (NI)</a>	395	14		1
<a href="#">Nuts (GB)</a>	388	48		3
<a href="#">Olive oil (GB)</a>	392	18	Yes	0
<a href="#">Olive oil (NI)</a>	390	3		0
<a href="#">Peppers (GB)</a>	395	30		0
<a href="#">Peppers (NI)</a>	394	14		0
<a href="#">Peppers (processed) (GB)</a>	391	24	Yes	2
<a href="#">Potatoes (GB)</a>	390	61		0
<a href="#">Potatoes (NI)</a>	390	14		0
<a href="#">Raspberry (GB)</a>	368	29		0
<a href="#">Raspberry (NI)</a>	368	10		0
<a href="#">Spring greens and kale (GB)</a>	366	30	Yes	1
<a href="#">Spring greens and kale (NI)</a>	366	12	Yes	1
<a href="#">Wheat (GB)</a>	396	36		0
<a href="#">Wheat (NI)</a>	395	30		0

Samples collected in GB must comply with GB set MRLs unless the goods are qualifying Northern Ireland goods and are subject to unfettered access under the terms of the UK Internal Market Act 2020. For samples collected in NI, under the Northern Ireland Protocol, certain aspects of EU food law, including compliance with EU MRLs apply.



## Other issues

### Suspected unauthorised uses

HSE passed details to the enforcement team of samples grown in GB or NI that contained a residue which does not have a plant protection product (PPP) with that active authorised for use on that crop, in GB and NI.

- 2 GB blackberries cyantraniliprole.
- 1 GB broccoli with triallate. HSE Enforcement found no evidence of misuse. These findings have been referred to the HSE regulatory division for further follow up with approval holders.
- 2 GB broccoli with triallate.
- 1 NI mushroom with deltamethrin
- 1 GB potato with 1,4-dimethylnaphthalene. HSE Enforcement found no evidence of misuse and spray records supplied to HSE support that, 1,4-dimethylnaphthalene was not used. HSE's investigation concluded that contaminated storage crates may have been the source of the residue.
- 1 GB potato with imazalil. HSE Enforcement found no evidence of misuse. This sample was removed from the survey as it was later identified as not intended for human consumption.
- 1 GB raspberries with thiacloprid.
- 2 GB raspberries with thiacloprid from samples arising from the same grower. HSE Enforcement concluded that thiacloprid had been used legally while the active was still authorised on raspberries.
- 9 GB spring greens and kale with triallate.
- 2 GB spring greens and kale with triallate. HSE Enforcement found no evidence of misuse and spray records showed that triallate was not used.
- 2 GB spring greens and kale with triallate and prosulfocarb.
- 1 NI spring greens and kale with triallate.
- 1 NI spring greens and kale with triallate. HSE Enforcement found no evidence of misuse and spray records showed that triallate was not used.
- 1 NI spring greens and kale with triallate and prosulfocarb.

### Organic samples with residues

HSE writes to the suppliers of samples of organic produce if they contain a pesticide residue. Defra's Organic Farming branch and the organic certification organisation are also informed.

- 1 GB sample of infant food (cereal based)
- 1 NI sample of mushrooms
- 1 GB sample of peppers
- 1 NI sample of peppers
- 1 NI sample of potatoes

## Further information

Further information on the individual sample details is in an accessible format at [Pesticide Residues in Food Quarterly Data](#).

This includes:

- brand name, sampling point and origin information
- pesticides sought and residues found
- HSE detailed risk assessments

## Introduction to the work of the Expert Committee on Pesticide Residues in Food (PRiF)

The PRiF's role is to give Ministers, the Director of the Health and Safety Executive (HSE) and the Chief Executive of the Food Standards Agency (FSA) independent advice on the UK government's national rolling programme of surveys, in particular:

- the planning of surveillance programmes for pesticide residues in the UK food supply and the evaluation of the results.
- procedures for sampling, sample processing, new methods of analysis, the assessment of variability of pesticide residues in food and related issues.

The Expert Committee on Pesticide Residues in Food was established in 2011. Our members have a broad range of expertise relating to the food supply industry. The main function of the Committee is to oversee Government's £2 million pesticide residues surveillance programme. Previously this work was carried out by the Pesticide Residues Committee.

Our Chair, Ann Davison has worked in consumer affairs for most of her career, running consumer organisations and networks. The committee also includes members with expertise in food science, food production and supply as well as two public interest experts.

Information on the membership of the PRiF is also available on the PRiF's website: [Expert Committee on Pesticide Residues in Food](#)

## UK National Monitoring Programmes

HSE, working under Defra, and the Scottish and Welsh governments authority, has official responsibility to organise a monitoring programme of GB food for pesticide residues. Similarly, HSE working under the Department of Agriculture, Environment and Rural affairs authority has official responsibility to organise a monitoring programme of NI food for pesticide residues, including participating in the EU multi-annual control programme.

The programmes are made up of a risk-based rolling programme of surveys and statutory programmes required by GB or EU law. It is a surveillance programme, which is designed based upon evidence gathered in the previous years, including previous results, PRiF advice and border control information. It is not an enforcement programme and its design is generally not adjusted during the year. HSE is also responsible for considering the safety of people who eat the food (in co-operation with the Food Standards Agency if necessary) and for following up adverse or unexpected results. HSE are also responsible for determining whether food is compliant with the law, specifically, whether any pesticide residue found is within the Maximum Residue Level.

Maximum Residue Levels (MRLs) reflect levels of pesticides that could occur in food which has been treated in accordance with good agricultural practice. Where pesticides do not give rise to readily detectable residues, or are not authorised for use on particular commodities, MRLs are set at the lowest level which can be identified in routine laboratory analysis. This provides a mechanism for statutory controls on pesticides in food which is put into circulation and for monitoring the correct use of these chemicals.

## HSE assessment of risk

HSE conducts a screening assessment of all the residues we find in the PRiF programme. If screening identifies any dietary intakes exceeding the relevant health-based reference values, then we present more detailed risk assessments, to consider whether there are any implications for health. Detailed risk assessments, where needed, are presented in [section 3](#). If we understand that a pesticide residue has a risk of genotoxicity (has potential to cause damage to genetic material), we will include this in the commentary.

Pesticide dietary intakes are assessed using models that combine data on the levels of residues in food with food dietary consumption values. If intakes are within the health-based reference values, then taking account of the precautions built into the model assessments we conclude that an effect on health is not anticipated. If dietary intakes exceed the reference values this does not automatically mean there are expected adverse health effects. However, this acts as a 'trigger' for HSE to consider these cases more thoroughly.

HSE conducts both short-term (acute) and long-term (chronic) assessments based on the residues found in the PRiF surveys. Each of these is tailored accordingly. Further information on the nature of HSE's assessments and approach is provided in the bullet points below, and in more detail, with reference to international assessment contexts in [section 3](#) and on HSE's website (<https://www.hse.gov.uk/pesticides/pesticides-registration/data-requirements-handbook/consumer-exposure.htm>)

- For acute assessment, we use short-term estimation values that use the highest residue found in a commodity and short-term consumption values for calculating short-term dietary intakes. These are then compared to the ARfD, a suitable health-based reference value for effects that could be caused by a single day or one-off consumption of a higher than usual residue. For acute assessment we consider the variation in residues that could occur within a residue sample, and a variability (multiplication) factor is included for that purpose, in order to address exposure to a higher than usual residue in a single item, such as a single apple or potato.
- For chronic assessment, we use long-term estimation values (based on median residues and long-term consumption values for calculating long-term dietary intakes) for each commodity survey and compare to the ADI, a suitable health-based reference value for lifetime. The issue is more fully considered in regulatory contexts pre-authorisation and at the time of MRL review. Then the issue is considered across all commodities (so more precautionary) by pesticide levels determined in GAP compliant trials, intended to address highest likely residues that might arise following pesticide use according to label recommendations.

For fruit and vegetables that have peel or skin that might not be consumed we present alternative risk assessments for 'without peel -flesh only' where peel versus pulp residue distribution data is available. As standard, we present a 'worst case' assessment for when all of the peel is consumed with the fruit.

- We calculate dietary intakes for different consumer groups, from infants, toddlers, and children of varying age, to adults, elderly, and vegetarians, to take account of

people with low bodyweights and varying dietary habits. As such the assessments we perform are protective for all consumers.

- For multiple residues, we consider the 'cocktail effect' – the possible implications to health of more than one pesticide being found in samples. We currently focus in detail on selected groups that we think are a priority to consider based on toxicity considerations and prevalence.

## Table 2: 2021 Survey Design

### Fruit and vegetables

Food	Sampling points	Sampled during	Reporting
Asparagus (GB)	Retail Outlets and Supply chain	Quarters 2 and 4	Quarters 2 and 4
Asparagus (NI)	Retail Outlets	Quarters 2 and 4	Quarters 2 and 4
Aubergine (GB)	Retail Outlets and Supply chain	Quarterly	Quarterly
Aubergine (NI)	Retail Outlets	Quarterly	Quarterly
Banana (GB)	Retail Outlets and Supply chain	Quarterly	Quarterly
Banana (NI)	Retail Outlets	Quarterly	Quarterly
Beans with pods (GB)	Retail Outlets and Supply chain	Quarterly	Rolling and Quarterly
Beans with pods (NI)	Retail Outlets	Quarterly	Rolling and Quarterly
Berries and small fruit (GB)	Retail Outlets and Supply chain	Quarterly	Quarterly
Berries and small fruit (NI)	Retail Outlets	Quarterly	Quarterly
Broccoli (GB)	Retail Outlets and Supply chain	Quarterly	Quarterly
Broccoli (NI)	Retail Outlets	Quarterly	Quarterly
Grapefruit (GB)	Retail Outlets and Supply chain	Quarter 2 - 4	Quarter 2 - 4
Grapefruit (NI)	Retail Outlets	Quarter 2 - 4	Quarter 2 - 4
Grapes (GB)	Retail Outlets and Supply chain	Quarterly	Rolling and Quarterly
Grapes (NI)	Retail Outlets	Quarterly	Rolling and Quarterly
Melon (GB)	Retail Outlets and Supply chain	Quarterly	Quarterly
Melon (NI)	Retail Outlets	Quarterly	Quarterly
Mushrooms (GB)	Retail Outlets	Quarterly	Quarterly
Mushrooms (NI)	Retail Outlets	Quarterly	Quarterly
Peppers (GB)	Retail Outlets and Supply chain	Quarterly	Quarterly
Peppers (NI)	Retail Outlets	Quarterly	Quarterly
Potatoes (GB)	Supply chain	Quarterly	Rolling and Quarterly
Potatoes (NI)	Retail Outlets	Quarterly	Rolling and Quarterly
Raspberry (GB)	Retail Outlets	Quarterly	Quarterly
Raspberry (NI)	Retail Outlets	Quarterly	Quarterly
Spring green and kale (GB)	Retail Outlets	Quarter 2 - 4	Quarter 2 - 4
Spring green and kale (NI)	Retail Outlets	Quarter 2 - 4	Quarter 2 - 4

### Animal products

Food	Sampling points	Sampled during	Reporting
Beef (GB)	Retail Outlets	Quarterly	Quarterly
Beef (NI)	Retail Outlets	Quarterly	Quarterly
Cheese (soft) (GB)	Retail Outlets	Quarterly	Quarters 1, 2 and 4

Cheese (soft) (NI)	Retail Outlets	Quarterly	Quarterly
Eggs (GB)	Retail Outlets	Quarterly	Quarterly
Eggs (NI)	Retail Outlets	Quarterly	Quarters 1, 3 and 4
Fish (white) (NI)	Retail Outlets	Quarterly	Quarterly
Milk (GB)	Retail Outlets	Quarterly	Quarterly
Milk (NI)	Retail Outlets	Quarterly	Quarterly

### Cereal products

Food	Sampling points	Sampled during	Reporting
Bread (GB)	Retail Outlets	Quarters 2 - 4	Quarter 3 and 4
Bread (NI)	Retail Outlets	Quarters 2 - 4	Quarter 3 and 4
Rice (GB)	Retail Outlets	Quarter 1 and 3	Quarter 1 and 3
Rice (NI)	Retail Outlets	Quarter 1 and 3	Quarter 1 and 3
Wheat (GB)	Retail Outlets	Quarterly	Quarter 2 and 4
Wheat (NI)	Retail Outlets	Quarterly	Quarter 2 and 4

### Miscellaneous products

Food	Sampling points	Sampled during	Reporting
Edible seeds (GB)	Retail Outlets	Quarter 2 and 3	Quarter 3
Infant food (cereal based) (GB)	Retail Outlets	Quarter 3	Quarter 4
Infant food (cereal based) (NI)	Retail Outlets	Quarter 3	Quarter 4
Nuts (GB)	Retail Outlets	Quarters 3 and 4	Quarter 4
Olive Oil (GB)	Retail Outlets	Quarterly	Quarters 2 - 4
Olive oil (NI)	Retail Outlets	Quarterly	Quarters 2 - 4
Plant based protein (GB)	Retail Outlets	Quarter 3	Quarter 3
Processed peppers (GB)	Retail Outlets	Quarter 3	Quarter 4
Soya based drink (GB)	Retail Outlets	Quarter 3	Quarter 3

### Sampling points

- Retail outlets: samples bought by market research contractor shoppers.
- Supply Chain: samples taken by inspectors from the Animal and Plant Health Agency from a range of points in the supply chain (wholesalers, retail depots, ports and import points).

### Reporting

- Results for certain higher-priority foods are produced, followed up and published more frequently at [Data.gov.uk](https://data.gov.uk)
- All results are published in the quarterly report. Some surveys are included in every quarter, some are every other quarter and some in just one quarter.

### Table 3: Summary of Results

Food	Analysed	With residues at or below the MRL <sup>2</sup>	With residues above the MRL <sup>3</sup>	With residues of non- approved pesticides (UK only)	With multiple residues	Organic samples tested	Organic samples with residues
Asparagus (GB)	24	1	0	0	0	0	0
Asparagus (NI)	6	0	0	0	0	0	0
Aubergine (GB)	25	19	2	0	10	2	0
Aubergine (NI)	11	8	1	0	4	0	0
Banana (GB)	13	8	2	0	9	2	0
Banana (NI)	13	11	0	0	11	2	0
Beans with Pods (GB)	32	19	5	0	16	0	0
Beans with pods (NI)	8	5	1	0	2	0	0
Beef (GB)	24	0	0	0	0	1	0
Beef (NI)	16	1	1	0	0	3	0

<sup>2</sup> In analytical terms this is a reportable value between LOD and the MRL

<sup>3</sup> Samples collected in GB must comply with GB set MRLs unless the goods are qualifying Northern Ireland goods and are subject to unfettered access under the terms of the UK Internal Market Act 2020. For samples collected in NI, under the Northern Ireland Protocol, certain aspects of EU food law, including compliance with EU MRLs apply.



Food	Analysed	With residues at or below the MRL <sup>2</sup>	With residues above the MRL <sup>3</sup>	With residues of non- approved pesticides (UK only)	With multiple residues	Organic samples tested	Organic samples with residues
Berries and small fruits (GB)	31	22	4	2	22	1	0
Berries and small fruits (NI)	10	8	0	0	6	0	0
Bread (GB)	54	47	1	0	28	0	0
Bread (NI)	20	18	0	0	11	0	0
Broccoli (GB)	31	20	0	2	8	3	0
Broccoli (NI)	10	7	0	0	4	2	0
Cheese (soft) (GB)	48	23	2	0	1	0	0
Cheese (soft) (NI)	15	0	0	0	0	0	0
Eggs (GB)	24	1	0	0	0	4	0
Eggs (NI)	14	0	0	0	0	5	0
Fish (white) (NI)	16	7	0	0	1	0	0
Grapefruit (GB)	24	24	0	0	24	0	0
Grapefruit (NI)	14	13	1	0	14	0	0

Food	Analysed	With residues at or below the MRL <sup>2</sup>	With residues above the MRL <sup>3</sup>	With residues of non- approved pesticides (UK only)	With multiple residues	Organic samples tested	Organic samples with residues
Grapes (GB)	18	18	0	0	17	0	0
Grapes (NI)	12	12	0	0	11	0	0
Infant food (cereal based) (GB)	36	1	13 <sup>4</sup>	0	1	13	1
Infant food (cereal based) (NI)	10	1	0	0	0	2	0
Melon (GB)	30	27	0	0	21	0	0
Melon (NI)	11	9	0	0	6	0	0
Milk (GB)	78	0	0	0	0	6	0
Milk (NI)	16	0	0	0	0	0	0
Mushroom (GB)	18	8	0	0	4	4	0
Mushroom (NI)	14	10	1	1	6	1	1
Nuts (GB)	48	16	3	0	0	3	0
Olive oil (GB)	18	7	0	0	1	2	0

<sup>4</sup> These are all residues of chlorate that are unlikely to be from pesticide use. See p71 for more information.

Food	Analysed	With residues at or below the MRL <sup>2</sup>	With residues above the MRL <sup>3</sup>	With residues of non- approved pesticides (UK only)	With multiple residues	Organic samples tested	Organic samples with residues
Olive oil (NI)	3	0	0	0	0	1	0
Peppers (GB)	30	30	0	0	21	1	1
Peppers (NI)	14	10	0	0	8	2	1
Peppers (processed) (GB)	24	10	2	0	9	0	0
Potatoes (GB)	61	36	0	0	8	1	0
Potatoes (NI)	14	5	0	0	2	4	1
Raspberry (GB)	29	22	0	1	16	0	0
Raspberry (NI)	10	6	0	0	4	0	0
Spring greens and kale (GB)	30	28	1	11	28	1	1
Spring greens and kale (NI)	12	9	1	2	9	0	0
Wheat (GB)	36	30	0	0	9	5	0
Wheat (NI)	30	25	0	0	8	4	0

**Table 4: Summary of samples sent to FSA for follow up**

Sample ID	Date of Sampling	Description	Country of Origin	Retail Outlet	Address	Brand Name	Packer / Manufacturer	Pesticide residues found in mg/kg (MRL)	INFOSAN Ref. No;
4259/2021	17/11/2021	Yard Long Beans	India	MH Malik Produce Ltd	P9 Produce Hall, Western International Market, Hayes Road, Southall, UB2 5YG	None stated	Soya Dhillon Export Dhillon House, Plot no. 72, Sector 23, Turbhe, Navi, Mumbai 400,706	omethoate 0.1 (MRL = 0.01*)	
								captan (sum) 0.8 (MRL = 0.03*)	
								carbendazim (sum) 0.1 (MRL = 0.2)	
								dimethoate 0.02 (MRL = 0.01*)	
								dithiocarbamates 0.2 (MRL = 1)	
								hexaconazole 0.08 (MRL = 0.01*)	
4193/2021	05/10/2021	Yard Long Beans	India	MK Impex Ltd	P16 Produce Hall, Western International Market, Hayes Road, Southall UB2 5XJ	None stated	MK Exports B/4/03 Groma House, Plot no. 14 - C, Sector 19, Vashi, Navi Mumbai - 400703, India	dithiocarbamates 0.9 (MRL = 1)	I-002-906*
								dimethoate 0.4 (MRL = 0.01*)	
								cypermethrin (sum) 0.07 (MRL = 0.7)	I-002-906*
								profenofos 0.3 (MRL = 0.01*)	
								omethoate 0.2 (MRL = 0.01*)	
hexaconazole 0.2 (MRL = 0.01*)									

Sample ID	Date of Sampling	Description	Country of Origin	Retail Outlet	Address	Brand Name	Packer / Manufacturer	Pesticide residues found in mg/kg (MRL)	INFOSAN Ref. No;
3995/2021	19/10/2021	Guar Beans	India	AR Butt	Unit 37/39, St James Market, Essex Street, Bradford BD4 7PN	None stated	Indo British Exports B-1 1st Floor, Export Building, APMC, Veg Market, Vashi, Navi Mumbai, 400705, India	chlorpyrifos 0.03 (MRL = 0.01*)	
								monocrotophos 0.004 (MRL = 0.01*)	
								omethoate 0.003 (MRL = 0.01*)	
4248/2021	07/12/2021	Cavendish bananas	Ecuador	Alam & Sons	32-40 St James Market, Essex Street, Bradford BD4 7PN	None stated	CIA Coexfruit S.A. Machala, El Oro	azoxystrobin 0.03 (MRL = 2)	
								bifenthrin 0.02 (MRL = 0.1)	
								chlorpyrifos 0.03 (MRL = 0.01*)	
								myclobutanil 0.1 (MRL = 2)	
4171/2021	05/10/2021	Plantain bananas	Guatemala	MM Exotic Ltd	62A New Spitalfields Market, Leyton E10 5SQ	None stated	Colomba Unit 2, Leaside IND EST, 60 German Road, London N17 0QH	chlorpyrifos 0.05 (MRL = 0.01*)	
								thiabendazole 0.3 (MRL = 6)	

\* <https://www.fao.org/food-safety/emergencies/infosan/en/>

## Table 5: Summary of MRL Exceedances

Sample ID	Food Type	Country of Origin	Pesticide Detected	Residue Detected (mg/kg)	MRL <sup>+</sup> (mg/kg)	MRL exceedance after allowing for measurement uncertainty
<b>Aubergine (GB)</b>						
2677/2021	Aubergine	UK	flonicamid (sum)	1	0.5	No
4256/2021	Baby Aubergine	Kenya	profenofos	0.02	0.01*	No
<b>Aubergine (NI)</b>						
5621/2021	Aubergine	Spain	chlorate	0.7	0.4	No
<b>Banana (GB)</b>						
4248/2021	Banana (eating)	Ecuador	chlorpyrifos	0.03	0.01*	Yes
4171/2021	Plantain	Guatemala	chlorpyrifos	0.05	0.01*	Yes
<b>Beans with pods (GB)</b>						
3995/2021	Speciality Beans	India	chlorpyrifos	0.03	0.01*	Yes
4193/2021	Speciality Beans	India	dimethoate	0.4	0.01*	Yes
			hexaconazole	0.2	0.01*	Yes
			omethoate	0.2	0.01*	Yes
			profenofos	0.3	0.01*	Yes

4238/2021	Speciality Beans	India	profenofos	0.04	0.01*	Yes
4259/2021	Speciality Beans	India	captan (sum)	0.8	0.03*	Yes
			dimethoate	0.02	0.01*	No
			hexaconazole	0.08	0.01*	Yes
			omethoate	0.1	0.01*	Yes
4299/2021	Speciality Beans	India	dinotefuran	0.02	0.01	No
<b>Beans with pods (NI)</b>						
0669/2021	Green Beans	Morocco	linuron	0.02	0.01*	Yes
<b>Beef (NI)</b>						
5978/2021	Diced Irish stewing beef	Ireland	DDAC (sum)	0.2	0.1	No
<b>Berries and small fruits (GB)</b>						
0623/2021	Fresh: Blackberries	Wales (UK)	cyantraniliprole	0.04	0.01*	Yes
5533/2021	Fresh: Blackberries	UK	cyantraniliprole	0.03	0.01*	Yes
5914/2021	Fresh: Blackberries	Wales (UK)	cyantraniliprole	0.04	0.01*	Yes
0709/2021	Frozen: Blackberries	UK	flonicamid (sum)	0.06	0.03*	No

**Bread (GB)**

0777/2021	Speciality Bread: Scones	UK	BAC (sum)	0.4	0.1	Yes
-----------	--------------------------	----	-----------	-----	-----	-----

**Cheese (soft) (GB)**

0851/2021	Mozzarella	Italy	chlorate	0.6	0.1	Yes
-----------	------------	-------	----------	-----	-----	-----

5502/2021	Mozzarella	Italy	chlorate	0.3	0.1	Yes
-----------	------------	-------	----------	-----	-----	-----

**Grapefruit (NI)**

5656/2021	Red Grapefruit	South Africa	sulfoxaflor	0.3	0.15	No
-----------	----------------	--------------	-------------	-----	------	----

**Infant food (cereal based) (GB)**

0955/2021	Rusks	UK	chlorate	0.02	0.01*	No
-----------	-------	----	----------	------	-------	----

0980/2021	Creamed Porridge	EU	chlorate	0.5	0.01*	Yes
-----------	------------------	----	----------	-----	-------	-----

1411/2021	Rusks	UK	chlorate	0.02	0.01*	No
-----------	-------	----	----------	------	-------	----

1778/2021	Rusks	UK	chlorate	0.02	0.01*	Yes
-----------	-------	----	----------	------	-------	-----

			diphenylamine	0.02	0.01*	Yes
--	--	--	---------------	------	-------	-----

1954/2021	Rusks	UK	chlorate	0.02	0.01*	Yes
-----------	-------	----	----------	------	-------	-----

2335/2021	Rusks	UK	chlorate	0.02	0.01*	No
-----------	-------	----	----------	------	-------	----

2790/2021	Rusks	UK	chlorate	0.02	0.01*	No
-----------	-------	----	----------	------	-------	----



2873/2021	Rusks	UK	chlorate	0.02	0.01*	No
2984/2021	Rusks	UK	chlorate	0.02	0.01*	Yes
3147/2021	Rusks	UK	chlorate	0.03	0.01*	Yes
3209/2021	Reduced Sugar Rusks	UK	chlorate	0.03	0.01*	Yes
5785/2021	Organic Raspberry and Apple Soft Oaty Bars	UK	Fosetyl (sum)	0.5	0.01*	Yes
5811/2021	Rusks	UK	chlorate	0.02	0.01*	Yes
<b>Mushroom (NI)</b>						
0667/2021	Button	UK	deltamethrin	0.08	0.05	No
<b>Nuts (GB)</b>						
0217/2021	Brazil Nuts	Brazil	inorganic bromide	57	50	No
1830/2021	Brazil Nuts	UK	inorganic bromide	56	50	No
1406/2021	Cashew Nuts	the Netherlands	inorganic bromide	63	50	No
<b>Peppers (processed) (GB)</b>						
0466/2021	Green Sliced Jalapeno Peppers	Turkey	famoxadone	0.02	0.01*	No
0958/2021	Pepper Quarters	Poland	ethephon	0.06	0.05*	No

**Spring greens and kale (GB)**

---

5561/2021	Kale	UK	prosulfocarb	0.02	0.01*	No
-----------	------	----	--------------	------	-------	----

---

**Spring greens and kale (NI)**

---

5658/2021	Kale	UK	prosulfocarb	0.03	0.01*	Yes
-----------	------	----	--------------	------	-------	-----

---

\* **Maximum Residue Levels set at the LOD (LOD MRL):** These MRLs are set at a default level, i.e., at the limit of determination (LOD) where analytical methods can reasonably detect the presence of the pesticide. Either insufficient trials data are available on which to set a maximum residue level, or there may be no use of the pesticide on that crop permitted. However, they may be permitted elsewhere.

+ Samples collected in GB must comply with GB set MRLs unless the goods are qualifying Northern Ireland goods and are subject to unfettered access under the terms of the UK Internal Market Act for 2020. For samples collected in NI, under the Northern Ireland Protocol., certain aspects of EU food law, including compliance with EU set MRLs, continue to apply.

# Section 1: findings by food

## Asparagus (GB)

### Samples tested

24 samples were tested for up to 390 pesticide residues

- 24 samples were imported from outside the EU

### Pesticide residues detected from those sought

- 23 samples contained no residues from those sought
- 1 sample contained a residue above the reporting limit
- None of the samples were labelled as organic.

### Risk assessments

The single residue detected by the laboratory would not be expected to have an effect on health.

### Combined risk assessments

None of the samples contained more than one residue, so we did not carry out a combined risk assessment.

### Additional Comments by the PRiF

None

### Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Asparagus (NI)

### Samples tested

6 samples were tested for up to 389 pesticide residues

- 6 samples were imported from outside the EU

### Pesticide residues detected from those sought

- 6 samples contained no residues from those sought
- None of the samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic

### Risk assessments

The laboratory did not detect any residues, so we did not carry out a risk assessment

### Additional Comments by the PRiF

None

### Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

# Aubergine (GB)

## Samples tested

25 samples were tested for up to 373 pesticide residues

- 3 samples came from the UK
- 1 sample was imported from outside the EU
- 21 samples came from the EU

## Pesticide residues detected from those sought

- 4 samples contained no residues from those sought
- 21 samples contained residues above the reporting limit
- 2 samples contained residues above the MRL
- 2 samples were labelled as organic. None contained residues from those sought

## Risk assessments

The residues detected by the laboratory would not be expected to have an effect on health.

## Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

## Additional Comments by the PRiF

None

## Residues measured above the MRL

The laboratory detected 2 residues above the MRL in aubergines. Details are available in [Table 5](#).

## Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

# Aubergine (NI)

## Samples tested

11 samples were tested for up to 373 pesticide residues

- 11 samples came from the EU

## Pesticide residues detected from those sought

- 2 samples contained no residues from those sought
- 9 samples contained residues above the reporting limit
- 1 sample contained residues above the MRL
- None of the samples were labelled as organic.

## Risk assessments

The residues detected by the laboratory would not be expected to have an effect on health.

## Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

## Additional Comments by the PRiF

None

## Residues measured above the MRL

The laboratory detected 1 residue above the MRL in aubergine. Details are available in [Table 5](#).

## Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Banana (GB)

### Samples tested

13 samples were tested for up to 393 pesticide residues

#### Banana (eating)

- 12 samples were imported from outside the EU

#### Plantain

- 1 sample was imported from outside the EU

### Pesticide residues detected from those sought

- 3 samples contained no residues from those sought
- 10 samples contained residues above the reporting limit
- 2 samples contained residues above the MRL
- 2 samples were labelled as organic. None contained residues from those sought

### Risk assessments

Following screening assessment there were two findings of the pesticides chlorpyrifos, above the MRL of 0.01\* mg/kg, where the effect on health needed to be considered in more detail.

One of the findings was in plantain, where the chlorpyrifos residue was found at a level of 0.05 mg/kg. For plantain, we use consumption data for banana, as we do not hold sufficient consumption data for plantain specifically, so this assessment also covers the other finding of chlorpyrifos of 0.03 mg/kg in a sample of Cavendish bananas.

HSE always undertakes assessments that consider both when the peel is not eaten, and one where it is assumed that all of the peel is eaten. These assessments are detailed in [section 3](#) and should be consulted for the full assessment of risk.

If the banana/plantain is consumed without the peel a short-term effect on health is not expected.

HSE has conducted a worst-case form of the assessment, assuming that all the peel is consumed with the fruit.

In this case the highest residue observed was chlorpyrifos at 0.05 mg/kg in plantain. As outlined in HSE's full risk assessment (section 3), EFSA's 2019 statement on the human health assessment of chlorpyrifos indicated that no toxicological reference values could be determined for chlorpyrifos. Although it is not possible to conclude on the short-term assessment with certainty whether or not presence of food residues of chlorpyrifos at this level would have any effect on health after eating large portions (97.5th percentile consumption) of banana/plantain (with all of the peel), HSE has stated a number of reasons why the detailed assessment they have performed is especially precautionary. Please refer to the full assessment of risk in section 3 for further details.

Despite this, an effect on health would only be anticipated if a number of factors came together at the same time: the high residue found in the banana/plantain sample (0.05 mg/kg) being consumed by the most critical consumer infants, high residue in single fruit

item, peak consumption levels (97.5th percentile), and a large proportion of peel from the fruit being consumed.

The EFSA 2019 statement on the human health assessment of chlorpyrifos also includes a consideration of the potential for genotoxicity (whether damage to genetic material can occur). We conclude that on a precautionary basis any findings of chlorpyrifos are undesirable due to the uncertainty regarding genotoxicity. Due to the low levels of chlorpyrifos (highest only 0.05 mg/kg) in the plantain/banana we consider any risks of adverse health effects are low.

Other risk assessment screening work undertaken did not indicate any other expectation of effects on health. Please refer to [‘how HSE perform the risk assessments’](#) for further details.

## Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

## Additional Comments by the PRiF

### Chlorpyrifos

The potential short-term effect of the chlorpyrifos residues we needed to consider in more detail depends on whether the peel is eaten. We would not expect an effect on health if the peel was not eaten. For the plantain or banana eaten with the peel we cannot conclude whether residues of chlorpyrifos at this level would have any effect on health due to the uncertainties around the toxicological values. We have concluded any risk of adverse health effect from chlorpyrifos due to genotoxicity to be low as explained in the detailed assessment. We note that market approval for chlorpyrifos has been withdrawn in many jurisdictions. This is consistent with our view that any residue of chlorpyrifos is undesirable, due to uncertainty in the genotoxicity safety assessment.

## Residues measured above the MRL

The laboratory detected 2 residues above the MRL, one in bananas and one in plantain. Details are available in [Table 5](#).

HSE have passed details of the following sample to FSA for further consideration. Further details are in [Table 4](#).

- Cavendish banana from Ecuador containing chlorpyrifos at 0.03 mg/kg
- Plantain from Guatemala containing chlorpyrifos at 0.05 mg/kg

## Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)



- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Banana (NI)

### Samples tested

13 samples were tested for up to 392 pesticide residues

### Bananas (eating)

- 13 samples were imported from outside the EU

### Pesticide residues detected from those sought

- 2 samples contained no residues from those sought
- 11 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- 2 samples were labelled as organic. None contained residues from those sought

### Risk assessments

Following screening assessment there was one sample of banana containing 1.4 mg/kg thiabendazole, where the effect on health needed to be considered in more detail.

HSE always undertakes assessments that consider both when the peel is not eaten, and one where it is assumed that all of the peel is eaten. These assessments are detailed in [section 3](#) and should be consulted for the full assessment of risk.

If the bananas are consumed without the peel, an effect on health is not expected.

HSE have conducted a worst-case form of the assessment, assuming that all the peel is consumed with the fruit. In this case, an effect on health would be unlikely. Additionally, an effect on health would only be anticipated if a number of factors came together at the same time: the high residue found in the banana sample being consumed by the most critical consumer – infant, high residue in a single fruit item, peak consumption levels (97.5th percentile), and a large proportion of peel from the fruit being consumed.

Other risk assessment screening work undertaken did not indicate any other expectation of effects on health. Please refer to [‘how HSE perform the risk assessments’](#) for further details.

### Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

## Additional Comments by the PRiF

Following the HSE risk assessment we needed to consider a sample of banana containing thiabendazole further. We would not expect an effect on health if the peel was not eaten and even if all of the peel was eaten an effect is unlikely.

## Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Beans with pods (GB)

### Samples tested

32 samples were tested for up to 390 pesticide residues

#### Dwarf Beans

- 1 sample was imported from outside the EU

#### Fine Beans

- 8 samples were imported from outside the EU

#### Green Beans

- 8 samples were imported from outside the EU

#### Runner Beans

- 2 samples were imported from outside the EU

#### Speciality Beans

- 8 samples were imported from outside the EU
- 5 samples came from the EU

### Pesticide residues detected from those sought

- 8 samples contained no residues from those sought
- 24 samples contained residues above the reporting limit
- 5 samples contained residues above the MRL
- None of the samples were labelled as organic.

### Risk assessments

Following screening assessment there were two pesticides (chlorpyrifos and dimethoate (and its metabolite omethoate)) where the effect on health needed to be considered in more detail.

Chlorpyrifos was found in a sample of Guar beans at a level of 0.03 mg/kg, above the MRL of 0.01\* mg/kg. Based on the HSE assessment of short-term risk for chlorpyrifos (see section 3), we conclude an effect on health is not expected.

As outlined in HSE's full risk assessment (section 3), EFSA issued a 2019 statement on the human health assessment of chlorpyrifos which included a consideration of the potential for genotoxicity (whether damage to genetic material can occur). We conclude that on a precautionary basis any findings of chlorpyrifos are undesirable due to the uncertainty regarding genotoxicity. Due to the low level of chlorpyrifos (only 0.03 mg/kg) in the Guar beans we consider any risks of adverse health effects are low.

Dimethoate and omethoate are chemically related pesticides and for toxicology purposes are considered together. Omethoate is also the main metabolite of dimethoate. Residues of dimethoate and/or omethoate were found in several samples of beans. We have provided a detailed risk assessment (see section 3 for full details) for a sample of yard long beans containing the highest levels observed at 0.4 mg/kg and 0.2 mg/kg respectively, above the MRLs of 0.01\* mg/kg. The EFSA Conclusion (2018) for dimethoate has indicated that no toxicological reference values could be determined for dimethoate and omethoate, due to a lack of a fully supporting toxicological database. We think that at the anticipated highest exposures following consumption of yard long beans containing these residues of dimethoate and omethoate, there is unlikely to be a risk of ill health effects based on short term toxicity.

The EFSA Conclusion (2018) for dimethoate also includes a consideration of the potential for genotoxicity (whether damage to genetic material can occur). We conclude that on a precautionary basis any findings of dimethoate and omethoate are undesirable due to the uncertainty regarding genotoxicity at low doses. Due to the low levels of dimethoate and omethoate (up to 0.4 mg/kg dimethoate and 0.2 mg/kg omethoate) in the beans with pods we consider any risks of adverse health effects are low.

Monocrotophos was found in a sample of Guar beans at level of 0.004 mg/kg below the MRL of 0.01\* mg/kg. Although at a low level, where a short-term effect on health is not expected, there is uncertainty about the potential for monocrotophos to cause genetic damage at low doses. Therefore, on a precautionary basis we consider any findings of monocrotophos in food undesirable. Due to the low level of monocrotophos (only 0.004 mg/kg) in the Guar beans we consider any risks of adverse health effects are low.

Please refer to the full assessment of risk in [section 3](#) for further details.

Other risk assessment screening work undertaken did not indicate any other expectation of effects on health. Please refer to '[how HSE perform the risk assessments](#)' for further details.

## **Combined risk assessments**

Some samples contained residues of more than one pesticide. Some of these residues are from pesticides which belong to similar chemical groups and may have similar toxicological effects. So, the risk assessors needed to consider their possible impacts on human health, both on their own and in combination.

HSE carried out a combined risk assessment of the relevant samples, as these contained different pesticides which might have similar effects; these pesticides are known to inhibit the enzyme acetylcholinesterase (please see the [glossary](#) on page 159).

One sample of Guar beans contained chlorpyrifos, omethoate and monocrotophos. HSE's assessment of this combination concluded that these pesticides together at the levels found will be unlikely to inhibit acetyl cholinesterase, the known effect from exposure to each of these residues. HSE concludes that a short-term effect on health is unlikely.

One sample of yard long beans contained dimethoate, it's metabolite omethoate and profenofos. CRD's combined risk assessment on this combination concluded that it is not anticipated that the profenofos residues will contribute significantly to the overall combined intake when compared to the single substance assessment conducted for dimethoate and its metabolite omethoate in this sample. HSE's assessment of this combination concluded that these pesticides together at the levels found will be unlikely to inhibit acetyl

cholinesterase, the known effect from exposure to each of these residues. HSE concludes that a short-term effect on health is unlikely.

It is noted that each of chlorpyrifos, monocrotophos and dimethoate, and its metabolite omethoate, have shown some potential to be genotoxic (cause genetic damage). Please refer to the above comments (under risk assessment) and details in section 3 on '[Substances that might be genotoxic](#)' for HSE's conclusions regarding potential genotoxicity.

These combined assessments are detailed in [section 3](#) and should be consulted for the full assessment of risk.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

## **Additional Comments by the PRiF**

We needed to consider several samples of beans with pods containing either chlorpyrifos, monocrotophos or dimethoate and its metabolite omethoate in more detail. HSE's assessment of short-term risk, also considering any combinations of these and other residues of pesticides known to inhibit acetyl cholinesterase when found together in the same samples, concluded the likelihood of adverse health effects to be unlikely.

We also concluded any risk of adverse health effect from chlorpyrifos, monocrotophos or dimethoate, and its metabolite omethoate, due to genotoxicity to be low. We note that market approval for these pesticides has been withdrawn in many jurisdictions. This is consistent with our view that any residue of dimethoate, its metabolite omethoate, monocrotophos or chlorpyrifos is undesirable, due to uncertainty in the genotoxicity safety assessment.

## **Residues measured above the MRL**

The laboratory detected 11 residues above the MRL in beans with pods. Details are available in [Table 5](#).

HSE has passed details of the following sample to FSA for further consideration. Further details are in [Table 4](#).

- Yard long beans from India containing dimethoate at 0.4 mg/kg, omethoate at 0.2 mg/kg and profenofos at 0.3 mg/kg
- Yard long beans from India containing dimethoate at 0.02 mg/kg and omethoate at 0.1 mg/kg
- Guar beans from India containing chlorpyrifos at 0.03mg/kg, monocrotophos at 0.004 mg/kg and omethoate at 0.003 mg/kg

## **Further Information**

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Beans with pods (NI)

### Samples tested

8 samples were tested for up to 389 pesticide residues

#### Fine Beans

- 5 samples were imported from outside the EU

#### Green Beans

- 3 samples were imported from outside the EU

### Pesticide residues detected from those sought

- 2 samples contained no residues from those sought
- 6 samples contained residues above the reporting limit
- 1 sample contained residues above the MRL
- None of the samples were labelled as organic.

### Risk assessments

The residues detected by the laboratory would not be expected to have an effect on health.

### Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

### Additional Comments by the PRiF

None

### Residues measured above the MRL

The laboratory detected 1 residue above the MRL in beans with pods. Details are available in [Table 5](#).

### Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Beef (GB)

### Samples tested

24 samples were tested for up to 116 pesticide residues

- 21 samples came from the UK
- 3 samples came from the EU

The country of origin of samples may not be the same as the country where the beef was produced. It may be where the beef was processed, where it was packed for consumer purchase or the address of the brand owner.

### Pesticide residues detected from those sought

- 24 samples contained no residues from those sought
- None of the samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- 1 sample was labelled as organic. None contained residues from those sought

### Risk assessments

The laboratory did not detect any residues, so we did not carry out a risk assessment

### Additional Comments by the PRiF

None

### Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found



## Beef (NI)

### Samples tested

16 samples were tested for up to 38 pesticide residues

- 15 samples came from the UK
- 1 sample came from the EU

The country of origin of samples may not be the same as the country where the beef was produced. It may be where the beef was processed, where it was packed for consumer purchase or the address of the brand owner.

### Pesticide residues detected from those sought

- 14 samples contained no residues from those sought
- 2 samples contained residues above the reporting limit
- 1 sample contained residues above the MRL
- 3 samples were labelled as organic. None contained residues from those sought

### Risk assessments

The residues detected by the laboratory would not be expected to have an effect on health.

### Combined risk assessments

None of the samples contained more than one residue, so we did not carry out a combined risk assessment.

### Additional Comments by the PRiF

One sample of diced beef contained a residue of BAC and 1 sample of diced beef contained DDAC. These substances are widely used as biocides (disinfectants) during food preparation and processing. This is the most likely source of the residues.

### Residues measured above the MRL

The laboratory detected 1 residue above the MRL in beef. Details are available in [Table 5](#)

### Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Berries and small fruits (GB)

### Samples tested

31 samples were tested for up to 368 pesticide residues

#### Fresh: Blackberries

- 5 samples came from the UK

#### Fresh: Blueberries

- 5 samples came from the UK
- 13 samples were imported from outside the EU
- 2 samples came from the EU

#### Frozen: Blackberries

- 3 samples came from the UK

#### Frozen: Blueberries

- 1 sample came from the UK
- 1 sample came from the EU

#### Frozen: Cranberries

- 1 sample came from the UK

The country of origin of frozen samples may not be the same as the country where the berries were grown. It may be where the berries were processed, where they were packed for consumer purchase or the address of the brand owner.

### Pesticide residues detected from those sought

- 4 samples contained no residues from those sought
- 27 samples contained residues above the reporting limit
- 4 samples contained residues above the MRL
- 1 sample was labelled as organic. None contained residues from those sought.

### Risk assessments

The residues detected by the laboratory would not be expected to have an effect on health.

### Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE considers combined risk assessments is in [Section 3: HSE assessment of risk](#)

## **Additional Comments by the PRiF**

### **Interpretation of UK origin**

Where frozen berries are labelled as a UK product this reflects either the name of the business the food is marketed under or the address of the business that has imported the food. Usually, the fruit was not grown in the UK.

### **Residues measured above the MRL**

The laboratory detected 4 residues above the MRL in blackberries. Details are available in [Table 5](#).

### **Further Information**

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Berries and small fruits (NI)

### Samples tested

10 samples were tested for up to 368 pesticide residues

#### Fresh: Blackberries

- 1 sample came from the UK
- 1 sample was imported from outside the EU
- 1 sample came from the EU

#### Fresh: Blueberries

- 6 samples were imported from outside the EU

#### Frozen: Blackberries

- 1 sample came from the UK

The country of origin of frozen samples may not be the same as the country where the berries were grown. It may be where the berries were processed, where they were packed for consumer purchase or the address of the brand owner.

### Pesticide residues detected from those sought

- 2 samples contained no residues from those sought
- 8 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

### Risk assessments

The residues detected by the laboratory would not be expected to have an effect on health.

### Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE considers combined risk assessments is in [Section 3: HSE assessment of risk](#).

### Additional Comments by the PRiF

#### Interpretation of UK origin

Where frozen berries are labelled as a UK product this reflects either the name of the business the food is marketed under or the address of the business that has imported the food. Usually, the fruit was not grown in the UK.

## Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Bread (GB)

### Samples tested

54 samples were tested for up to 393 pesticide residues

#### Ordinary Bread: Other

- 3 samples came from the UK

#### Ordinary Bread: White

- 27 samples came from the UK

#### Ordinary Bread: Wholemeal

- 12 samples came from the UK

#### Speciality Bread: Brioche

- 2 samples came from the UK

#### Speciality Bread: Croissants

- 1 sample came from the UK

#### Speciality Bread: Crumpets

- 5 samples came from the UK

#### Speciality Bread: Muffins

- 1 sample came from the UK

#### Speciality Bread: Pancakes

- 1 sample came from the UK

#### Speciality Bread: Scones

- 1 sample came from the UK

#### Speciality Bread: Waffles

- 1 sample came from the UK

The country of origin of samples may not be the same as the country where the bread was produced. It may be where the wheat was processed, where it was packed for consumer purchase or the address of the brand owner.

### Pesticide residues detected from those sought

- 6 samples contained no residues from those sought
- 48 samples contained residues above the reporting limit
- 1 sample contained residues above the MRL
- None of the samples were labelled as organic.

More information on how MRLs for individual, unprocessed ingredients also apply to processed and compound foods, and how MRLs were adjusted to take account of processing, is in [section 4](#).

## **Risk assessments**

The residues detected by the laboratory would not be expected to have an effect on health.

## **Combined risk assessments**

Some samples contained residues of more than one pesticide. Two residues in one sample are from pesticides which belong to similar chemical groups and may have similar toxicological effects. So, the risk assessors needed to consider their possible impacts on human health, both on their own and in combination.

HSE carried out a combined risk assessment of the relevant sample. We would not expect any of these combinations to have an effect on health.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

## **Additional Comments by the PRiF Speciality bread**

Each year we include a type of speciality bread in the monitoring programme. This year we are looking at morning bakery products. This includes items such as crumpets, bagels, waffles, croissants, and pancakes amongst other similar items.

## **Residues measured above the MRL**

The laboratory detected 1 residue above the MRL in bread (speciality). Details are available in [Table 5](#)

## **Further Information**

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Bread (NI)

### Samples tested

20 samples were tested for up to 392 pesticide residues

#### Ordinary Bread: Brown

- 2 samples came from the UK
- 1 sample came from the EU

#### Ordinary Bread: Other

- 1 sample came from the UK

#### Ordinary Bread: White

- 11 samples came from the UK
- 2 samples came from the EU

#### Ordinary Bread: Wholemeal

- 3 samples came from the UK

The country of origin of samples may not be the same as the country where the bread was produced. It may be where the wheat was processed, where it was packed for consumer purchase or the address of the brand owner.

### Pesticide residues detected from those sought

- 2 samples contained no residues from those sought
- 18 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

More information on how MRLs for individual, unprocessed ingredients also apply to processed and compound foods, and how MRLs were adjusted to take account of processing, is in [section 4](#).

### Risk assessments

The residues detected by the laboratory would not be expected to have an effect on health.

### Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)



## Additional Comments by the PRiF

None

## Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Broccoli (GB)

### Samples tested

31 samples were tested for up to 370 pesticide residues

#### Fresh

- 25 samples came from the UK
- 2 samples were imported from outside the EU
- 4 samples came from the EU

### Pesticide residues detected from those sought

- 11 samples contained no residues from those sought
- 20 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- 3 samples were labelled as organic. None contained residues from those sought

### Risk assessments

Three samples of broccoli contained residues of triallate, a herbicide (weed killer), at levels of either 0.01 or 0.02 mg/kg, below the MRL of 0.1\* mg/kg. Although low level residues, where a short-term effect on health is not expected, there is uncertainty about the potential for triallate to cause genetic damage at low doses. Therefore, on a precautionary basis we consider any findings of triallate in food undesirable. Due to the low level of triallate (highest only up to 0.02 mg/kg) in broccoli we consider any risks of adverse health effects are low.

Other risk assessment screening work undertaken did not indicate any other expectation of effects on health. Please refer to [‘how HSE perform the risk assessments’](#) for further details.

### Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

### Additional Comments by the PRiF

We have asked HSE to update us on the regulatory issues concerning the residues of triallate detected. For the moment this aspect will be picked up in the routine domestic review programme. HSE is following up these residues with growers as this use is not permitted on this crop.

### Further Information

- [Summary table of results](#)
- [Survey Design](#)

- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Broccoli (NI)

### Samples tested

10 samples were tested for up to 370 pesticide residues

#### Fresh

- 4 samples came from the UK
- 1 sample was imported from outside the EU
- 5 samples came from the EU

### Pesticide residues detected from those sought

- 3 samples contained no residues from those sought
- 7 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- 2 samples were labelled as organic. None contained residues from those sought

### Risk assessments

The residues detected by the laboratory would not be expected to have an effect on health.

### Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

### Additional Comments by the PRiF

None

### Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## **Cheese (soft) (GB)**

### **Samples tested**

48 samples were tested for up to 113 pesticide residues

#### **Brie**

- 1 sample came from the UK
- 10 samples came from the EU

#### **Camembert**

- 1 sample came from the UK
- 9 samples came from the EU

#### **Cottage Cheese**

- 3 samples came from the UK

#### **Cream Cheese**

- 1 sample came from the EU

#### **Feta**

- 6 samples came from the EU

#### **Mozzarella**

- 1 sample came from the UK
- 4 samples came from the EU

#### **Ricotta**

- 7 samples came from the EU

#### **Soft Cheese**

- 2 samples came from the UK
- 3 samples came from the EU

The country of origin of samples may not be the same as the country where the cheese was produced. It may be where the cheese was processed, where it was packed for consumer purchase or the address of the brand owner.

### **Pesticide residues detected from those sought**

- 23 samples contained no residues from those sought
- 25 samples contained residues above the reporting limit
- 2 samples contained residues above the MRL
- None of the samples were labelled as organic.

### **Risk assessments**

The residues detected by the laboratory would not be expected to have an effect on health.

## Combined risk assessments

One sample contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in this sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

## Additional Comments by the PRiF

### Chlorate

We have not marked two samples where we measured chlorate over the MRL for milk as breaches of the legislation. These were a sample of ricotta cheese made in Italy and a sample of cottage cheese made in the UK. The brand owners have provided evidence that these residues are from treated potable (drinking) water used during production processes. Legislation allows for chlorate MRLs to be exceeded where the source of the chlorate is treated water or biocides appropriately used post-harvest. HSE have concluded that these findings are not breaches of the MRL. For more information on chlorate see [Section 4. 'issues arising in this report Chlorate'](#)

### DDT

DDT is known to be a soil contaminant. At the low levels found in this report we consider this example likely to be as a result of historic use. More information on DDT can be found in [Section 4 'issues arising in this report. DDT'](#)

## Residues measured above the MRL

The laboratory detected 4 residues above the MRL in cheese. Two were considered compliant having arisen from processing. Details of the remaining residues are available in [Table 5](#)

## Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Cheese (soft) (NI)

### Samples tested

15 samples were tested for up to 38 pesticide residues

#### Brie

- 3 samples came from the UK
- 2 samples came from the EU

#### Camembert

- 2 samples came from the EU

#### Feta

- 3 samples came from the EU

#### Mascarpone

- 1 sample came from the EU

#### Mozzarella

- 1 sample came from the EU

#### Ricotta

1 sample came from the EU

#### Soft Cheese

- 2 samples came from the EU

The country of origin of samples may not be the same as the country where the cheese was produced. It may be where the cheese was processed, where it was packed for consumer purchase or the address of the brand owner.

### Pesticide residues detected from those sought

- 15 samples contained no residues from those sought
- None of the samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

### Risk assessments

The laboratory did not detect any residues, so we did not carry out a risk assessment.

### Additional Comments by the PRiF

None

### Further Information

- [Summary table of results](#)
- [Survey Design](#)

- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found



## Eggs (GB)

### Samples tested

24 samples were tested for up to 116 pesticide residues

#### Duck

- 2 samples came from the UK

#### Hens

- 22 samples came from the UK

### Pesticide residues detected from those sought

- 23 samples contained no residues from those sought
- 1 sample contained residues above the reporting limit
- None of the samples contained residues above the MRL
- 4 samples were labelled as organic. None contained residues from those sought

### Risk assessments

The single residue detected by the laboratory would not be expected to have an effect on health.

### Combined risk assessments

None of the samples contained more than one residue, so we did not carry out a combined risk assessment.

### Additional Comments by the PRiF

#### DDT

We looked further at a sample of egg containing DDT below the MRL. HSE is aware that such instances might be a result of contaminated soil from historic use of DDT, as hens are known to swallow soil particles or grit to aid in the breakdown of food material in the gizzard. We informed Defra's Veterinary Medicine Division (VMD) of these findings. VMD has concluded that this was probably not a residue resulting from current use but instead reflects the potential of findings of DDT as a historical environment contaminant. This is accounted for in the existing MRL. More information on DDT can be found in Section 4 ['issues arising in this report. DDT'](#)

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Eggs (NI)

### Samples tested

14 samples were tested for up to 117 pesticide residues

### Hens

- 14 samples came from the UK

### Pesticide residues detected from those sought

- 14 samples contained no residues from those sought
- None of the samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- 5 samples were labelled as organic. None contained residues from those sought

### Risk assessments

The laboratory did not detect any residues, so we did not carry out a risk assessment.

### Additional Comments by the PRiF

None

### Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Fish (white) (NI)

### Samples tested

16 samples were tested for up to 38 pesticide residues

#### Basa

- 2 samples were imported from outside the EU

#### Cod

- 7 samples were imported from outside the EU

#### Haddock

- 1 sample was imported from outside the EU

#### Hake

- 3 samples were imported from outside the EU
- 1 sample came from the EU

#### Sea bream

- 2 samples were imported from outside the EU

Where no sea area information is available, the country of origin on the packaging does not necessarily indicate where the fish was caught or farmed. It could be where it was landed or processed or where it was packed for retail sale.

### Pesticide residues detected from those sought

- 9 samples contained no residues from those sought
- 7 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

### Risk assessments

The residues detected by the laboratory would not be expected to have an effect on health.

### Combined risk assessments

One sample contained residues of more than one pesticide. Some of these residues are from pesticides which belong to similar chemical groups and may have similar toxicological effects. So, the risk assessors needed to consider their possible impacts on human health, both on their own and in combination.

HSE carried out a combined risk assessment of the relevant sample. We would not expect any of these combinations to have an effect on health.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

## Additional Comments by the PRiF

### BAC

5 samples, 4 of cod and 1 of hake, contained a residue of BAC. This substance is widely used as biocides (disinfectants) during food preparation and processing. This is the most likely source of the residue.

### Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

# Grapefruit (GB)

## Samples tested

24 samples were tested for up to 391 pesticide residues

- 23 samples were imported from outside the EU
- 1 sample came from the EU

## Pesticide residues detected from those sought

- All samples contained residues
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

## Risk assessments

Following screening assessment there were two pesticides, imazalil and thiabendazole, where the effect on health needed to be considered in more detail.

HSE always undertakes assessments that consider both when the peel is not eaten, and one where it is assumed that all of the peel is eaten. These assessments are detailed in [section 3](#) and should be consulted for the full assessment of risk.

For both of these pesticides, if the grapefruit are consumed without the peel an effect on health is not expected.

HSE has conducted a worst-case form of the assessment, assuming that all the peel is consumed with the fruit. In this case, for both thiabendazole and imazalil an effect on health would be unlikely. Additionally, an effect on health would only be anticipated if a number of factors came together at the same time: the high residue found in the grapefruit sample being consumed by the most critical consumer – infants, high residue in a single fruit item, peak consumption levels (97.5th percentile), and a large proportion of peel from the fruit being consumed.

Other risk assessment screening work undertaken did not indicate any other expectation of effects on health. Please refer to [‘how HSE perform the risk assessments’](#) for further details.

## Combined risk assessments

All samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

## Additional Comments by the PRiF

Following the HSE risk assessment, we needed to consider residues of two pesticides, imazalil and thiabendazole in more detail. We would not expect an effect on health if the peel was not eaten and even if all of the peel was eaten an effect is unlikely.

## Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Grapefruit (NI)

### Samples tested

14 samples were tested for up to 390 pesticide residues

- 12 samples were imported from outside the EU
- 2 samples came from the EU

### Pesticide residues detected from those sought

- All samples contained residues
- 1 sample contained residues above the MRL
- None of the samples were labelled as organic.

### Risk assessments

Following screening assessment there were two pesticides, thiabendazole and imazalil, where the effect on health needed to be considered in more detail.

The highest level of thiabendazole at (2.9 mg/kg) is lower than the level of thiabendazole (3.3 mg/kg) found in a sample reported for Grapefruit-GB. The highest level of imazalil at (2.4 mg/kg) is lower than the level of imazalil (2.7 mg/kg) reported in a sample for Grapefruit-GB. Therefore, the risk assessments carried out for thiabendazole and imazalil and the conclusions (see Grapefruit-GB) also apply to Grapefruit-NI.

As such, in both these cases, we conclude that an effect on health based on short-term toxicity would not be expected if the peel is removed prior to consuming, and unlikely even if the grapefruit was consumed with all of its peel. Please see [section 3](#) for full details of HSE's assessment.

Other risk assessment screening work undertaken did not indicate any other expectation of effects on health. Please refer to [‘how HSE perform the risk assessments’](#) for further details.

### Combined risk assessments

All samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

### Additional Comments by the PRiF

Please see our comment for grapefruit GB survey where we found higher levels of these residues. Following the HSE risk assessment, we needed to consider residues of two pesticides imazalil and thiabendazole in more detail. We would not expect an effect on health if the peel was not eaten and even if all of the peel was eaten an effect is unlikely.

## Residues measured above the MRL

The laboratory detected 1 residue above the MRL in grapefruit. Details are available in [Table 5](#)

## Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found



## Grapes (GB)

### Samples tested

18 samples were tested for up to 397 pesticide residues

- 3 samples were imported from outside the EU
- 15 samples came from the EU

### Pesticide residues detected from those sought

- All samples contained residues
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

### Risk assessments

Following screening assessment there was one sample containing ethephon at 0.9 mg/kg, below the MRL of 1.0 mg/kg, where the effect on health needed to be considered in more detail.

Based on the HSE assessment of short-term risk for ethephon (see [section 3](#)), we conclude an effect on health would be unlikely.

Other risk assessment screening work undertaken did not indicate any other expectation of effects on health. Please refer to [‘how HSE perform the risk assessments’](#) for further details.

### Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

### Additional Comments by the PRiF

Following the HSE risk assessment, we needed to consider one sample in more detail, with a residue of ethephon at 0.9 mg/kg. We consider that an effect on health would be unlikely.

Grapes are available throughout the year, imported from all around the world and so they have been surveyed as part of our rolling reporting programme [GOV.UK](#) for a number of years

### Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)

- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Grapes (NI)

### Samples tested

12 samples were tested for up to 396 pesticide residues

- 4 samples were imported from outside the EU
- 8 samples came from the EU

### Pesticide residues detected from those sought

- All samples contained residues
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

### Risk assessments

The residues detected by the laboratory would not be expected to have an effect on health.

### Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

### Additional Comments by the PRiF

None

### Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Infant food (cereal based) (GB)

### Samples tested

36 samples were tested for up to 397 pesticide residues

- 28 samples came from the UK
- 8 samples came from the EU

The country of origin of samples may not be the same as the country where the infant food was produced. It may be where the infant food processed, where they were packed for consumer purchase or the address of the brand owner.

### Pesticide residues detected from those sought

- 22 samples contained no residues from those sought
- 14 samples contained residues above the reporting limit
- 13 samples contained residues above the MRL
- 13 samples were labelled as organic. 1 contained residues from those sought

### Risk assessments

A sample of creamed porridge infant food contained a residue of chlorate at 0.5 mg/kg where the effect on health needed to be considered in more detail. Based on the HSE Chemicals Regulation Division's risk assessment (please see [Section 3](#)) of the residue detected we consider a short-term effect on health to be unlikely.

This conclusion is based on assessment of peak daily amounts of infant food consumed (all types of infant foods consumed throughout the day). However, it is noted that the highest residue of chlorate in infant food in this report (0.5 mg/kg) was found in jars of creamed porridge. The porridge is marketed as a breakfast cereal. If the porridge containing chlorate at 0.5 mg/kg accounted for half an infant's daily diet, then there would be no exceedance of the ARfD and an effect on health would not be expected. If it accounted for the entire daily diet, then the ARfD would be exceeded but an effect on health would still be unlikely.

Chlorate was found in 13 of the 46 infant food samples, and aside from the highest residue of chlorate of 0.5 mg/kg assessed here, the next highest residue of chlorate was 0.03 mg/kg. A chlorate residue at this level (0.03 mg/kg) would lead to dietary intakes within the ARfD, and an effect on health would not be expected.

Other risk assessment screening work undertaken did not indicate any other expectation of effects on health. Please refer to ['how HSE perform the risk assessments'](#) for further details.

### Combined risk assessments

One sample contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in this sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

## Comments from Department of Health and Social Care (DHSC)

In accordance with the UK-wide Nutrition Related Labelling, Composition and Standards (NLCS) provisional common framework, it is the responsibility of UK authorities to conduct the appropriate risk assessment and risk management processes (including seeking scientific evaluation) of amendments to GB domestic legislation.

The PRiF and industry stakeholders have raised concerns to the NLCS policy group which is established under the NLCS framework, and comprises of officials from across the UK governments around the ability of industry to meet the required maximum residue level (MRL) set for infant and follow-on formula and infant food under Commission Delegated Regulation (EU) 2016/127 on infant and follow-on formula and the Processed Cereal-based Foods and Baby Foods for Infants and Young Children (England) Regulations 2003 respectively.

The NLCS policy group are therefore considering the requirement on pesticides as per Article 4 of Commission Delegated Regulation (EU) 2016/127 and regulation 7 of the Processed Cereal-based Foods and Baby Foods for Infants and Young Children (England) Regulations 2003.

The NLCS framework was established to maintain a consistent and co-ordinated policy approach for Nutrition Legislation across the UK, and to manage divergence with Northern Ireland. The current approach to the Protocol on Ireland/Northern Ireland (NIP) means that EU legislation relating to nutrition, as detailed in Annex 2 of the NIP, continues to be directly applicable in Northern Ireland (NI). Therefore, any decision made will need to take into account the impact on Northern Ireland and the UK internal market.

A summary of the EU Regulations which the UK authorities have considered under the arrangements agreed in the NLCS provisional common framework between 01.01.2021 - 31.12.2021 have been published and are available on [gov.uk](https://www.gov.uk).

## Additional Comments by the PRiF

### Chlorate

We needed to consider residues of chlorate in more detail and following HSE's short-term assessment of risk, concluded that we would not expect any effects on health at a level of 0.03 mg/kg, and that an effect on health was unlikely if the highest residue of 0.5 mg/kg was found.

The legal Maximum Residue Level for chlorate in foods for infants is set at 0.01 mg/kg, and there is no provision to take into account unavoidable residues from sources other than pesticide use. We have been updated by the manufacturer and their trade groups of the strategies the industry is adopting to minimise the presence of chlorate without compromising microbiological safety.

We are satisfied that the origin of the residues we have found is chlorate is from drinking water or other potable water used in manufacturing, and/or chlorate as a residue from disinfectant use. We have reminded manufacturers of our on-going advice not to take any action that could affect microbiological safety without careful consideration.

We had previously raised our consideration that the default MRL was not always achievable in the case of chlorate with the lead government department, the Department of Health and Social Care. We are grateful for the update provided from the Nutrition

Related Labelling, Composition and Standards (NLCS) Group and the ongoing work they are doing to consider possible steps that can be taken.

## **Residues measured above the MRL**

The laboratory detected 14 residues above the MRL in infant food (cereal based). Details are available in [Table 5](#)

## **Further Information**

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

# Infant food (cereal based) (NI)

## Samples tested

10 samples were tested for up to 395 pesticide residues

- 7 samples came from the UK
- 3 samples came from the EU

The country of origin of samples may not be the same as the country where the infant food was produced. It may be where the infant food processed, where they were packed for consumer purchase or the address of the brand owner.

## Pesticide residues detected from those sought

- 9 samples contained no residues from those sought
- 1 sample contained residues above the reporting limit
- None of the samples contained residues above the MRL
- 2 samples were labelled as organic. None contained residues from those sought

## Risk assessments

The single residue detected by the laboratory would not be expected to have an effect on health.

## Combined risk assessments

None of the samples contained more than one residue, so we did not carry out a combined risk assessment.

## Additional Comments by the PRiF

None

## Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Melon (GB)

### Samples tested

30 samples were tested for up to 372 pesticide residues

#### Cantaloupe

- 4 samples were imported from outside the EU

#### Galia

- 5 samples were imported from outside the EU

#### Honeydew

- 11 samples were imported from outside the EU

#### Watermelon

- 10 samples were imported from outside the EU

The country of origin of prepared samples may not be the same as the country where the melon was grown. It may be where the melon was processed, where it was packed for consumer purchase or the address of the brand owner.

### Pesticide residues detected from those sought

- 3 samples contained no residues from those sought
- 27 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

### Risk assessments

Following screening assessment there was one sample of melon containing 1.1 mg/kg imazalil, where the effect on health needed to be considered in more detail.

HSE always undertake assessments that consider both when the peel is not eaten, and one where it is assumed that all of the peel is eaten. These assessments are detailed in section 3 and should be consulted for the full assessment of risk.

If the melon is consumed without the peel, an effect on health is not expected.

HSE has conducted a worst case form of the assessment, assuming that all the peel is consumed with the fruit, and we conclude that an effect on health would be unlikely. Additionally, an effect on health would only be anticipated if a number of factors came together at the same time: the high residue found in the melon sample being consumed by the most critical consumer – 11-14 year old, high residue in a single fruit item, peak consumption levels (97.5th percentile), and a large proportion of peel from the fruit being consumed.

Other risk assessment screening work undertaken did not indicate any other expectation of effects on health. Please refer to [‘how HSE perform the risk assessments’](#) for further details.



## Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

## Additional Comments by the PRiF

Following the HSE risk assessment we needed to consider a sample of melon containing imazalil. We would not expect an effect on health if the melon peel was not eaten. Even if all the peel was consumed with the fruit, we consider an effect on health unlikely.

## Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Melon (NI)

### Samples tested

11 samples were tested for up to 372 pesticide residues

#### Galia

- 1 sample was imported from outside the EU

#### Honeydew

- 5 samples were imported from outside the EU

#### Watermelon

- 4 samples were imported from outside the EU

#### Yellow

- 1 sample was imported from outside the EU

The country of origin of prepared samples may not be the same as the country where the melon was grown. It may be where the melon was processed, where it was packed for consumer purchase or the address of the brand owner.

### Pesticide residues detected from those sought

- 2 samples contained no residues from those sought
- 9 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

### Risk assessments

The residues detected by the laboratory would not be expected to have an effect on health.

### Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

### Additional Comments by the PRiF

None

### Further Information

- [Summary table of results](#)
- [Survey Design](#)

- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Milk (GB)

### Samples tested

78 samples were tested for up to 113 pesticide residues

#### Cow's milk

- 76 samples came from the UK

#### Goats milk

- 2 samples came from the UK

### Pesticide residues detected from those sought

- 78 samples contained no residues from those sought
- None of the samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- 6 samples were labelled as organic. None contained residues from those sought

### Risk assessments

The laboratory did not detect any residues, so we did not carry out a risk assessment.

### Additional Comments by the PRiF

None

### Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Milk (NI)

### Samples tested

16 samples were tested for up to 114 pesticide residues

### Cow's milk

- 16 samples came from the UK

### Pesticide residues detected from those sought

- 16 samples contained no residues from those sought
- None of the samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

### Risk assessments

The laboratory did not detect any residues, so we did not carry out a risk assessment.

### Additional Comments by the PRiF

None

### Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Mushroom (GB)

### Samples tested

18 samples were tested for up to 395 pesticide residues

#### Button & Other

- 4 samples came from the UK
- 4 samples came from the EU

#### Chestnut

- 7 samples came from the UK
- 1 sample came from the EU

#### Flat

- 1 sample came from the UK

#### Forestiere

- 1 sample came from the EU

### Pesticide residues detected from those sought

- 10 samples contained no residues from those sought
- 8 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- 4 samples were labelled as organic. None contained residues from those sought

### Risk assessments

The residues detected by the laboratory would not be expected to have an effect on health.

### Combined risk assessments

Some samples contained residues of more than one pesticide. Two residues in one sample are from pesticides which belong to similar chemical groups and may have similar toxicological effects. So, the risk assessors needed to consider their possible impacts on human health, both on their own and in combination.

HSE carried out a combined risk assessment of the relevant sample. We would not expect any of these combinations to have an effect on health.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

### Additional Comments by the PRiF

We would not expect any of the residues detected to have an effect on health.

Chlormequat and mepiquat residues detected are unlikely to be from unauthorised use of either of these active substances on mushrooms. Both are used on cereal crops, and residues can be transmitted from straw used as a growing medium into the mushrooms.

## Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Mushroom (NI)

### Samples tested

14 samples were tested for up to 395 pesticide residues

### Button & Other

- 8 samples came from the UK
- 6 samples came from the EU

### Pesticide residues detected from those sought

- 3 samples contained no residues from those sought
- 11 samples contained residues above the reporting limit
- 1 sample contained residues above the MRL
- 1 sample was labelled as organic. 1 contained residues from those sought

### Risk assessments

The residues detected by the laboratory would not be expected to have an effect on health.

### Combined risk assessments

Some samples contained residues of more than one pesticide. Two residues in one sample are from pesticides which belong to similar chemical groups and may have similar toxicological effects. So, the risk assessors needed to consider their possible impacts on human health, both on their own and in combination.

HSE carried out a combined risk assessment of the relevant sample. We would not expect any of these combinations to have an effect on health.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

### Additional Comments by the PRiF

We would not expect any of the residues detected to have an effect on health.

As with the chlormequat findings in mushroom GB samples chlormequat and mepiquat residues detected are unlikely to be from unauthorised use of either of these active substances on mushrooms. Both are used on cereal crops, and residues can be transmitted from straw used as a growing medium into the mushrooms.

### Residues measured above the MRL

The laboratory detected 1 residue above the MRL in mushrooms. Details are available in [Table 5](#)

### Further Information

- [Summary table of results](#)
- [Survey Design](#)



- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Nuts (GB)

### Samples tested

48 samples were tested for up to 388 pesticide residues

#### Almonds

- 3 samples came from the UK
- 5 samples were imported from outside the EU

#### Brazil Nuts

- 2 samples came from the UK
- 3 samples were imported from outside the EU

#### Cashew Nuts

- 8 samples came from the UK
- 6 samples were imported from outside the EU
- 2 samples came from the EU

#### Peanuts

- 5 samples came from the UK
- 2 samples were imported from outside the EU
- 1 sample came from the EU

#### Pecan

- 1 sample came from the UK

#### Walnuts

- 3 samples came from the UK
- 7 samples were imported from outside the EU

The country of origin of samples may not be the same as the country where the nuts were grown. It may be where the nuts were processed, where they were packed for consumer purchase or the address of the brand owner.

### Pesticide residues detected from those sought

- 29 samples contained no residues from those sought
- 19 samples contained residues above the reporting limit
- 3 samples contained residues above the MRL
- 3 samples were labelled as organic. None contained residues from those sought

### Risk assessments

The residues detected by the laboratory would not be expected to have an effect on health.

### Combined risk assessments

None of the samples contained more than one residue, so we did not carry out a combined risk assessment.

## **Additional Comments by the PRiF**

We detected residues of bromide in 5 samples of Brazil nuts, 1 sample of cashews and 1 sample of walnuts. We think these residues are a result of naturally occurring levels of bromine and agree with HSE conclusion that these are not breaches of the MRL. The MRLs for these foods were set at a level intended to take account of naturally occurring levels, and we have measured residues only just above those levels.

The MRLs for these foods were set some years ago at a level that should take account of naturally occurring levels.

## **Residues measured above the MRL**

The laboratory detected 3 residues above the MRL in nuts. Details are available in [Table 5](#)

## **Further Information**

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Olive oil (GB)

### Samples tested

18 samples were tested for up to 392 pesticide residues

#### Blended

- 1 sample came from the UK

#### Extra Virgin Olive Oil

- 1 sample came from the UK
- 15 samples came from the EU

#### Virgin Olive Oil

- 1 sample came from the EU

The country of origin of samples may not be the same as the country the olive oil was produced. It may be where the olive oil was processed, where they were packed for consumer purchase or the address of the brand owner.

### Pesticide residues detected from those sought

- 11 samples contained no residues from those sought
- 7 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- 2 samples were labelled as organic. None contained residues from those sought

More information on how MRLs for individual, unprocessed ingredients also apply to processed and compound foods, and how MRLs were adjusted to take account of processing, is in [section 4](#)

### Risk assessments

Following screening assessment there were two pesticides, chlorpyrifos and dimethoate, each found below their respective MRLs of 0.01\* mg/kg, where the effect on health needed to be considered in more detail.

Based on the HSE assessment of short-term risk for chlorpyrifos (see [section 3](#)), we conclude an effect on health is not expected.

As outlined in HSE's full risk assessment ([section 3](#)), EFSA issued a 2019 statement on the human health assessment of chlorpyrifos which included a consideration of the potential for genotoxicity (whether damage to genetic material can occur). We conclude that on a precautionary basis any findings of chlorpyrifos are undesirable due to the uncertainty regarding genotoxicity. Due to the low level of chlorpyrifos (only 0.003 mg/kg) in the olive oil we consider any risks of adverse health effects are low.

Based on the HSE assessment of short-term risk for dimethoate (see [section 3](#)), we conclude an effect on health is not expected.

In 2018, EFSA reviewed dimethoate and concluded that no toxicological reference values could be determined for dimethoate, due to a lack of a fully supporting toxicological database (EFSA Conclusion for dimethoate, 2018). At the anticipated highest exposure,

an effect on health is not expected based on short-term toxicity. Please refer to [section 3](#) for full details of HSE's assessment.

The EFSA Conclusion (2018) for dimethoate also includes a consideration of the potential for genotoxicity (whether damage to genetic material can occur) for dimethoate. We conclude that on a precautionary basis any findings of dimethoate are undesirable due to the uncertainty regarding genotoxicity at low doses. Due to the low level of dimethoate (only 0.007 mg/kg) in the olive oil we consider any risks of adverse health effects are low.

Other risk assessment screening work undertaken did not indicate any other expectation of effects on health. Please refer to ['how HSE perform the risk assessments'](#) for further details.

## Combined risk assessments

One sample contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in this sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

## Additional Comments by the PRiF

We needed to consider residues of two pesticides, chlorpyrifos and dimethoate, in more detail and following HSE's assessment of risk, concluded the likelihood of adverse health effects based on short term toxicity to be not expected.

We also concluded any risk of adverse health effect from chlorpyrifos or dimethoate due to genotoxicity to be low. We note that market approval for both pesticides has been withdrawn in many jurisdictions. This is consistent with our view that any residue of chlorpyrifos is undesirable, due to uncertainty in the genotoxicity safety assessment.

## Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Olive oil (NI)

### Samples tested

3 samples were tested for up to 390 pesticide residues

#### Extra Virgin Olive Oil

- 2 samples came from the EU

#### Virgin Olive Oil

- 1 sample came from the EU

The country of origin of samples may not be the same as the country the olive oil was produced. It may be where the olive oil was processed, where they were packed for consumer purchase or the address of the brand owner.

### Pesticide residues detected from those sought

- 3 samples contained no residues from those sought
- None of the samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- 1 sample was labelled as organic. None contained residues from those sought

More information on how MRLs for individual, unprocessed ingredients also apply to processed and compound foods, and how MRLs were adjusted to take account of processing, is in [section 4](#)

### Risk assessments

The laboratory did not detect any residues, so we did not carry out a risk assessment.

### Additional Comments by the PRiF

None

### Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

# Peppers (GB)

## Samples tested

30 samples were tested for up to 395 pesticide residues

### Fresh

- 1 sample came from the UK
- 2 samples were imported from outside the EU
- 27 samples came from the EU

The country of origin of samples may not be the same as the country where the peppers were grown. It may be where the peppers were processed, where they were packed for consumer purchase or the address of the brand owner.

## Pesticide residues detected from those sought

- All samples contained residues
- None of the samples contained residues above the MRL
- 1 sample was labelled as organic. 1 contained residues from those sought

## Risk assessments

The residues detected by the laboratory would not be expected to have an effect on health.

## Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

## Additional Comments by the PRiF

None

## Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Peppers (NI)

### Samples tested

14 samples were tested for up to 394 pesticide residues

#### Fresh

- 14 samples came from the EU

The country of origin of samples may not be the same as the country where the peppers were grown. It may be where the peppers were processed, where they were packed for consumer purchase or the address of the brand owner.

- Pesticide residues detected from those sought
- 4 samples contained no residues from those sought
- 10 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- 2 samples were labelled as organic. 1 contained residues from those sought

### Risk assessments

The residues detected by the laboratory would not be expected to have an effect on health.

### Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

### Additional Comments by the PRiF

None

### Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found



## Peppers (processed) (GB)

### Samples tested

24 samples were tested for up to 391 pesticide residues

- 3 samples came from the UK
- 18 samples were imported from outside the EU
- 3 samples came from the EU

The country of origin of samples may not be the same as the country where the peppers were produced. It may be where the peppers were processed, where they were packed for consumer purchase or the address of the brand owner.

### Pesticide residues detected from those sought

- 12 samples contained no residues from those sought
- 12 samples contained residues above the reporting limit
- 2 samples contained residues above the MRL
- None of the samples were labelled as organic.

### Risk assessments

Following screening assessment there was one pesticide, chlorpyrifos, found below the MRL of 0.01\* mg/kg, where the effect on health needed to be considered in more detail.

Based on the HSE assessment of short-term risk for chlorpyrifos (see [section 3](#)), we conclude an effect on health is not expected.

As outlined in HSE's full risk assessment ([section 3](#)), EFSA issued a 2019 statement on the human health assessment of chlorpyrifos which included a consideration of the potential for genotoxicity (whether damage to genetic material can occur). We conclude that on a precautionary basis any findings of chlorpyrifos are undesirable due to the uncertainty regarding genotoxicity. Due to the low level of chlorpyrifos (only 0.003 mg/kg) in the processed pepper we consider any risks of adverse health effects are low.

Other risk assessment screening work undertaken did not indicate any other expectation of effects on health. Please refer to ['how HSE perform the risk assessments'](#) for further details.

### Combined risk assessments

Some samples contained residues of more than one pesticide. Two residues in one sample are from pesticides which belong to similar chemical groups and may have similar toxicological effects. So, the risk assessors needed to consider their possible impacts on human health, both on their own and in combination.

HSE carried out a combined risk assessment of the relevant sample. We would not expect any of these combinations to have an effect on health.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

## **Additional Comments by the PRiF**

We needed to consider a residue of chlorpyrifos in more detail. Following HSE's assessment of risk, we concluded the likelihood of adverse health effects based on short term toxicity to be not expected.

We also concluded any risk of adverse health effect from chlorpyrifos due to genotoxicity to be low. We note that market approval for chlorpyrifos has been withdrawn in many jurisdictions. This is consistent with our view that any residue of chlorpyrifos is undesirable, due to uncertainty in the genotoxicity safety assessment.

## **Residues measured above the MRL**

The laboratory detected 2 residues above the MRL in peppers (processed). Details are available in [Table 5](#)

## **Further Information**

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

# Potatoes (GB)

## Samples tested

61 samples were tested for up to 390 pesticide residues

- 60 samples came from the UK
- 1 sample came from the EU

## Pesticide residues detected from those sought

- 25 samples contained no residues from those sought
- 36 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- 1 sample was labelled as organic. None contained residues from those sought

## Risk assessments

The residues detected by the laboratory would not be expected to have an effect on health.

## Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

## Additional Comments by the PRiF

None

## Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

# Potatoes (NI)

## Samples tested

14 samples were tested for up to 390 pesticide residues

- 14 samples came from the UK

## Pesticide residues detected from those sought

- 9 samples contained no residues from those sought
- 5 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- 4 samples were labelled as organic. 1 contained residues from those sought

## Risk assessments

The residues detected by the laboratory would not be expected to have an effect on health.

## Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

## Additional Comments by the PRiF

None

## Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

# Raspberry (GB)

## Samples tested

29 samples were tested for up to 368 pesticide residues

### Fresh

- 12 samples came from the UK
- 9 samples were imported from outside the EU
- 3 samples came from the EU

### Frozen

- 4 samples came from the UK
- 1 sample was imported from outside the EU

The country of origin of samples may not be the same as the country where the raspberries were grown. It may be where the raspberries were processed, where they were packed for consumer purchase or the address of the brand owner.

## Pesticide residues detected from those sought

- 7 samples contained no residues from those sought
- 22 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

## Risk assessments

The residues detected by the laboratory would not be expected to have an effect on health.

## Combined risk assessments

None

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

## Additional Comments by the PRiF

### Interpretation of UK origin

Where frozen berries are labelled as a UK product this reflects either the name of the business the food is marketed under or the address of the business that has imported the food. Usually, the fruit was not grown in the UK.

## Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)

- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

# Raspberry (NI)

## Samples tested

10 samples were tested for up to 368 pesticide residues

### Fresh

- 4 samples were imported from outside the EU
- 5 samples came from the EU

### Frozen

- 1 sample came from the UK

The country of origin of samples may not be the same as the country where the raspberries were grown. It may be where the raspberries were processed, where they were packed for consumer purchase or the address of the brand owner.

## Pesticide residues detected from those sought

- 4 samples contained no residues from those sought
- 6 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

## Risk assessments

The residues detected by the laboratory would not be expected to have an effect on health.

## Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

## Additional Comments by the PRiF

### Interpretation of UK origin

Where frozen berries are labelled as a UK product this reflects either the name of the business the food is marketed under or the address of the business that has imported the food. Usually, the fruit was not grown in the UK.

## Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)

- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found



## Spring greens and kale (GB)

### Samples tested

30 samples were tested for up to 366 pesticide residues

#### Kale

- 17 samples came from the UK

#### Spring Greens

- 13 samples came from the UK

The country of origin of samples may not be the same as the country where the spring greens and kale were grown. It may be where the spring greens and kale were processed, where they were packed for consumer purchase or the address of the brand owner.

### Pesticide residues detected from those sought

- 1 sample contained no residues from those sought
- 29 samples contained residues above the reporting limit
- 1 sample contained residues above the MRL
- 1 sample was labelled as organic. 1 contained residues from those sought

### Risk assessments

Following screening assessment there was one sample of kale containing cypermethrin at 0.04 mg/kg, below the MRL of 1.0 mg/kg, where the effect on short term health needed to be considered in more detail.

Cypermethrin is an insecticide which is available in different isomeric forms. We understand that the application made to the crop was alpha-cypermethrin, so we have conducted the assessment based on alpha-cypermethrin, which is the more toxic form.

This assessment has used the specific ARfD for alpha-cypermethrin. In this case, HSE's assessment concludes that a short-term effect on health is unlikely. [Full risk assessment](#) is available at page 119

Thirteen samples of kale or spring greens contained residues of triallate, a herbicide (weed killer), at levels ranging 0.01 to 0.08 mg/kg, below the MRL of 0.1\* mg/kg. Although low level residues, where a short-term effect on health is not expected, there is uncertainty about the potential for triallate to cause genetic damage at low doses. Therefore, on a precautionary basis we consider any findings of triallate in food undesirable. Due to the low level of triallate (highest only up to 0.08 mg/kg) in the spring greens or kale we consider any risks of adverse health effects are low.

Other risk assessment screening work undertaken did not indicate any other expectation of effects on health. Please refer to ['how HSE perform the risk assessments'](#) for further details.

### Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide

residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

## **Additional Comments by the PRiF**

We have asked HSE to update us on the regulatory issues concerning triallate. For the moment this aspect will be picked up in the routine domestic review programme. HSE is following up these residues with growers as this use is not permitted.

## **Residues measured above the MRL**

The laboratory detected 1 residue above the MRL in kale. Details are available in [Table 5](#)

## **Further Information**

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Spring greens and kale (NI)

### Samples tested

12 samples were tested for up to 366 pesticide residues

#### Kale

- 8 samples came from the UK

#### Spring Greens

- 4 samples came from the UK

The country of origin of samples may not be the same as the country where the spring greens and kale were grown. It may be where the spring greens and kale were processed, where they were packed for consumer purchase or the address of the brand owner.

### Pesticide residues detected from those sought

- 2 samples contained no residues from those sought
- 10 samples contained residues above the reporting limit
- 1 sample contained residues above the MRL
- None of the samples were labelled as organic.

### Risk assessments

Three samples of kale or spring greens contained residues of triallate, a herbicide (weed killer), at levels ranging 0.01 to 0.08 mg/kg, below the MRL of 0.1\* mg/kg. Although low level residues, where a short-term effect on health is not expected, there is uncertainty about the potential for triallate to cause genetic damage at low doses. Therefore, on a precautionary basis we consider any findings of triallate in food undesirable. Due to the low level of triallate (highest only up to 0.08 mg/kg) in the spring greens or kale we consider any risks of adverse health effects are low.

Other risk assessment screening work undertaken did not indicate any other expectation of effects on health. Please refer to [‘how HSE perform the risk assessments’](#) for further details.

### Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

### Additional Comments by the PRiF

We have asked HSE to update us on the regulatory issues concerning triallate. For the moment this aspect will be picked up in the routine domestic review programme. HSE is following up these residues with growers as this use is not permitted.

## Residues measured above the MRL

The laboratory detected 1 residue above the MRL in kale. Details are available in [Table 5](#)

## Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Wheat (GB)

### Samples tested

36 samples were tested for up to 396 pesticide residues

#### Wheat

- 33 samples came from the UK
- 1 sample came from the EU

#### Wholemeal Wheat Flour

- 2 samples came from the UK

The country of origin of samples may not be the same as the country where the wheat was grown. It may be where the wheat was processed, where it was packed for consumer purchase or the address of the brand owner.

### Pesticide residues detected from those sought

- 6 samples contained no residues from those sought
- 30 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- 5 samples were labelled as organic. None contained residues from those sought

More information on how MRLs for individual, unprocessed ingredients also apply to processed and compound foods, and how MRLs were adjusted to take account of processing is in [section 4](#)

### Risk assessments

The residues detected by the laboratory would not be expected to have an effect on health.

### Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

### Additional Comments by the PRiF

None

### Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)

- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found

## Wheat (NI)

### Samples tested

30 samples were tested for up to 395 pesticide residues

#### Wheat

- 25 samples came from the UK
- 4 samples came from the EU

#### Wholemeal Wheat Flour

- 1 sample came from the UK

The country of origin of samples may not be the same as the country where the wheat was produced. It may be where the wheat was processed, where it was packed for consumer purchase or the address of the brand owner.

### Pesticide residues detected from those sought

- 5 samples contained no residues from those sought
- 25 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- 4 samples were labelled as organic. None contained residues from those sought

More information on how MRLs for individual, unprocessed ingredients also apply to processed and compound foods, and how MRLs were adjusted to take account of processing, is in [section 4](#)

### Risk assessments

The residues detected by the laboratory would not be expected to have an effect on health.

### Combined risk assessments

Some samples contained residues of more than one pesticide. Two residues in one sample are from pesticides which belong to similar chemical groups and may have similar toxicological effects. So, the risk assessors needed to consider their possible impacts on human health, both on their own and in combination.

HSE carried out a combined risk assessment of the relevant sample. We would not expect any of these combinations to have an effect on health.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#)

### Additional Comments by the PRiF

None

### Further Information

- [Summary table of results](#)

- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
  - Brand name, sampling point and origin information
  - Pesticides sought and residues found



# Section 2: Sample details and supplier responses

## Sample details

The sample details are published on [Pesticide Residues in Food Quarterly Data](#) as a dataset in ODS format.

### About sample information

The following information is available on each sample collected this quarter:

- Date and place of collection
- Description (e.g., 'runner bean', organic milk);
- Country of origin or manufacture;
- Brand name and packer/manufacturer; and
- Residues detected (results shown in green indicate residues above the MRL).
- Where the brand name of a sample is given the produce involved may have been on sale in other retail premises at the same time.

The description and country of origin are taken from labelling on the food or at the point of sale. The country of origin of processed food may not be the country where the unprocessed produce was produced. This is true even of food that has undergone minimal processing, such as meat that has been butchered or frozen vegetables.

Samples with residues above the MRL are in bold, green text.

Some brand name details have been withheld – these will be published once enquiries are complete.

### The Government's 'brand naming' policy

The Government has decided that brand name information should be published as part of the Government food chemical surveillance programme. Brand names have been published for most pesticide residue surveys since 1998. Certain samples are excluded from the release of brand name information. These include samples taken as part of any pesticide residues enforcement programme and those taken as part of surveys to study individual people/farms. This policy was reviewed in 2000/1, when Ministers agreed to its continuation.

Where we find residues above an MRL or the presence of non-authorized plant protection product, brand owners/retailers/ growers are notified of the result in advance of publication of reports and given four weeks to comment.

### Interpreting brand name information

There is no ready definition of what constitutes a brand in all cases. For clearly branded produce like breakfast cereals or biscuits the "brand owner" is shown. In the case of "own brand" goods this may be one of the multiple retailers. For fruit and vegetables, the retailer is generally shown. For meat, milk, and most other animal products the retailer is also generally shown. Finally, for all commodities the country of origin is shown where this was displayed either on the produce or in the store.

Our programme aims to take samples of produce in approximate proportion to the market share of the main retailers. This has been done to ensure we obtain an accurate representation of a sector (e.g., fruit and vegetables).

Individual programmes are not capable of generating statistically valid information on residues in particular crops from particular retailers. This would require the collection of a much larger number of samples: either substantially increasing costs or greatly reducing the range of different foods sampled in any one year. Therefore, results from an individual survey cannot be taken as a fair representation of the residues status of any particular brand.

However, we do collect samples from a variety of outlets in a range of locations, over a period of years. Successive programmes should therefore help generate information on the typical residues profile of particular types of produce and on major trends in the incidence and levels of pesticides. It should be noted that this quarterly report is not intended to give a comprehensive comparison with previous surveys of the same commodities.

A particular issue arises in relation to the country of origin of fruit and vegetables. The origins included in the reports are those recorded either on the produce or in the store. However, it is not uncommon for mixing to occur on shop shelves. We have responded by increasing the proportion of pre-packed goods sampled. However, pre-packed samples are not available for some produce in some stores, and it could also introduce bias to surveys if loose produce were not sampled. Loose produce is therefore sampled but the origin of the sample should be interpreted with a degree of caution.

## Action taken by HSE

HSE wrote to:

- The suppliers of all samples containing residues above the MRL
- The authorities of the exporting countries of all samples containing residues above the MRL
- The suppliers of GB and NI samples that contained residues of actives which do not have a plant protection product authorised for the crop they were detected in.
- The Organics branch of Defra about samples that were labelled as organic and contained any residues of pesticides.
- The suppliers and certification organisation of all organic samples containing any residues of pesticides.

Recipients of the letters are given 4 weeks to provide a statement for inclusion in the report. The Expert Committee on Pesticide Residues in Food reviews any replies received.

## Supplier responses

We did not receive any supplier responses for inclusion in this report.

## Section 3: HSE assessment of risk

The surveillance programme is designed to enable the regulatory authorities to check that pesticides are being found at levels, as expected, under the MRLs. This confirms that the regulatory processes are working correctly, and as part of this, that pesticide users are complying with any specified conditions that were part of the authorisation. In addition, this work checks that dietary intakes of residues are within acceptable limits. This may be more challenging when pesticide residues are found in food products that have not been grown in the UK or EU, notably when older pesticides have been used. One of the roles of the PRiF, using the work of HSE, is to call out any pesticide residue which is higher than expected and explain more about any risks to consumers from this.

This section details how risks from dietary intakes are assessed.

### When assessments are carried out

HSE performs screening assessment for each residue and commodity combination to identify residue levels that would lead to intakes above the relevant health-based reference doses (these are also sometimes referred to as toxicological reference values “TRVs”). Further information on this screening approach is available on request from HSE. We then present detailed assessments for every case where the actual residue level found could lead to an intake by any group above the reference dose.

### Assessing Dietary intakes

Assessing the acceptability of dietary intakes is complicated. HSE carries out consumer risk assessments for both short-term (peak) and long-term intakes. These assessments use information on food consumption collected in UK dietary surveys in conjunction with the residue levels we find. Occasionally, HSE uses additional pesticide specific information on the losses of residues that occur during preparation and/or cooking of food.

### How the assessment is carried out

#### Short-term risk assessment

HSE calculate short-term intakes (also called NESTIs) using consumption data for high-level (97.5<sup>th</sup> percentile) consumers, based on single-day consumption values and the highest residue found in a food commodity. The residue found is multiplied by a variability factor to take account of the fact that residues may vary between individual items that make up the sample analysed. This is why in some of our detailed risk assessments we refer to some of the general variability factors (of 5, 7 and 10) that are applied in short-term risk assessments. Sometimes, regulatory assessment of data for a pesticide can support an alternative specific value of the variability factor, and where justified, HSE will apply these to the risk assessment and explain this. The estimated intake is compared to the Acute Reference Dose (ARfD). This is done for ten consumer groups: adults, infants, toddlers, 4-6 year olds, 7-10 year olds, 11-14 year olds, 15-18 year olds, vegetarians, elderly living in residential homes and elderly living in their own homes.

#### Long-term risk assessment

HSE also calculate long-term intakes (NEDIs) for high-level (97.5<sup>th</sup> percentile) consumers, but in this case the consumption data are high-level long-term values rather than peak single-day events. Similarly, the residue values used reflect long-term average levels (we

use the median value across each commodity type) rather than occasional high values. Again, these estimates are made for the ten consumer groups. In this case the estimated intake is compared to the Acceptable Daily Intake (ADI).

### **Where do we get Reference Doses from?**

The reference doses (ADI, ARfD or other suitable health-based reference values such as Tolerable Daily Intake (TDI, sometimes used instead of an ADI)) are set following regulatory assessment. In the UK, these values will be reviewed prior to establishment by the Expert Committee on Pesticides (ECP). We currently use reference values from a range of respected sources, including the EU and values set by EFSA. Up to 2019, the UK was part of a harmonised approach to the assessment of pesticide substances in the EU, and we have taken part in the peer review of previously established values. PRiF will also use values from other respected international sources, such as those established by the Joint FAO/WHO Meeting on Pesticide Residues (JMPR, which evaluates and publishes residues and toxicological evaluations of pesticides) and levels set by regulatory authorities in other countries. For a small number of pesticides, the reference doses used have been determined by HSE (e.g., prothiofos, tecnazene). These have not been independently peer-reviewed and should therefore be regarded as provisional.

### **Further explanation of the models we use and application to the PRiF work**

#### **We use Deterministic models**

The assessments we use are 'deterministic' which means we use a defined level of input (such as a median or highest residue and 97.5<sup>th</sup> percentile dietary consumption values). More information on the deterministic intake assessments is available on HSE's website: [The HSE Pesticide Website](#) then search for Consumer Exposure. Here you will find information and further links. See below for an explanation of probabilistic models, where the inputs into the assessments can be varied and more realistic assessments reflecting a range of possible scenarios can be modelled.

#### **Detailed Risk Assessment work is carried out before pesticides are authorised**

The fundamental full complement of risk assessment work for pesticides is done at the pre-authorisation stage considering the residues data packages when trials reflecting the label uses of the pesticides are assessed. These trials profile the highest likely residues that might arise when the pesticides are used as intended (crops and permitted doses of use). MRLs are set on the basis of these data sets, and the post-approval monitoring work then serves as a check for whether residues found are in line with this prior expectation. The risk assessments supporting the MRLs assess the highest residue observed in each crop (or animal product) for short-term assessment. For the long-term MRL assessments, the median residues, across all crops and animal products are taken together to assess combined intakes over the long-term as 'total dietary intakes', taking account of all possible food exposures. Authorisation for pesticide products can only be granted where these assessments of dietary intake do not exceed the health-based reference values.

#### **MRL considerations**

Although MRLs are not safety levels, an MRL would not be established if the residue concentrations measured in the supervised trials used to support the MRL would give rise to health concerns. In most cases residues present at the MRL result in intakes below the

ARfD and the ADI. So even if the MRL is exceeded this does not always lead to an intake above the ARfD or ADI.

### **What happens if we find a pesticide intake above an ADI or an ARfD?**

In addition, an estimated intake in the monitoring work that exceeds the ADI or ARfD does not automatically result in concerns for consumer health, because a protective approach is used in setting the ADI and ARfD. In the unusual circumstance of an intake exceeding the ADI or ARfD, HSE undertake an evaluation of the toxicological data, and we present details of this assessment.

When we present the outcomes of risk assessments, we provide a conclusion on the possible impact on human health based on the degree of concern following the HSE assessment of risk. These conclusions keep to the following order of increasing severity:

Effects on health are not expected (toxicological reference values not exceeded) < unlikely risk (of effects on health) < low risk < higher risk (exposures are undesirable<sup>5</sup>). Most detailed consumer intake assessments that we present with the PRiF reports are for short-term exposure rather than chronic exposure. This is because in most cases the monitoring data show the majority of samples tested contain residues below the reporting limit and so chronic exposure would not present a concern. The reporting limits are set at suitable low levels based on analytical laboratory procedures so that dietary intakes are typically far below the health-based reference values.

Monitoring data presented in the PRiF quarterly reports provides a “snapshot view” of the residues found in specific foods in a survey usually conducted over a 12 week period and limited to around 30 samples for most commodities. It is important not to use isolated findings of higher than expected residues in small surveys to make judgements on long-term effects over a lifetime exposure.

For PRiF work, long-term exposure assessments use median residue levels, rather than the highest residues found. For quarterly assessment (data obtained over three months only) we currently only assess long-term dietary assessment commodity by commodity and not as total dietary intakes across commodities. Even where a number of samples in a PRiF commodity survey do contain the same pesticide, it is very rare that the ‘screening assessment’ we undertake leads to the need for a more detailed assessment to be presented (only where the dietary intakes exceed the ADI or TDI).

We do not see a high number of PRiF samples for any pesticide that exceed an MRL. This means that, over time, median residue levels found in PRiF monitoring don’t often exceed the median residues used in the trials assessed at the time of MRL setting and so don’t often require consideration of long-term effects. When HSE does need to assess long-term effects, it is likely to overestimate exposure to an assessed pesticide residue in a single food item. For a single commodity risk assessment, we assume high level (97.5<sup>th</sup> percentile) consumption at the median residue level in that food for each day of lifetime. Although the HSE long-term assessments by their quarterly nature are indicative only, the assessment we perform is conservative. Furthermore, alternative published assessments (for example those considering trends and large bodies of data) are available which further

---

<sup>5</sup> Furthermore, PRiF will always conclude that on a precautionary basis any findings of genotoxic substances in food are undesirable (please see the explanation regarding genotoxicity in the below section on ‘Implications for health’).

consider the long-term exposure to pesticides<sup>6</sup>. Some pesticides contribute more significantly to long-term dietary intakes across commodities based on their toxicology and prevalence (such as chlorpyrifos, cyfluthrin, deltamethrin, diazinon, dieldrin, dimethoate, dithiocarbamates, fenamiphos, fipronil, imazalil, lambda-cyhalothrin, omethoate and pirimiphos-methyl), based on the chronic exposure assessments presented in EFSA, 2020<sup>7</sup>).

We will continue to have focus on residues of consumer relevance, when they are found, in both UK and NI produced and imported produce. As pesticide use changes, including when there are impacts of regulatory action on pesticides, the profiles of residues in the monitoring can change over time. For example, chlorpropham, which was previously found at levels well above the reporting limits in potatoes, is no longer permitted for use in the UK and EU, and this pesticide will no longer have dietary intakes which take up a substantial portion of its health based reference values.

## Implications for health

Where intakes exceed a reference dose, it is necessary for the underlying toxicological studies (animal studies) to be considered to enable the significance for the consumer of such an exceedance to be understood. Toxicological studies supplied by the registrants in the regulatory data packages are conducted using different doses to determine the nature of any ill health effects as well as the levels at which such effects can be expected to occur.

Toxicological studies that we refer to and use in the HSE risk assessments are conducted using test animals to identify the highest experimental dose that causes no detectable adverse effects (the NOAEL). Where there is more than one relevant toxicological study, the lowest appropriate NOAEL for the most sensitive adverse effect is typically used. There is some uncertainty in extrapolating between animals and people and it is therefore important to use a 'safety factor' to account for sources of variation. This safety factor is incorporated (by dividing the NOAEL by the safety factor) in deriving a reference dose, either an ADI or an ARfD, to which consumer intakes are compared. A safety factor therefore extrapolates from the animal testing to the general population. Factors in the order of x100 are commonly used, x 10 for animal to man, and x10 for within human population differences in sensitivity. However, toxicologists may propose different values (e.g., from 5 to 1000) based on scientific reasoning in accordance with study designs and the quality of the data that has been generated from the studies.

In order to ensure exposures to pesticides do not pose unacceptable risk to humans a wide range of investigations are performed. Most of these are performed on experimental animals because the only end-points that can be examined in human volunteers are those involving observation or blood and urine sampling. Human volunteer studies involving pesticides are not generated in current regulatory work. There is debate at the

---

<sup>6</sup> Total diet studies (e.g. those performed by US FDA [US FDA total diet study program](#), various Nougadère A et al., publications in Environment International journal on TDS in France); see also EFSA evaluations of chronic exposures to pesticides (2020 and 2021 examples included in the list of EFSA publications on cumulative exposure to pesticides outlined in this section (under Multiple residues)). annual EU monitoring data published on the EFSA website: 2018 report (published in 2020) noted below at <https://doi.org/10.2903/j.efsa.2020.6057>).

<sup>7</sup> EFSA (European Food Safety Authority), Medina-Pastor P and Triacchini G, 2020.

The 2018 European Union report on pesticide residues in food. EFSA Journal 2020;18(4):6057, 103 pp. <https://doi.org/10.2903/j.efsa.2020.6057>

international level as to whether human studies that have been generated should be used for risk assessment purposes. The UK policy is not to use these data in pre-authorisation assessments which support the registration of a pesticide; the JMPR chose to apply judgement in the appropriate use of these data if available. The HSE risk assessments will usually refer to test animal species, such as dog, rat, and rabbit. All toxicological work is undertaken based on principles of minimising animal distress. Where scientifically valid human data are available the risk assessments will refer to these as they reduce the uncertainty in the assessment. Therefore, human data is only referred to in more limited circumstances.

Acute (short-term) toxicology is not a concern for all pesticides, as some are not acutely toxic. In terms of the pesticides that have been found in fruit and vegetables through the surveillance programme an acute risk assessment would not be necessary on the following examples: maleic hydrazide, diphenylamine, kresoxim-methyl, and quintozone.

During regulatory assessment, careful consideration is given to any pesticides that may exhibit any potential to be genotoxic (able to damage genetic material) in live animals. In the PRiF programme we note residue types that have been shown in the toxicological data sets to have genotoxic potential or those where data are suggestive of genotoxicity but not certain. There is a small number of cases of older pesticides, likely found only in imported foods, that might be genotoxic. These are examples where modern data to investigate the true genotoxic potential are not expected to be made available. In such situations, we might conclude on a precautionary basis that any findings of these pesticides are undesirable due to the uncertainty regarding genotoxicity, and at low residue levels any risks of adverse health effects are low due to the limited levels of exposure anticipated. PRiF uses low reporting limits for these pesticides to detect these residues even at very low levels, as we know they are of particular interest to consumers.

## **Consumption data and refining the risk assessment**

### **Consumption values**

As the surveillance programme monitors residues in all types of food, from raw commodities (e.g., potatoes) to processed (e.g., wine), dried (e.g., dried fruit) and composite foods (e.g., fruit bread), consumer risk assessments are specifically tailored to address processed and mixed food products. Sometimes this can be affected by availability of consumption data. For example, for pâté, we assess this using consumption data for liver (all types of liver), and for fish pâté we use consumption data for fish (all sources and types of fish). However, we use specific consumption data where FSA have provided data to us (e.g., data on orange juice, dried grapes, and bread). Consumption data are available for most raw commodities, but where data are limited then we will suggest using alternative data. This may involve considering other commodities (e.g., using potato data as 'surrogate' for sweet potato), or alternative sources of consumption data such as EU PRIMo or JMPR consumption and dietary assessment models, to consider items that do not currently feature in UK data sets. Where alternative data are used in our screening and written assessments we explain this in our presentation of the risk assessment work for each quarterly report (for examples, please see the bullet points before the table of detailed risk assessments in [section 3](#)).

### **Fruit and vegetables with removable peel**

For fruit and vegetables that have peel or skin that might not be consumed we present alternative risk assessments for 'without peel -flesh only' where peel versus pulp residue



distribution data is available. As standard, we present an assessment for 'all of the peel' consumed. It is not expected that consumers will always eat peel, so these standard assessments are considered to be highly cautious and not necessarily realistic. Further data are being generated to better understand whether some people do eat the peel of these, and if so, how much of the peel they tend to eat.

## **Dithiocarbamate Residues**

Dithiocarbamate residues are determined as carbon disulphide which is a common product from different dithiocarbamate pesticides. For the risk assessment we take a precautionary approach. For short-term assessment the worst case dithiocarbamate residue is calculated by assuming the residue is derived from thiram (a molecular weight conversion is applied to estimate the level of residue based on thiram) and this is compared to the ARfD for thiram. Where it can be confirmed that a specific dithiocarbamate was applied the equivalent residue of the specific active substance is estimated and the intake compared to the appropriate reference dose. We only present a detailed risk assessment when dietary intake exceeds either the thiram or other suitable reference dose.

The analysis of dithiocarbamates is further complicated by an expectation that some types of crops, such as members of *Brassicaceae* (e.g., watercress) and *Caricaceae* (e.g., papaya) might contain natural sources of sulphur compounds that could be also determined as carbon disulphide during analysis in the laboratory. The PRiF will consider and explain in the report whether residues reported as dithiocarbamates could be from natural sources or whether they have arisen as a result of fungicide treatment.

## **Probabilistic Modelling**

The standard 'deterministic' calculations of consumer exposure used in regulatory assessment and the HSE risk assessments for PRiF work use realistic consumption data and residue levels. However, they tend to overestimate intakes in most circumstances. This is due to the assumptions used; fruit and vegetables would contain high levels of residue in an individual unit and that these would be consumed by high-level consumers. They do not take into account the possible range of residue levels and consumption distributions that may occur in reality. These possible combinations of residues and consumption levels can be taken into account using modelling/simulation techniques to produce probability distributions of residue intake levels to indicate the range of consumer intakes, presented as a probabilistic assessment of consumer exposure. These techniques are not yet routinely used to estimate dietary intakes of pesticide residues.

## **Multiple residues and other developments in risk assessment for pesticides**

The risk assessment process is not standing still. We are aware that some consumers are concerned by the 'cocktail effect'- the possible implications of residues of more than one chemical occurring in, say, a single portion of fruit or vegetables or the interaction between mixtures of pesticides and veterinary medicines at residue levels.

Where more than one pesticide residue is found in a sample, we consider the need for further assessment. The question of which pesticides should be assessed together remains a challenge due to the complexity of the mixtures. In the PRiF work currently, we consider some combinations that we think are a priority (based on toxicological profile or prevalence of the co-occurring residues that are related to one another chemically). If

more than one triazole, or more than one organophosphate/carbamate is found or the following combinations captan/folpet, BAC/DDAC, chlormequat/mepiquat, we will undertake an additional risk assessment. In a 'first step' screening assessment approach we will consider whether the sum of the dietary intakes of each pesticide taken together in that commodity (when expressed as a % of its own reference value) exceeds a total of 100. If this value is not exceeded, then we do not anticipate that there would be an effect on human health and the assessment is not considered further. If this value (of 100) is exceeded (in the initial screen) then we would present a more detailed risk assessment, including a table to show the dietary intakes of each of the pesticides within the group, in the report. Further information is available on: [The HSE Pesticide Website](#). Search for the Data Requirements Introduction and Index and follow the 'consumer exposure' links.

International research is aimed at improving the regulatory assessment of mixtures of residues to help understand whether there are any health implications from any observed combinations of pesticide residues in food. In our work, PRiF aims to keep our assessments at a high regulatory standard taking account of current knowledge in the field. We will look to adapt as new risk assessment approaches develop. We are keen to ensure our reports reflect consumer concerns.

The Food Standards Agency (FSA) asked the Committee on Toxicity of Chemicals in Food, Consumer Products and the Environment (COT) to assess these concerns. Their report "Risk Assessment of Mixtures of Pesticides" was published in 2002<sup>8</sup>.

The Committee concluded that the probability of any health hazard from exposures to mixtures is likely to be small. Nonetheless, it identified areas of uncertainty in the risk assessment process and made recommendations for further work. These fell under the broad headings of regulatory, surveillance, research, and public information issues. An action plan to take forward the recommendations was published by the FSA. A number of research projects were commissioned by the FSA to help progress the action plan.

Further to the work done by COT in 2002, combined assessment methodology has been taken forward at the international level, especially the European Food Safety Authority (EFSA) to develop methodology.

Much of the existing recent work on cumulative exposures to pesticides uses probabilistic models and large EU wide monitoring data sets. Notable work includes the EFSA publications on cumulative exposure cited below. On the basis of the work to date, including consumer assessment case studies, EFSA concludes, with varying degrees of certainty for all the population groups assessed, that consumer risk from dietary cumulative exposure is below the thresholds established by EU risk managers. Further information can be obtained from EFSA's publications, news updates, and FAQs:

- EFSA Feb 2021 (Statement): Comparison of cumulative dietary exposure to pesticide residues for the reference periods 2014–2016 and 2016–2018 [link](#)
- EFSA Feb 2021 (Scientific Report) Cumulative dietary risk assessment of chronic acetylcholinesterase inhibition by residues of pesticides [link](#)
- EFSA April 2020: News Pesticides: first cumulative risk reports published This work is centred on two case studies (outlined below) [link](#)

---

<sup>8</sup> [Foods Standards Agency Risk Assessment of Mixtures of Pesticides \(COT Report, 2002\)](#)

- EFSA April 2020: Cumulative risk assessment of pesticides: FAQ [link](#)
- EFSA April 2020: Cumulative dietary risk characterisation of pesticides that have acute effects on the nervous system [link](#)
- EFSA April 2020: Cumulative dietary risk characterisation of pesticides that have chronic effects on the thyroid [link](#)
- EFSA news update (Jan 2016) Pesticides: breakthrough on cumulative risk assessment [link](#)
- EFSA Sept 2019: Establishment of (CAGs) cumulative assessment groups (effects on thyroid) [link](#)
- EFSA Sept 2019: Establishment of (CAGs) cumulative assessment groups (effects on the nervous system) [link](#)
- EFSA Jan 2014: Outcome of the public consultation on the Scientific Opinion on the identification of pesticides to be included in cumulative assessment groups (CAGs) on the basis of their toxicological profile [link](#)
- EFSA Dec 2013: Scientific Opinion on the relevance of dissimilar mode of action and its appropriate application for cumulative risk assessment of pesticides residues in food [link](#)
- EFSA Sept 2009: Scientific Opinion on Risk Assessment for a Selected Group of Pesticides from the Triazole Group to Test Possible Methodologies to Assess Cumulative Effects from Exposure through Food from these Pesticides on Human Health [link](#)
- EFSA May 2008: Opinion of the Scientific Panel on Plant Protection products and their Residues to evaluate the suitability of existing methodologies and, if appropriate, the identification of new approaches to assess cumulative and synergistic risks from pesticides to human health with a view to set MRLs for those pesticides in the frame of Regulation (EC) 396/2005 [link](#)

Further publications on topics related to consumer risk assessment that are under development are as follows:

- UK Committee on Carcinogenicity (2019) guidance note (COC Guidance Statement G09) on LTL exposure assessment. [COC 2019 LTL Guidance](#)
- EFSA: Update: use of the benchmark dose approach in risk assessment (2016) [BMDL link](#)
- WHO guidance on genotoxicity (2020). [EHC 240 \(updated 2nd Ed\) genotoxicity](#)

HSE (UK) is participating in a number of international initiatives related to residues and risk assessment (OECD Working group on residue definitions, and the ongoing JMPR programme of evaluation work/attendance at CCPR (CODEX) and participating in JMPR/CCPR discussions of a technical nature on general considerations for risk assessment.

Further advances in risk assessment methodology will be taken into account in developing the approach to risk assessments in the future.

## Risk Assessment - dietary intake assessments

The screening assessment uses the internationally agreed approach to long-term (chronic) and short-term (acute) consumer exposure assessment with UK food consumption data as detailed within the UK NEDI and NESTI models which are available on the [HSE website](#).

Screening assessments have been done for all pesticides to check that predicted intakes are within the relevant health-based reference values. A short-term (acute) exposure assessment is not done for pesticides which are not acutely toxic where it has been established that an ARfD is not required. EU toxicological endpoints can be found in the [EU Pesticides database](#).

Toxicological reference values set by the JMPR (The Joint FAO/WHO Meeting on Pesticide Residues) can be found in individual pesticide evaluations at [JMPR Evaluations](#) (an up to date index to pesticide evaluations is available in the latest report). (an up to date index to pesticide evaluations is available in the latest report).

The screening assessment uses the internationally agreed approach to long-term (chronic) and short-term (acute) consumer exposure assessment with UK food consumption data as detailed within the UK NEDI and NESTI models which are available on the [HSE website](#).

For the Q4 (2021) assessments, the following approaches have been taken to refine these assessments according to case-by-case issues and to ensure that appropriate consumption values are used for less frequently consumed commodities where available food consumption data may be limited:

- Data on asparagus were used for adults and vegetarians and data on broccoli were used for the other groups for asparagus; this was considered reasonable after taking account of the low numbers of consumers in the survey data for other groups and the comparison to other data.
- Data on aubergine were used, although there are a low number of consumers in several of the sub-groups for aubergine. However, use of these consumption data was considered reasonable after comparison with alternative data.
- Data on banana were used for plantain
- Data on beans with pods were used for all forms of green beans, including speciality beans
- Data on meat (excluding poultry and offal) were used for all forms of beef
- Data on fish were used for all forms of white fish
- Data on melon were used for all forms of melon and watermelon
- Data on blackberries and raspberries were used for screening assessment of raspberries, blackberries, and blueberries; use of these consumption data was considered reasonable after comparison with alternative data.

- Data on cheese were used for all forms of cheese
- Data on oil was used for all types of olive oil
- Consumption values were provided by the Food Standards Agency for grapefruit to identify the various contributions of grapefruit in the diet. The consumption values used here (and as used in 2018 and 2016) cover consumption for fresh grapefruit excluding juice, canned grapefruit and grapefruit consumed in other recipe forms. For the current assessments, the consumer groups covered represent infants (4 to 18 months old), toddlers (1.5 to 3 years), 4-6 year olds, 7-10 year olds, 11-14 year olds, 15-18 year olds, and elderly. These are slightly different to the groups used previously. Vegetarians, elderly residential and elderly in their own home are not reported separately. For some of the consumer groups, the number of grapefruit consumers in the surveys (from years 2008 to 2012) are very low e.g., infants which is not unexpected. The highest number of young consumers out of all the young consumer groups was three persons for infants (3 out of 2863 infant consumers). These data are used as they represent the best available data. Adults and elderly consume fresh grapefruit at a higher frequency (a higher number of consumers) than the other consumer groups, and of these adults is the most critical consumer group.
- Data on cabbage were used for samples of whole spring greens and kale. For samples of sliced or shredded kale or greens data on cabbage without the use of a variability factor were used. As pieces of sliced/shredded kale or greens are small, a whole vegetable consideration which takes account of unit to unit variability is not considered relevant.
- For large speciality mushrooms (large flat mushrooms), data on mushrooms with a variability factor of 7 and a unit weight of 85 g were used. As per ordinary mushrooms, no variability factor was used for the smaller button mushrooms.
- Data on bread were used for all forms of bread, including speciality breads.
- Data on flour were used for all forms of wheat flour. These consumption values were provided by the Food Standards Agency; the consumer groups covered represent infants (6 - 12 months), toddlers (1.5 to 3 years), 4-6 year olds, 7-10 year olds, 11-14 year olds, 15-18 year olds, adults, vegetarians and elderly [elderly residential and elderly in their own home are not reported separately].
- Data on peanuts were used for all nuts; use of these consumption data was considered reasonable after comparison with data for individual types of nuts which had low numbers of consumers for some groups.
- Data on peppers were used for all types of processed peppers.
- Specific infant consumption data on infant food were used.

## Plantain (Banana)

Crop	Pesticide	Highest residue (mg/kg)	Adult	Intake (mg/kg bw/day) Critical group <sup>†</sup>	ARfD (mg/kg bw)	Source
Plantain (Banana)	Chlorpyrifos	0.05	0.00061	0.0042 (infant) 0.0028 (toddler) 0.0021 (4-6 year old) 0.0015 (7-10 year old) 0.00087 (11-14 year old) 0.00074 (15-18 year old) 0.00071 (vegetarian) 0.00064 (elderly – residential) 0.00061 (adult) 0.00056 (elderly – own home)	Not established	EU, 2019

### Comment on risk assessment

EFSA (2019)<sup>9</sup> has indicated that no toxicological reference values could be determined for chlorpyrifos, due to concerns over genotoxicity. Additionally, EFSA raised concerns over neurological effects in the developing foetus and young child. Chlorpyrifos is not approved in the EU and UK and pesticide products containing chlorpyrifos were withdrawn in 2020.

HSE considers that for short-term risk assessment, an indicative toxicological reference value of 0.0003 mg/kg bw can be used based on the LOAEL set by EFSA for a developmental neurotoxicity study and applying a safety factor of 1000 to account for the severe nature of the findings (effects on brain measurements in a developmental neurotoxicity study). Toxicologists usually use

<sup>9</sup> [EFSA 2019 statement on human health- chlorpyrifos](#)

safety factors of between 100 and a 1000 when a NOAEL cannot be determined within a study. HSE proposed indicative toxicological reference value is conservative as it uses the highest uncertainty factor applied by toxicologists and is based on a LOAEL from a study with repeated dosing. Overall, HSE's approach is considered precautionary in protecting the nervous system in the developing foetus and child.

### **Plantain (Banana) flesh after peeling**

The dietary intakes calculated for when the peel is removed prior to consumption, indicate that there are no exceedances of indicative toxicological reference value and, in this case, a short term effect on health is not expected.

The below risk assessment only applies if all of the peel is consumed. This is because it has been reported that only 2% of the residue of chlorpyrifos remains (EFSA, 2017) in the flesh when the fruit is peeled.

### **Whole plantain (banana), including all the peel**

We cannot conclude whether residues of chlorpyrifos at this level would have any effect on health. As well as the precautionary nature of the toxicological assessment (see above), any effect on health would depend on a number of factors which would need to come together at the same time (the high residue found in the sample being consumed by the most critical consumer infant (0.05 mg/kg of chlorpyrifos in plantain/bananas), high residue in single fruit item, peak consumption levels (97.5th percentile), and a large proportion of peel from the fruit being consumed).

The following risk assessment, assuming all of the peel is consumed, is presented, although the PRiF consider this to be a 'worst case' form of assessment for the reasons explained above:

The intakes for all of the consumer groups exceeded the HSE proposed indicative toxicological reference value. The highest intake was for infants.

If infants ate large portions of plantain (banana), including the peel, containing chlorpyrifos at 0.05 mg/kg their intake could be around 1400 % of the above mentioned HSE proposed indicative toxicological reference value for short term exposure. This intake is approximately 70 times lower than the lowest intake in repeat-dose animal studies which was reported to cause effects in a developmental neurotoxicity study where pregnant rats were dosed from day 6 of pregnancy through until the pups were 11 days old.

Toxicologists usually apply a factor of between 100 and 1000 to this dose to take into account the uncertainties caused by using animal data and possible differences in susceptibility between people. Given the nature of the findings, HSE took a precautionary approach and applied a factor of 1000. We consider this significant reduction in the uncertainty factor from 1000 to 70 undesirable. The developmental neurotoxicity study in which the effects on the brain measurements were observed reported no behavioural or developmental deficits, and there is an indication that the changes in brain measurement might be reversible. The interpretation of this study by regulatory assessors is uncertain, and, despite the precautionary nature of HSE's assessment, it is

not possible to conclude on whether there might be any adverse short term health effects after eating large portions (97.5th percentile consumption) of plantain (banana), including the peel, containing the level found in this report.

This estimate assumes that peel of plantain/banana is consumed. However, if the peel is not consumed then intakes in all groups are within the indicative reference value and a short-term effect on health is not expected.

Please refer to the section below on [‘Substances that might be genotoxic’](#) for HSE’s conclusions regarding potential genotoxicity.

### **Banana**

<b>Crop</b>	<b>Pesticide</b>	<b>Highest residue (mg/kg)</b>	<b>Adult</b>	<b>Intake (mg/kg bw/day) Critical group<sup>†</sup></b>	<b>ARfD (mg/kg bw)</b>	<b>Source</b>
Banana	Thiabendazole	1.4	0.017	0.12 (infant)	0.1	EU, 2017

### **Comment on risk assessment**

#### **Banana flesh after peeling**

The dietary intakes calculated for when the peel is removed prior to consumption, indicate that there are no exceedances of the ARfD and, in this case, an effect on health is not expected. This is in line with the risk assessment performed when the MRL was established.

The below risk assessment only applies if all of the peel is consumed. This is because it has been reported that only 3 % of the residue of thiabendazole remains (EFSA, 2016) in the flesh when the fruit is peeled.

#### **Whole banana, including all the peel**

We consider that an effect on health would be unlikely. Any effect on health would depend on a number of factors which would need to come together at the same time (the high residue found in the sample being consumed by the most critical consumer infant (1.4 mg/kg of thiabendazole in bananas), high residue in single fruit item, peak consumption levels (97.5th percentile), and a large proportion of peel from the fruit being consumed).

The following risk assessment assuming all of the peel is consumed, is presented, although the PRiF consider this to be a ‘worst case’ form of assessment for the reasons explained above:

The intakes for infants exceeded the ARfD. Intakes for all other consumer groups were below the reference dose. If infants ate large portions of bananas (including the peel) containing thiabendazole at 1.4 mg/kg, their intake of thiabendazole could be 117



% of the Acute Reference Dose. This intake is 80 times lower than a dose which caused no observed adverse effect in a developmental study in rats over 11 days. The European Food Safety Authority used this study as the basis of the ARfD. Toxicologists usually apply a factor of 100 to this dose to take into account uncertainties caused by using animal data and possible differences in susceptibility between people. We consider the reduced factor of 80 still sufficient to make an effect on health unlikely.

This estimate assumes that peel of bananas is consumed. However, if the peel is not consumed then the risk assessment that is the basis for the MRL applies, intakes in all groups are within the ARfD and an effect on health is not expected.

Overall, HSE concludes that a short-term effect on health would be unlikely in the case of consumption including all of the peel, and not expected in the case of peeling banana prior to consumption.

#### Beans with pods

Crop	Pesticide	Highest residue (mg/kg)	Adult	Intake (mg/kg bw/day) Critical group <sup>†</sup>	ARfD (mg/kg bw)	Source
Beans with pods (Guar beans)	Chlorpyrifos	0.03	0.000069	0.00015 (infant) 0.00015 (toddler) 0.00011 (4-6 year old child) 0.000083 (vegetarian) 0.000082 (15-18 year old child) 0.000069 (adult) 0.000065 (elderly – own home) 0.000061 (7-10 year old child) 0.000059 (11-14 year old child) 0.000033 (elderly – residential)	No toxicological reference values established	EU, 2019

## Comment on risk assessment

EFSA (2019)<sup>10</sup> has indicated that no toxicological reference values could be determined for chlorpyrifos, due to concerns over genotoxicity. Additionally, EFSA raised concerns over neurological effects in the developing foetus and young child. Chlorpyrifos is not approved in the EU and UK and pesticide products containing chlorpyrifos were withdrawn in 2020.

HSE considers that for short-term risk assessment, an indicative toxicological reference value of 0.0003 mg/kg bw can be used based on the LOAEL set by EFSA for a developmental neurotoxicity study and applying a safety factor of 1000 to account for the severe nature of the findings (effects on brain measurements in a developmental neurotoxicity study). Toxicologists usually use safety factors of between 100 and a 1000 when a NOAEL cannot be determined within a study. The HSE proposed indicative toxicological reference value is conservative as it uses the highest uncertainty factor applied by toxicologists and is based on a LOAEL from a study with repeated dosing. Overall, the HSE approach is considered precautionary in protecting the nervous system in the developing foetus and child.

None of the intakes exceeded the HSE proposed indicative toxicological reference value for short term assessment. Based on the low intakes, HSE concludes that a short term effect on health is not expected.

Please refer to the section below on [‘Substances that might be genotoxic’](#) for HSE’s conclusions regarding potential genotoxicity

### Beans with pods

Crop	Pesticide	Highest residue (mg/kg)	Adult	Intake (mg/kg bw/day) Critical group <sup>†</sup>	ARfD (mg/kg bw)	Source
Beans with pods (Yard long beans)	Dimethoate and Omethoate	0.4 (D: dimethoate) and	D: 0.00092	0.0020 (infant) 0.0020 (toddler) 0.0015 (4-6 year old child) 0.0011 (vegetarian) 0.0011 (15-18 year old child) 0.00092 (adult) 0.00087 (elderly – own home) 0.00081 (7-10 year old child)	Not established	EU, 2019

<sup>10</sup> [EFSA 2019 statement on human health- chlorpyrifos](#)

		0.2 (O: omethoate)	O: 0.00046	0.00078 (11-14 year old child) 0.00044 (elderly – residential) 0.0010 (infant) 0.0010 (toddler) 0.00075 (4-6 year old child) 0.00056 (vegetarian) 0.00055 (15-18 year old child) 0.00046 (adult) 0.00043 (elderly – own home) 0.00040 (7-10 year old child) 0.00039 (11-14 year old child) 0.00022 (elderly – residential)	Not established	EU, 2019
--	--	--------------------	------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------	----------

### Comment on risk assessment

The EFSA Conclusion (2018) for dimethoate has indicated that no toxicological reference values could be determined for dimethoate and its metabolite omethoate, due to a lack of a fully supporting toxicological database. Neither dimethoate nor omethoate is approved in the EU and the UK, and pesticide products containing dimethoate were withdrawn in the EU and UK in 2020.

For dimethoate, EFSA (2018) stated an indicative value for a hypothetical toxicological reference value for short term exposure of 0.0001 mg/kg bw. Using this indicative value, estimated dietary intakes for all consumer groups exceed this reference value. The intakes of omethoate for all consumer groups exceeded this hypothetical short term toxicological reference value for dimethoate. The highest intake was infants and toddlers.

If infants and toddlers ate large portions of beans with pods containing dimethoate at 0.4 mg/kg their intake could be 2000 % of the above mentioned hypothetical toxicological reference value. If infants and toddlers ate large portions of beans with pods containing omethoate at 0.2 mg/kg their intake could be 1000 % of this hypothetical toxicological reference value for dimethoate. This indicative toxicological reference value is a precautionary value intended to protect the nervous system in the developing foetus and child, which has been set well below intakes which caused no observed effects in animal studies. The JMPR

(September, 2019) established an ARfD for dimethoate of 0.02 mg/kg bw; this supports the view that the proposed hypothetical reference value from the EFSA Conclusion is precautionary.

These exposures are undesirable, but it is not clear if they may cause any adverse effect. The estimated exposures are unlikely to inhibit acetylcholinesterase<sup>11</sup>, the basis of previous evaluations of the safety of dimethoate and omethoate. Based on this assessment, HSE concludes that a short term effect on health is unlikely after eating large portions (97.5th percentile consumption) of beans with pods containing the levels found in this report.

Please refer to the section below on [“Substances that might be genotoxic”](#) for HSE’s conclusions regarding potential genotoxicity.

### **Grapefruit**

<b>Crop</b>	<b>Pesticide</b>	<b>Highest residue (mg/kg)</b>	<b>Adult</b>	<b>Intake (mg/kg bw/day) Critical group<sup>†</sup></b>	<b>ARfD (mg/kg bw)</b>	<b>Source</b>
Grapefruit	Imazalil	2.7	0.039	0.15 (infant)	General population 0.1 Pregnant and nursing females 0.05	EFSA, 2007

### **Comment on risk assessment**

#### **Grapefruit flesh after peeling**

The dietary intakes calculated for when the peel is removed prior to consumption, indicate that there are no exceedances of the ARfD and, in this case, an effect on health is not expected. This is in line with the risk assessment performed when the MRL was established.

The below risk assessment only applies if all of the peel is consumed. This is because it has been reported that only 7 % of the residue of imazalil remains (EFSA, 2018) in the flesh when the fruit is peeled.

---

<sup>11</sup> This enzyme, acetylcholinesterase, is included in the Glossary on page 160

## **Whole grapefruit, including all the peel**

### *Pregnant and nursing females*

The intakes for adults, 11-14 year old children, 15-18 year old children and vegetarians are all below the ARfD of 0.05 mg/kg bw for pregnant and nursing females.

### *General population*

We consider that an effect on health would be unlikely. Any effect on health would depend on a number of factors which would need to come together at the same time (the high residue found in the sample being consumed by the most critical consumer infants (2.7 mg/kg of imazalil in grapefruit), high residue in single fruit item, peak consumption levels (97.5th percentile), and a large proportion of peel from the fruit being consumed).

The following risk assessment assuming all of the peel is consumed, is presented, although the PRiF consider this to be a 'worst case' form of assessment for the reasons explained above:

The intakes for infants exceeded the ARfD for the general population. If infants ate large portions of grapefruit containing imazalil at 2.7 mg/kg, their intake of imazalil could be 147% of the Acute Reference Dose. This intake is 67 times lower than a dose which caused no observed adverse effect in a rabbit developmental study, used as the basis of the ARfD (the ARfD is based on a NOAEL of 10 mg/kg bw for reduced bodyweight gain and food consumption in dams). The European Food Safety Authority used this study as the basis of the ARfD. Toxicologists usually apply a factor of 100 to this dose to take into account uncertainties caused by using animal data and possible differences in susceptibility between people. We consider the reduced factor of 67 still enough to account for these uncertainties, also noting that an ARfD based on maternal toxicity in a developmental study with repeated dosing (13 days) is likely to be very protective for the general population. Based on this assessment an effect on health is unlikely.

This estimate assumes that peel of grapefruit is consumed. However, if the peel is not consumed then the risk assessment that is the basis for the MRL applies, intakes in all groups are within the ARfD and an effect on health is not expected.

### **Grapefruit**

<b>Crop</b>	<b>Pesticide</b>	<b>Highest residue (mg/kg)</b>	<b>Adult</b>	<b>Intake (mg/kg bw/day) Critical group<sup>†</sup></b>	<b>ARfD (mg/kg bw)</b>	<b>Source</b>
Grapefruit	Thiabendazole	3.3	0.048	0.18 (infant)	0.1	EU, 2017

## Comment on risk assessment

### Grapefruit flesh after peeling

The dietary intakes calculated for when the peel is removed prior to consumption, indicate that there are no exceedances of the ARfD and, in this case, an effect on health is not expected. This is in line with the risk assessment performed when the MRL was established.

The below risk assessment only applies if all of the peel is consumed. This is because it has been reported that only 5% of the residue of thiabendazole remains (EFSA, 2021) in the flesh when the fruit is peeled.

### Whole grapefruit, including all the peel

We consider that an effect on health would be unlikely. Any effect on health would depend on a number of factors which would need to come together at the same time (the high residue found in the sample being consumed by the most critical consumer infants (3.3 mg/kg of thiabendazole in grapefruit, high residue in single fruit item, peak consumption levels (97.5th percentile), and a large proportion of peel from the fruit being consumed).

The following risk assessment assuming all of the peel is consumed, is presented, although the PRiF consider this to be a 'worst case' form of assessment for the reasons explained above:

The intakes for infants exceeded the ARfD. If infants ate large portions of grapefruit containing thiabendazole at 3.3 mg/kg, their intake of thiabendazole could be 180% of the Acute Reference Dose. This intake is 56 times lower than a dose which caused no observed adverse effects in a developmental study in rats over 11 days. The European Food Safety Authority used this study as the basis of the ARfD. Toxicologists usually apply a factor of 100 to this dose to take into account the uncertainties caused by using animal data and possible differences in susceptibility between people. We consider the reduced factor of 56 still sufficient to make an effect on health unlikely.

This estimate assumes that peel of grapefruit is consumed. However, if the peel is not consumed then the risk assessment that is the basis for the MRL applies, intakes in all groups are within the ARfD and an effect on health is not expected.

### Grapes

Crop	Pesticide	Highest residue (mg/kg)	Adult	Intake (mg/kg bw/day) Critical group <sup>†</sup>	ARfD (mg/kg bw)	Source
Grapes	Ethephon	0.9	0.018	0.055 (toddler)	0.05	EU, 2008

## Comment on risk assessment

The intakes for toddlers exceeded the ARfD. If toddlers ate large portions of grapes containing ethephon at 0.9 mg/kg, their intake of ethephon could be 110% of the Acute Reference Dose. This intake is 109 times lower than a dose which caused no observed adverse effect in a 28 day oral dog study. The European Food Safety Authority used this study as the basis of the ARfD.

Toxicologists usually apply a factor of 100 to this dose to take into account uncertainties caused by using animal data and possible differences in susceptibility between people. However, in this case the factor was larger (120) to ensure consistency with the findings of human volunteer studies. We consider the reduced factor of 109 (from 120) still sufficient to make an effect on health unlikely. More detail on the factors applied is on page 145 of this report.

Based on this assessment, we conclude that a short term effect on health is unlikely.

## Infant food

Crop	Pesticide	Highest residue (mg/kg)	Adult	Intake (mg/kg bw/day) Critical group <sup>†</sup>	ARfD (mg/kg bw)	Source
Infant food	Chlorate	0.5	N/A	0.043	0.036	EFSA, 2015

## Comment on risk assessment

The intakes for infants exceeded the ARfD. This assessment is for peak daily amounts of infant food consumed (all types of infant foods consumed throughout the day). If infants ate large portions of infant food containing chlorate at 0.5 mg/kg, their intake of chlorate could be 119% of the Acute Reference Dose. This intake is 1.2 times higher than a dose which caused no observed adverse effect in a controlled clinical study with daily dosing for 12 weeks. The European Food Safety Authority used this valid human study as the basis of the ARfD; an uncertainty factor was not considered necessary to take into account uncertainties caused by possible differences in susceptibility between people. In this study, there were no changes in the extensive battery of biochemical and physiological parameters monitored in any of the people from which clinical observations were made at the tested dose. Based on the residue intake representing a small (19%) exceedance of the dose tested in the clinical trial, HSE concludes that an effect on health is unlikely.

This conclusion is based on assessment of peak daily amounts of infant food consumed (all types of infant foods consumed throughout the day). However, it is noted that the highest residue of chlorate in infant food in this report (0.5 mg/kg) was found in jars of creamed porridge. The porridge is marketed as a breakfast cereal. If the porridge containing chlorate at 0.5 mg/kg accounted for half an infant's daily diet, then there would be no exceedance of the ARfD and an effect on health would not be expected. If it accounted for the entire daily diet, then the ARfD would be exceeded but an effect on health would still be unlikely.

Chlorate was found in 13 of the 46 infant food samples, and aside from the highest residue of chlorate of 0.5 mg/kg assessed here, the next highest residue of chlorate was 0.03 mg/kg. A chlorate residue at this level (0.03 mg/kg) would lead to dietary intakes within the ARfD, and an effect on health would not be expected.

### Spring Greens & Kale

Crop	Pesticide	Highest residue (mg/kg)	Adult	Intake (mg/kg bw/day) Critical group <sup>†</sup>	ARfD (mg/kg bw)	Source
Kale	Cypermethrin (alpha-cypermethrin)	0.04	0.00049	0.0017 (infants) 0.0013 (4-6 year old)	0.00125	EU, 2019

### Comment on risk assessment

The risk assessment has been conducted assuming the residue of cypermethrin is derived from use of alpha-cypermethrin. Toxicological studies suggest that alpha-cypermethrin is the more toxic isomeric form and there is a specific ARfD for alpha-cypermethrin. The analytical data (chromatograms) supports the understanding that the residue found comes from use of alpha-cypermethrin; this also fits with the authorised uses on kale (alpha-cypermethrin is authorised for use on this crop and not cypermethrin).

Cypermethrin is an insecticide which is available in different isomeric forms. We understand that the application made to the crop was alpha-cypermethrin, so we have conducted the assessment based on alpha-cypermethrin, which is the more toxic form.

The intakes for infants and 4-6 year old children exceed the ARfD. The highest intake is for infants.

If infants ate large portions of kale containing alpha-cypermethrin at 0.04 mg/kg, their intake of alpha-cypermethrin could be 138% of the Acute Reference Dose of 0.00125 mg/kg bw. This intake is 150 times lower than a dose which caused minor adverse effects in a developmental neurotoxicity study in rats. The European Food Safety Authority used this study as the basis of the ARfD. Toxicologists usually apply a factor of 100 to this dose to take into account uncertainties caused by using animal data and possible differences in susceptibility between people. The factor used for alpha-cypermethrin was two-fold greater (200) to account for the absence of a dose level in the study where no adverse effects were seen. We consider the reduced factor of 150 (from 200) still sufficient to make an effect on health unlikely.

### Melon

Crop	Pesticide	Highest residue (mg/kg)	Adult	Intake (mg/kg bw/day) Critical group <sup>†</sup>	ARfD (mg/kg bw)	Source
------	-----------	-------------------------	-------	---------------------------------------------------	-----------------	--------



Melon	Imazalil	1.1	0.041	0.053 (11-14 year old child)	General population 0.1 Pregnant and nursing females 0.05	EFSA, 2007
-------	----------	-----	-------	------------------------------	-------------------------------------------------------------------	------------

### **Comment on risk assessment**

#### **Melon flesh after peeling**

The dietary intakes calculated for when the peel is removed prior to consumption, indicate that there are no exceedances of the ARfD and, in this case, an effect on health is not expected. This is in line with the risk assessment performed when the MRL was established.

The below risk assessment only applies if all of the peel is consumed. This is because it has been reported that only 12% of the residue of imazalil remains (EFSA, 2017) in the flesh when the fruit is peeled.

#### **Whole melon, including all the peel**

We consider that an effect on health would be unlikely. Any effect on health would depend on a number of factors which would need to come together at the same time (the high residue found in the sample being consumed by the most critical consumer 11-14 year old child (1.1 mg/kg of imazalil in melon), high residue in single fruit item, peak consumption levels (97.5th percentile), and a large proportion of peel from the fruit being consumed).

The following risk assessment assuming all of the peel is consumed, is presented, although the PRiF consider this to be a 'worst case' form of assessment for the reasons explained above:

### *Pregnant and nursing females*

The intakes for 11-14 year old children exceeded the ARfD. If 11-14 year old children ate large portions of melon containing imazalil at 1.1 mg/kg, their intake of imazalil could be 107% of the Acute Reference Dose. This intake is 94 times lower than a dose which caused no observed adverse effect in a 13 day repeat dose rabbit developmental study (the ARfD is based on a NOAEL of 5 mg/kg bw for foetal toxicity (increased resorptions; a marker of early foetal deaths)). The European Food Safety Authority used this study as the basis of the ARfD. Toxicologists usually apply a factor of 100 to this dose to take into account uncertainties caused by using animal data and possible differences in susceptibility between people. We consider the reduced factor of 94 still sufficient to make an effect on health unlikely.

### *General population*

The intakes for all consumer groups are below the ARfD of 0.1 mg/kg bw for general population.

These estimates assume that peel of melon is consumed. However, if the peel is not consumed then the risk assessment that is the basis for the MRL applies, intakes in all groups are within the ARfDs and an effect on health is not expected.

### **Olive oil**

<b>Crop</b>	<b>Pesticide</b>	<b>Highest residue (mg/kg)</b>	<b>Adult</b>	<b>Intake (mg/kg bw/day) Critical group<sup>†</sup></b>	<b>ARfD (mg/kg bw)</b>	<b>Source</b>
Olive oil	Chlorpyrifos	0.003	0.000004	0.000009 (4-6 year old child) 0.000009 (toddler) 0.000008 (infant) 0.000007 (7-10 year old child) 0.000006 (vegetarian) 0.000005 (11-14 year old child) 0.000005 (15-18 year old child) 0.000004 (adult) 0.000004 (elderly – residential) 0.000003 (elderly – own home)	No toxicological reference values established	EU, 2019

## Comment on risk assessment

EFSA (2019)<sup>12</sup> has indicated that no toxicological reference values could be determined for chlorpyrifos, due to concerns over genotoxicity. Additionally, EFSA raised concerns over neurological effects in the developing foetus and young child. Chlorpyrifos is not approved in the EU and UK and pesticide products containing chlorpyrifos were withdrawn in 2020.

HSE considers that for short-term risk assessment, an indicative toxicological reference value of 0.0003 mg/kg bw can be used based on the LOAEL set by EFSA for a developmental neurotoxicity study and applying a safety factor of 1000 to account for the severe nature of the findings (effects on brain measurements in a developmental neurotoxicity study). Toxicologists usually use safety factors of between 100 and a 1000 when a NOAEL cannot be determined within a study. The HSE proposed indicative toxicological reference value is conservative as it uses the highest uncertainty factor applied by toxicologists and is based on a LOAEL from a study with repeated dosing. Overall, the HSE approach is considered precautionary in protecting the nervous system in the developing foetus and child.

None of the intakes exceeded the HSE proposed indicative toxicological reference value for short term assessment. Based on the low intakes, HSE concludes that a short term effect on health is not expected.

Please refer to the section below on [“Substances that might be genotoxic”](#) for HSE’s conclusions regarding potential genotoxicity.

### Olive oil

Crop	Pesticide	Highest residue (mg/kg)	Adult	Intake (mg/kg bw/day) Critical group <sup>†</sup>	ARfD (mg/kg bw)	Source
Olive oil	Dimethoate	0.007	0.000009	0.000022 (4-6 year old child) 0.000020 (toddler) 0.000018 (infant) 0.000017 (7-10 year old child) 0.000015 (vegetarian) 0.000012 (11-14 year old child) 0.000011 (15-18 year old child)	Not established	EU, 2019

<sup>12</sup> [EFSA 2019 statement on human health- chlorpyrifos](#)

				0.000009 (adult)		
				0.000008 (elderly – residential)		
				0.000007 (elderly – own home)		

### Comment on risk assessment

The EFSA Conclusion (2018) for dimethoate has indicated that no toxicological reference values could be determined for dimethoate due to the lack of a fully supporting toxicological database. Dimethoate is not approved in the EU and the UK, and pesticide products containing dimethoate were withdrawn in the EU and UK in 2020.

For dimethoate, EFSA (2018) stated an indicative value for a hypothetical toxicological reference value for short-term exposure of 0.0001 mg/kg bw. Using this indicative value, estimated dietary intakes of dimethoate do not exceed this reference value for any consumer group. This indicative toxicological reference value is a precautionary value intended to protect the nervous system in the developing foetus and child, which has been set well below intakes which caused no observed effects in animal studies. The JMPR (September 2019) established an ARfD for dimethoate of 0.02 mg/kg bw; this supports the view that the proposed hypothetical reference value from the EFSA Conclusion is precautionary. Based on the low short-term intakes, HSE concludes that a short-term effect on health is not expected.

Please refer to the section below on [“Substances that might be genotoxic”](#) for HSE’s conclusions regarding potential genotoxicity

### Processed pepper

Crop	Pesticide	Highest residue (mg/kg)	Adult	Intake (mg/kg bw/day) Critical group <sup>†</sup>	ARfD (mg/kg bw)	Source
Processed pepper	Chlorpyrifos	0.003	0.000040	0.000049 (7-10 year old child) 0.000049 (vegetarian) 0.000049 (toddler) 0.000040 (adult) 0.000035 (4-6 year old child) 0.000032 (11-14 year old child) 0.000030 (15-18 year old child) 0.000029 (elderly – own home) 0.000016 (elderly – residential)	No toxicological reference values established	EU, 2019

## Comment on risk assessment

EFSA (2019)<sup>13</sup> has indicated that no toxicological reference values could be determined for chlorpyrifos, due to concerns over genotoxicity. Additionally, EFSA raised concerns over neurological effects in the developing foetus and young child. Chlorpyrifos is not approved in the EU and UK, and pesticide products containing chlorpyrifos were withdrawn in 2020.

HSE considers that for short-term risk assessment, an indicative toxicological reference value of 0.0003 mg/kg bw can be used based on the LOAEL set by EFSA for a developmental neurotoxicity study and applying a safety factor of 1000 to account for the severe nature of the findings (effects on brain measurements in a developmental neurotoxicity study). Toxicologists usually use safety factors of between 100 and a 1000 when a NOAEL cannot be determined within a study. The HSE proposed indicative toxicological reference value is conservative as it uses the highest uncertainty factor applied by toxicologists and is based on a LOAEL from a study with repeated dosing. Overall, the HSE approach is considered precautionary in protecting the nervous system in the developing foetus and child.

None of the intakes exceeded the HSE proposed indicative toxicological reference value for short term assessment. Based on the low intakes, HSE concludes that a short term effect on health is not expected.

Please refer to the section below on [‘Substances that might be genotoxic’](#) for HSE’s conclusions regarding potential genotoxicity.

---

<sup>13</sup> [EFSA 2019 statement on human health- chlorpyrifos](#)

## Short-term dietary risk assessment – multiple assessments needed following screening assessment of samples

Samples which contain more than one pesticide from the groups we consider (samples containing more than one organophosphorus/carbamate or captan/folpet or DDAC/BAC or mepiquat/chlormequat or triazoles] and where a more detailed assessment was needed following screening.

Crop/Critical group	Pesticide	Residue mg/kg	Intake			ARfD	Source	
			mg/kg bw	%ARfD				
Beans (with pods) Infant	dimethoate	0.4	0.0020	-	}	Total	Not established	EU, 2019
	omethoate	0.2	0.0010	-			Not established	EU, 2019
	profenofos	0.3	0.0015	0.2	}	-	1	JMPR, 2007

### Comment on risk assessment:

The presence of profenofos in the sample does not significantly contribute to the overall combined intake when compared to the presence of dimethoate and its metabolite omethoate alone. The overall risk is covered by the single substance risk assessment for beans with pods (see the above details for HSE's single substance assessment for beans with pods for this residue finding of dimethoate [0.4 mg/kg] and omethoate [0.2 mg/kg]).

Overall, HSE concludes that, when taking into account the known effect that each of dimethoate, omethoate and profenofos can have on acetyl cholinesterase, the presence of these pesticides in the same sample is unlikely to inhibit acetyl cholinesterase.

HSE concludes that a short term effect on health for the combined residues of dimethoate, omethoate and profenofos in this sample of beans with pods (Yard long beans) is unlikely.

Please refer to the section below on "[Substances that might be genotoxic](#)" for HSE's conclusions regarding potential genotoxicity

Crop/Critical group	Pesticide	Residue mg/kg	Intake			ARfD	Source	
			mg/kg bw	%ARfD				
Beans (with pods) Infant	chlorpyrifos	0.03	0.00015	-	}	Total	Not established	EU, 2019
	monocrotophos	0.004	0.000020	1.0			0.002	JMPR, 1995
	omethoate	0.003	0.000015	-	}	-	Not established	EU, 2019

### **Comment on risk assessment:**

These highest calculated dietary intakes are low. The estimated highest intake of chlorpyrifos represents around 50% of the above noted indicative toxicological reference value of 0.0003 mg/kg bw for short-term assessment (see explanation above of this indicative value in the single substance assessment for beans with pods-chlorpyrifos above). The estimated highest intake of omethoate represents around 15% of the hypothetical short-term reference value for dimethoate proposed by EFSA (EFSA Conclusion for dimethoate, 2018). Furthermore, the presence of monocrotophos in the sample does not significantly contribute to the overall combined intake. At these low levels, these pesticides together will be unlikely to inhibit acetyl cholinesterase, the known effect from exposure to each of these residues.

Overall, HSE concludes that, when taking into account the known effect that each of chlorpyrifos, monocrotophos and omethoate can have on acetyl cholinesterase, the presence of these pesticides in the same sample is unlikely to inhibit acetyl cholinesterase.

HSE concludes that a short term effect on health for the combined residues of chlorpyrifos, monocrotophos and omethoate in this sample of beans with pods (Guar beans) is unlikely.

Please refer to the section below on [‘Substances that might be genotoxic’](#) for HSE’s conclusions regarding potential genotoxicity.

## **Long-term dietary risk assessments needed following screening assessment of samples**

As noted in [section 3](#) total long-term dietary assessments across all commodities are not performed for these quarterly assessments. The issue is more fully considered in regulatory contexts pre-authorisation and at the time of MRL review. Then the issue is considered across all commodities (so more precautionary) by pesticide levels determined in GAP compliant trials, intended to address highest likely residues that might arise following pesticide use according to label recommendations

However, for the PRiF quarterly assessments, HSE do perform a screening exercise for all of the residues found for an individual commodity to see if the long-term intakes (commodity by commodity) show any indication of exceedance of the ADI. If an exceedance was observed, then HSE would consider further and we would present a more detailed risk assessment.

In HSE's long-term exposure screening assessment for this report NI and GB samples were combined.

None of these individual commodity long-term exposure screening assessments performed in this quarter (for each of the pesticides found in this report) indicated potential for adverse long-term health effects. HSE assessed the dietary intakes to be below the ADI or other established long-term health-based reference value.

## **Substances that might be genotoxic (see explanation in the section on HSE's assessment of risk)**

During regulatory assessment, careful consideration is given to any pesticides that may exhibit any potential to be genotoxic (able to damage genetic material) in live animals, so we need to consider the significance to the consumer when these residues are found. There are small number of examples of older pesticides that might be genotoxic, where modern data to investigate the true genotoxic potential is not expected to be made available. It is likely that these will only be found in imported foods. For many of these old pesticides, the toxicological reference doses are low and PRiF uses low reporting limits to ensure that these residues are found even at very low levels, as we know they are of particular interest to consumers. The evaluation of possible health implications for PRiF findings is complex as tests for genotoxicity are commonly performed at higher doses (orders of magnitude higher) than the dietary exposure levels that are assessed in PRiF reports. As such it is difficult to conclude specifically, and to extrapolate the findings in the laboratory to the context of findings in the PRiF monitoring and the presence of residues at low levels in foods. Where relevant some reassurance that any risks are likely to be small can be gained if increased cancer incidence, which may be due to gene mutations, does not occur in long-term animal feeding studies, designed to detect such observations. Where relevant we will indicate this. Due to the uncertainty about the potential for genetic damage (genotoxicity) at low doses, PRiF will always conclude that on a precautionary basis any findings of genotoxic substances in food are undesirable.



## **Assessment of genotoxicity (Q4 2021) and conclusions:**

### **Residues found in this report that have genotoxic potential (concluded from laboratory studies on animals): omethoate, monocrotophos**

There is some evidence (*in vitro* and/or *in vivo*<sup>14</sup>) that these pesticides can damage genetic material (are genotoxic). There is some reassurance that risks of developing ill health effects following single or repeat exposures are likely to be low, since they did not increase cancer incidence in studies with repeat daily doses over their life-span in rats or mice. The doses used in both the genotoxicity tests and the cancer studies were orders of magnitude higher than the exposures estimated in this assessment. It is not known if lower doses which are not toxic also have this effect.

### **Residues found in this report where toxicological data are suggestive of genotoxicity but not certain: dimethoate, chlorpyrifos, triallate**

It is unclear whether these pesticides can damage genetic material (are genotoxic). There is some evidence from studies performed *in vitro* and/or *in vivo* that they may be genotoxic. Whilst there are negative results in the available *in vivo* studies, the currently recommended *in vivo* follow up studies, that may clarify the genotoxic potential of these pesticides, have not been performed. There is some reassurance that risks of developing ill health effects following single or repeat exposures are likely to be low, since they did not cause cancer in cancer or other long-term studies with repeat daily doses in animals over their life-span. The doses used in these studies were orders of magnitude higher than the exposures estimated in this assessment. It is not known if lower doses which are not toxic also have this effect.

**Conclusions:** Overall, we conclude that on a precautionary basis any residue findings of these pesticides are undesirable due to the uncertainty regarding genotoxicity at low doses; however, we consider any risks of adverse health effects are low at the highest levels of exposure after eating large portions (97.5th percentile consumption) of the foods containing the levels of these pesticides found in this report.

---

<sup>14</sup> *in vivo/in vitro*: see glossary pg 159

## Section 4: issues arising in this report and updates on previous reports

### Issues arising in this report

#### Chlorate

We have been testing a limited number of foods for chlorate since 2016. The pesticide sodium chlorate is a residual broad action weed killer that is not authorised for use in the EU or UK. However, we are confident that the residues we are detecting come from use of chlorine-based disinfectants used to maintain microbiological safety (control microorganisms that cause food poisoning). Because these residues are unavoidable, and important for the maintaining of microbiological control vital for food safety, we are not treating these results as breaches of the MRL. **We are not advising that food companies change their existing practices as a result of our findings, but they should be aware about the ongoing discussion in this area.**

We are only part of the work going on across government and beyond to consider what to do about chlorate residues in food and water.

#### MRLs after 20 June 2020

After detailed discussion and consultation with stakeholders, the EU agreed new MRLs for chlorate that came into force on 20 June 2020. All samples for foods, other than infant food, covered by this report were taken after 20 June 2020 when the new, higher MRLs were in place.

The new chlorate MRLs include a footnote referring specifically to taking account of the use of biocides during processing in addition to the MRLs for food as harvested or initially produced. The footnote exceptionally specifies that for considering compliance with chlorate MRLs, simple types of processing that do not affect the other residue levels, such as packing, washing, chopping and freezing can be taken into account.

The responsibility for providing evidence showing that residues from processing can be taken into account, lies with the food business operator, and so we will be interested to see such evidence where appropriate. HSE will decide whether the footnote can be applied and if so, this will be reflected in our reports.

The Food and Biocides Industry Group have produced more detailed information and guidance on this topic which is available on the Chilled Food Association's website at <https://www.chilledfood.org/fbig/>.

#### Infant food

Infant food MRLs are set under separate legislation managed by UK health departments. The footnote that applies to other foods cannot be used for infant foods, although residues occur for the same reasons. UK health departments are working with HSE and FSA to resolve this.

## **Best practice for use**

The Food Standards Agency is working with the food industry to develop and promote best practice in the use of sanitisers. This is important because the presence of low-level residues of chlorate in food results from measures taken by the food and water industries to protect food safety by reducing microbiological contamination of food and drink (including drinking water, which is a significant source of chlorate in food). Chlorate itself is not used as a disinfectant, but chlorine-based sanitisers can contain small amounts of chlorate.

## **Drinking Water**

In national legislation throughout the UK it is already a requirement to keep disinfection by-products as low as possible. This is usually achieved through management of disinfectant dosing and storage.

## **Advisory Committee on the Microbiological Safety of Food**

### **Microbiological safety of food**

The HSE is working with the Advisory Committee on the Microbiological Safety of Food to understand how changes to pesticide MRLs affect biocide use, microbiological food safety, and any change to the overall risk to consumers taking into account both chemical and microbiological safety.

### **Dietary intakes**

Since 2018 the Committee on Toxicity of Chemicals in Food, Consumer Products and the Environment (COT) has been considering chlorate as part of its on-going work looking at the chemicals in the diet of infants and young children (up to 5 years). The European Food Safety Authority's 2015 opinion on chlorate<sup>15</sup> establishes appropriate health-based guidance values for chlorate exposure to protect against acute and chronic risks to health.

## **DDT**

The use of DDT is banned or heavily restricted in many countries. It isn't allowed for use on food crops anymore, but it is still used in some countries outside the EU as a public health insecticide. Residues of DDT take a long time to break down in the environment and can accumulate in fatty tissue which is a major reason that it has been banned in the EU and many other countries.

Due to the bans and restrictions on use the levels in food have decreased substantially since the 1960s and 1970s. Even so, because it takes a long time to break down we do expect, and do see, occasional DDT residues in our monitoring results. Overall, the incidence and the size of residues have fallen steadily over time, which is what we would expect. In recent years none of our findings were unusual, unexpected or of concern. We can tell from the chemical form that we detect whether the residues we have found are from historic use (which is what we usually find). We explain this every time we publish DDT results to try to make it as clear as we can that the results show food producers are not using DDT today. However, there are occasional media stories about DDT and various links and associations, which do not make this distinction.

---

<sup>15</sup> [EFSA Journal 2015;13\(6\):4135 \[103 pp.\] Chlorate in food | EFSA \(europa.eu\)](https://www.efsa.europa.eu/en/efsajournal/doc/13636/attachment/68262.pdf)

The residues we find nowadays are at levels that would not be expected to have any effect on health, either in the short term or in the long term, when checked against today's understanding of the effect of DDT on health. As a committee, we take care to ensure we look thoroughly at this, and the Food Standards Agency is also actively involved in our considerations.

## Processing factors

As the surveillance programme monitors residues in all types of food, from raw commodities (e.g., potatoes) to processed (e.g., wine), dried (e.g., dried fruit) and composite foods (e.g., fruit bread), consumer risk assessments are specifically tailored to address processed and mixed food products. MRLs are generally set for raw commodities, although when MRLs are established the assessment of dietary intakes takes into account the potential for residues to remain in processed foods produced from the raw agricultural commodities. MRLs have been set for processed infant foods, and in future may be extended to other processed food products.

MRLs apply to all traded foods, including foods used as ingredients. The law specifies the level to apply to foods as they are traded. For almost all foods that means their raw, unprocessed form. But MRLs also apply to prepared and processed foods in which case the effect of processing needs to be taken into account.

In nearly all cases the MRL is set for the food in its raw, unprocessed form (the form of each food to which MRLs apply is listed in Annex I of Regulation 396/2005). These MRLs can be applied to processed foods using appropriate processing factors. Processing factors take account of the effect of processing on the food as traded. Different forms of processing may remove, concentrate, or dilute residues, and the effect may vary depending on the food and the pesticide concerned. Multiplying the processing factor by the original MRL gives a calculated MRL that can indicate the food was made with an ingredient or ingredients which had residues over the original MRL.

Calculating the MRLs for processed goods is dependent on the information available. HSE will contact the supplier if residues exceed the calculated MRL to give them an opportunity to provide relevant information to support the calculation.

Processing factors for olive oil. PRiF use the general principle that is used in the EU reports. The general principle that for virgin olive oil, and for fat soluble compounds a default x 5 factor can be applied (in the absence of a more specific experimentally derived PF), see the link below.

<https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32018R0555&from=EN>

Virgin olive oil (if no specific oil processing factor is available, a default factor of 5 may be applied for fat-soluble substances, taking into account an olive oil production standard yield of 20 % of the olive harvest; for non-fat- soluble substances a default oil processing factor of 1 may be used).

### Processing factors and MRLs used for bread

Food type	Pesticide	Processing factor	MRL for unprocessed grain (mg/kg)	Bread MRL (mg/kg)
Wholemeal wheat bread	Chlormequat	0.58	7	4.06
Wholemeal wheat bread	Fosetyl	0.98	2	1.96
Wholemeal wheat bread	Glyphosate	2.5	10	25
Wholemeal wheat bread	Mepiquat	0.94	3	2.82
Other wheat bread	Chlormequat	2	7	14
Other wheat bread	Cypermethrin	0.17	2	0.34
Other wheat bread	Fosetyl	0.82	2	1.64
Other wheat bread	Glyphosate	0.16	10	1.6
Other wheat bread	Mepiquat	0.16	3	0.48
Other wheat bread	Pirimiphos methyl	0.18	5	0.9

**Processing factors and MRLs used for olive oil.**

<b>Food type</b>	<b>Pesticide</b>	<b>Processing factor Default x5 for fat soluble</b>	<b>MRL for olives for oil production mg/kg</b>	<b>Olive oil MRL mg/kg</b>
Olives for oil production	Chlorpyrifos	5	0.01	0.05
Olive oil production	cypermethrin	5	0.05	0.25

## Follow-up from Previous Reports

### Quarter 4 2020

#### Cauliflower

Fonicamid (sum): Sample number 4634/2020

We passed details of a sample of cauliflower from the UK that contained fonicamid (sum) to HSE. Following investigation this case has been closed. HSE investigation concluded that insufficient tank cleaning measures had been followed, after a legal application of fonicamid to a crop of Brussels sprout, resulting in contamination and the detection of an unauthorised residue of fonicamid on cauliflower. The level of residue indicated there was no human health concern. HSE communicated with the supplier with advice on the correct use of pesticide products and measures to prevent contamination.

### Quarter 3 2021

#### GB Raspberry

Fonicamid (sum): Sample number 0355/2021

We passed details of a sample of raspberries from the UK that contained fonicamid (sum) to HSE. HSE enquiries are not yet complete, and an update will appear in a future report.

#### GB Spring greens and kale

Tebuconazole: Sample number 0363/2021

We passed details of a sample of spring greens from the UK that contained tebuconazole to HSE. HSE investigation concluded that insufficient tank cleaning measures had been followed, after a legal application of tebuconazole to a crop of cabbage resulting in contamination. The level of residue indicated there was no human health concern. HSE communicated with the supplier with advice on the correct use of pesticide products and measures to prevent contamination.

#### GB Spring greens and kale

Cyantraniliprole and Tebuconazole: Sample number 1419/2021

We passed details of a sample of spring greens from the UK that contained cyantraniliprole and tebuconazole to HSE. HSE investigation concluded that insufficient tank cleaning measures had been followed, after a legal application of cyantraniliprole and tebuconazole to a crop of cabbage resulting in contamination. The level of residue indicated there was no human health concern. HSE communicated with the supplier with advice on the correct use of pesticide products and measures to prevent contamination

## Brand name details of samples where follow-up action is now complete

Sample ID	Date of Sampling	Description	Country of Origin	Retail Outlet	Address	Brand Name	Packer / Manufacturer	Pesticide residues found in mg/kg (MRL)
4634/2020	06/10/2020	Cauliflower	UK	Jack Waters and Son Ltd	South Yorkshire Fresh Produce and Flower Centre, Parkway Drive, Sheffield S9 4WN			flonicamid (sum) 0.1 (MRL = 0.03*)
0363/2022	09/06/2021	Spring greens and kale	GB	Asda	Stanley Grove, Longsight, Manchester M12 4NH	Asda Grower's Selection	Asda Stores Ltd Asda House, South Bank, Great Wilson Street, Leeds LS11 5AD	azoxystrobin 0.02 (MRL = 6)
								difenoconazole 0.02 (MRL = 2)
								indoxacarb 0.2 (MRL = 0.4)
								Spirotetramat (partial sum) 0.3 (MRL = 7)
								tebuconazole 0.08 (MRL = 0.02*)
1419/2021	23/08/2021	Spring greens and kale	GB	Asda	8 Whalebone Lane, Dagenham RM8 1BB	Asda Grower's Selection	Asda Stores Ltd Asda House, South Bank, Great Wilson Street, Leeds LS11 5AD	azoxystrobin 0.1 (MRL = 6)
								cyantraniliprole 0.02 (MRL = 0.01*)
								difenoconazole 0.09 (MRL = 2)
								indoxacarb 0.04 (MRL = 0.4)



								spinosad (sum) 0.08 (MRL = 2)
								tebuconazole 0.03 (MRL = 0.02*)

## In our next report:

In Quarter 1 of 2022 we will look at results for:

### **Samples collected in GB**

- Apples (eating)
- Avocado
- Beans with pods
- Cabbage
- Cucumber
- Grapes
- Lettuce
- Milk
- Peaches and nectarines
- Pork
- Potatoes
- Spinach
- Strawberry
- Tomatoes

### **Samples collected in NI**

- Apples (eating)
- Beans with pods
- Cabbage
- Fish (sea)
- Game
- Grapes
- Lettuce
- Peaches and nectarines
- Potatoes
- Spinach
- Strawberry
- Tomatoes

## Section 5: background and reference

### Reasons for pesticide residue testing

Food safety is important. Modern food production processes have given us plentiful supplies of a wide range of good quality affordable produce.

In the food industry of today the production environment can be managed from the preparation of seeds used for crops, through to growth, harvesting and storage of the produce.

One of the ways the food industry controls the environment in which foodstuffs are produced is by applying pesticides. They help farmers and growers maximise the production of food stuffs by, for example, preventing weeds inhibiting the growth of the crop, or insects destroying or infesting them. Pesticides can also be used to help protect seeds or prolong the life of crops after they have been harvested. Biological and physical (cultural) controls are also used to protect crops or as part of an integrated system.

As pesticides are used to control unwanted pests, weeds, and diseases, they can potentially also harm people, wildlife and the environment. This is why the UK, in common with most other countries, imposes legally enforceable conditions as to how and when pesticides can be used. No pesticide can be supplied or used on a food or ornamental crops in the UK without Government authorisation. To obtain this authorisation the manufacturer of the pesticide must show that it does not present a concern for people's health or the environment. Naturally derived and synthetic pesticides are subject to the same regulation.

Once the authorisation has been granted Government authorities carry out follow up checks to ensure that the authorisation is providing the necessary degree of protection to users, consumers and the environment and that those who use pesticides are complying with conditions specified within it.

The Government authority responsible for checking pesticide residues in foodstuffs is the Health and Safety Executive. Defra's Expert Committee on Pesticide Residues in Food (PRiF) oversees and provides an independent check on this work. We know that the use of pesticides on crops may lead to traces (residues) of these chemicals in food and we expect to find these in our monitoring programme.

### Detail of reporting practice

#### Results by food commodity

- We include information about the survey (for instance where samples came from) for each commodity
- Detailed tabulated results are at the back of this report - these tables are also available for download from our website
- We summarise our findings and any follow-up action taken.

#### Risk assessments – single residues

- All results are screened by HSE to check for intakes above the toxicological reference values, the Acute Reference Dose (ARfD) or the Acceptable Daily Intake (ADI). HSE assumes a relatively high level of intake and also assumes that most produce is eaten whole including peel/skin even when these are rarely consumed

- Where intakes above the toxicological reference values are identified, we consider a detailed risk assessment prepared by HSE (at section II of this report).
- Our observations and the follow-up action taken are summarised in the section for that food.

### **Risk assessments – multiple combined residues**

- Residues of more than one pesticide from the same category/class of particular categories of pesticides, which have a similar toxicological mode of action, are initially screened by HSE to check for intakes that might need further combined assessment.
- Where combined intakes exceed the initial screen 'trigger', we consider a detailed combined risk assessment prepared by HSE (at section 3 of this report). Further details on the approach are explained in Section 3.
- Our observations and any follow-up action taken are summarised in the section for that food commodity.

### **Risk assessment - conclusions**

- Where, in the light of current knowledge and considering the usual level of scientific uncertainty the intake will not cause ill health the conclusion will say no effect on health is expected.
- Where, in the light of current knowledge the intake is not likely to cause ill health, the conclusion will be less definite and state that an effect on health is unlikely.
- Where scientific uncertainty is greater or if risk of adverse health effects could be higher more information is provided.

### **Residues in GB and NI produce of pesticides which do not have a PPP authorised for use on that crop in GB and NI.**

- All residues found in foods produced in GB or NI are checked by HSE to make sure there is a PPP containing that pesticide authorised for use on that crop.
- Where there is no GB or NI authorisation is identified, details of the sample are referred to the Enforcement Section for follow up.
- Our observations and any follow-up action taken to date are summarised in the section for that food commodity. We may have to withhold details of samples while investigations are underway, in which case the details will be published in a later report.

### **Residues above the MRL, after taking into account measurement uncertainty**

- Samples containing residues above the MRL are listed at Appendix B, and those which are clearly above the MRL after taking into account measurement uncertainty of plus or minus 50% are highlighted.
- Our observations and any follow-up action taken are summarised in the section for that food commodity.
- The results in our reports are rounded for publication but not adjusted for measurement uncertainty.
- We apply measurement uncertainty only to decide whether to highlight a result as over the MRL in the brand name annex. To do this we use the actual value reported by the laboratory before rounding. If after taking measurement uncertainty into account that value is found to be over the MRL the result will be highlighted in the brand name annex.

For example:

- The lab reports the results of duplicate analysis of a residue above an MRL at 0.023 mg/kg and 0.025 mg/kg giving an average value of 0.024mg/kg. For reporting purpose this value would be 0.02 mg/kg.

- If measurement uncertainty is then applied to the reported value of 0.02 mg/kg it could take the value to between 0.01 - 0.03 mg/kg. If the MRL is 0.01 mg/kg the lower value would be at the MRL and there is no exceedance.
- However, if measurement uncertainty is applied to the measured result, e.g., 0.024 mg/kg the value could then be in the range of 0.012 – 0.036 mg/kg. In this case the lower value is above the MRL and so will be treated as an exceedance.

### Residues in organic food

- We monitor pesticide residues in all the GB and NI food supply, including organic food.
- We are not responsible for checking compliance with the rules associated with organic production. However, when we do detect residues in an organic food we explain whether or not those residues indicate a breach of the rules and inform Defra's Organic Farming Branch.

### Brand Name Annex

- Full brand name details for samples included in this report are published in a brand name annex. Within this annex, samples with results of interest are highlighted.
- Brand name details are only published when enough follow-up work is completed for us to be reasonably sure whether a breach of the law or good practice has occurred.
- Therefore, sometimes brand name details are withheld pending completion of this work but are published in a later report.

## Pesticides analysed as multi-component analytes and their reporting limits

Why some results cover more than one substance

Both the legal controls and our analytical tests are aimed at checking food for the presence of residues of specific pesticides. Residues are the chemical traces left behind after pesticides are used. In most cases the residue of a pesticide is measured by first identifying the pesticide and then measuring the quantity of that pesticide in the food itself. But for some pesticides the residue remaining in the food is known to be chemically different from the original pesticide and so the laboratory needs to look for more than one component. There are various reasons why this happens, for example:

- the animal or plant can change the pesticide into related chemicals
- the pesticide can change in the environment into related chemicals
- some pesticides are mixtures of chemicals, so the relevant components of the mixture need to be checked for
- in the laboratory sample preparation and/or analysis may change pesticides into related chemicals
- related chemicals may be pesticides in their own right

The MRL setting process takes account of all these issues. The EU may set a complex residue definition to ensure that the identity and quantity of the residue found is representative of the pesticide present. A complex residue definition may be set where it is necessary for safety reasons or to be able to accurately identify the pesticide residue present in the food. This definition usually includes the actual pesticide, plus other related chemicals. These residues are usually reported together as a "sum". Sometimes different foods need different definitions because different pesticide residues are known to occur in that food. For instance, plants and animals may metabolise a pesticide differently, which forms different residues.

The full definitions of pesticides that we have found in our surveys are described in the table below. If you would like more detail about a particular residue definition, please get in touch. You can email us at [prif@hse.gov.uk](mailto:prif@hse.gov.uk) and other contact details are on the back cover.

Where the detailed individual analysis results tell us something useful, we mention that in our conclusions.

### How we calculate sums

Unless the definition says otherwise, the summed result is a simple addition. For individual components that are not detected that result is treated as a zero.

Where a residue definition says, “expressed as”, that means that the individual component results are adjusted by molecular weight before being added together. The residue definition is set this way so that the final calculated result for the whole definition is an expression of the level of the most toxic component, and so that value can be used directly in consumer risk assessment without further adjustment.

## Complex residue definitions used in our reports

There are a large number of pesticides used and types of food in the world. So other complex residue definitions may apply to food/pesticide combinations not yet considered by PRiF. You can look up all the EU MRL definitions for pesticide residues at the European Commission’s pesticide database at [EU-Pesticide Database](#)

Short name we use in our reports	Legal residue definition – These definitions apply to all foods unless otherwise stated
2,4-D (sum)	2,4-D (sum of 2,4-D and its esters expressed as 2,4-D)
abamectin (sum)	Abamectin (sum of Avermectin B1a, AvermectinB1b and delta-8,9 isomer of Avermectin B1a)
aldicarb (sum)	Aldicarb (sum of Aldicarb, its sulfoxide and its sulfone, expressed as Aldicarb)
aldrin and dieldrin	Aldrin and Dieldrin (Aldrin and dieldrin combined expressed as dieldrin), aka dieldrin (sum)
Amitraz	Amitraz (amitraz including the metabolites containing the 2,4 - dimethylaniline moiety expressed as amitraz)
BAC (sum)	Benzalkonium chloride (mixture of alkylbenzyltrimethylammonium chlorides with alkyl chain lengths of C <sub>8</sub> , C <sub>10</sub> , C <sub>12</sub> , C <sub>14</sub> , C <sub>16</sub> and C <sub>18</sub> )
benthiavalicarb (sum)	Benthiavalicarb (Benthiavalicarb-isopropyl (KIF-230 R-L) and its enantiomer (KIF-230 S-D) and diastereomers (KIF-230 R-L and KIF-230 S-D))
bixan (animal products)	Sum of bixafen and desmethyl bixafen expressed as bixafen This definition applies to animal products only
captan and folpet	Sum of captan and folpet aka captan/folpet This definition applies only to pome fruit (fruits such as apples and pears), strawberries, raspberries, currants, tomatoes, and beans. For all other foods there are separate MRLs for captan only and for folpet only.

carbendazim (animal products)	Carbendazim and thiophanate-methyl, expressed as carbendazim
Carbendazim (sum)	Carbendazim and benomyl (sum of benomyl and carbendazim expressed as carbendazim)
carbofuran (sum)	Carbofuran (sum of carbofuran and 3-hydroxy-carbofuran expressed as carbofuran)
chlordan (animal products)	Chlordane (sum of cis- and trans-isomers and oxychlordan expressed as chlordan) This definition applies to animal products only
chlordan (sum)	Chlordane (sum of cis- and trans- isomers) This definition applies to all foods except animal products
chlorpropham (potatoes)	Chlorpropham only This definition applies only to potatoes
chlorpropham (sum for animal products)	Chlorpropham and 4-hydroxychlorpropham-O-sulphonic acid (4-HSA), expressed as chlorpropham This definition applies only to animal products
chlorpropham (sum)	Chlorpropham (Chlorpropham and 3-chloroaniline, expressed as Chlorpropham) This definition applies to all foods except potatoes and animal products
DDAC (sum)	Didecyldimethylammonium chloride (mixture of alkyl-quaternary ammonium salts with alkyl chain lengths of C <sub>8</sub> , C <sub>10</sub> and C <sub>12</sub> )
DDT (sum)	DDT (sum of p,p'-DDT, o,p'-DDT, p-p'-DDE and p,p'-TDE (DDD) expressed as DDT)
Dichlorprop	Sum of Dichlorprop, including dichlorprop-p and its conjugates, expressed as dichlorprop
dicofol (sum)	Dicofol (sum of p, p' and o,p' isomers)
Dimethenamid	Dimethenamid-p (Dimethenamid-p including other mixtures of constituent isomers (sum of isomers))
dimethoate (sum)	Dimethoate (sum of dimethoate and omethoate expressed as dimethoate)
disulfoton (sum)	Disulfoton (sum of disulfoton, disulfoton sulfoxide and disulfoton sulfone expressed as disulfoton)

dithiocarbamates	Dithiocarbamates are a group of pesticides that are chemically similar. Testing for them individually in routine analysis is not possible, so MRLs are set for a test for the group.
endosulfan (sum)	Endosulfan (sum of alpha- and beta-isomers and endosulfan-sulphate expressed as endosulfan)
fenamiphos (sum)	Fenamiphos (sum of fenamiphos and its sulphoxide and sulphone expressed as fenamiphos)
fenchlorphos (sum)	Fenchlorphos (sum of fenchlorphos and fenchlorphos oxon expressed as fenchlorphos)
fensulfothion (sum)	Fensulfothion (sum of fensulfothion, its oxygen analogue and their sulfones, expressed as fensulfothion).
fenthion (sum)	Fenthion (fenthion and its oxygen analogue, their sulfoxides and sulfone expressed as parent)
fenvalerate & esfenvalerate (all isomers)	Fenvalerate (any ratio of constituent isomers (RR, SS, RS & SR) including esfenvalerate)
fipronil (infant food)	Sum of fipronil and fipronil-desulfinyl, expressed as fipronil This definition applies to foods for babies only
fipronil (sum)	Fipronil (sum Fipronil and sulfone metabolite (MB46136) expressed as Fipronil) This definition applies to all foods except foods for babies
flonicamid (sum)	Flonicamid (sum of flonicamid, TNFG and TNFA) This definition applies to all food except animal products. The full definition must be sought. Residues found are usually of the metabolites.
fluazifop-p-butyl (sum)	Fluazifop-P-butyl (fluazifop acid (free and conjugate))
Fosetyl (sum)	Fosetyl-Al (sum of fosetyl, phosphonic acid and their salts, expressed as fosetyl)
haloxyfop (sum)	Haloxyfop including haloxyfop-R (Haloxyfop-R methyl ester, haloxyfop-R and conjugates of haloxyfop-R expressed as haloxyfop-R)
Heptachlor (infant food)	Sum of heptachlor and trans heptachlor epoxide This definition applies to foods for babies only
Heptachlor (sum)	Heptachlor (sum of heptachlor and heptachlor epoxide expressed as heptachlor) This definition applies to all foods except infant foods



hexachlorocyclohexane (sum)	Hexachlorocyclohexane (HCH), sum of isomers, except the gamma isomer This definition applies to all foods except animal products (For animal products the alpha and beta isomers have separate MRLs)
Malathion	Malathion (sum of malathion and malaoxon expressed as malathion)
MCPA (animal products)	[Residue definition, animal products] MCPA, MCPB and MCPA thioethyl expressed as MCPA This definition applies to animal products only
MCPA (sum)	MCPA and MCPB (MCPA, MCPB including their salts, esters and conjugates expressed as MCPA) This definition applies to all foods except animal products
Mepanipyrim (sum)	Mepanipyrim and its metabolite (2-anilino-4-(2-hydroxypropyl)-6-methylpyrimidine) expressed as mepanipyrim
methiocarb (sum)	Methiocarb (sum of methiocarb and methiocarb sulfoxide and sulfone, expressed as methiocarb)
methomyl (sum)	Sum of methomyl and thiodicarb expressed as methomyl
Oxydemeton-methyl (sum)	Oxydemeton-methyl (sum of oxydemeton-methyl and demeton-S-methylsulfone expressed as oxydemeton-methyl)
parathion-methyl (sum)	Parathion-methyl (sum of Parathion-methyl and paraoxon-methyl expressed as Parathion-methyl)
Permethrin	Permethrin (sum of isomers)
phorate (sum)	Phorate (sum of phorate, its oxygen analogue and their sulfones expressed as phorate)
phosmet (sum)	Phosmet (phosmet and phosmet oxon expressed as phosmet) This definition applies to all foods except animal products
pirimicarb (sum)	Pirimicarb (sum of Pirimicarb and Desmethyl pirimicarb expressed as Pirimicarb) for certain animal products. Pirimicarb only for fruit and vegetables and some animal products.
Prothioconazole (sum)	Prothioconazole (sum of prothioconazole-desthio and its glucuronide conjugate, expressed as prothioconazoledesthio) This definition applies to animal products only
PTU & propineb	Sum of PTU and propineb This definition applies to food for babies only

quintozene (sum)	Quintozene (sum of quintozene and pentachloro-aniline expressed as quintozene)
Prochloraz (sum)	Prochloraz (sum of prochloraz and its metabolites containing the 2,4,6-Trichlorophenol moiety expressed as prochloraz)
Terbufos (sum)	Terbufos (sum of terbufos, its sulfoxide and sulfone This definition applies only to foods for babies)
thiamethoxam (sum)	Thiamethoxam (sum of thiamethoxam and clothianidin expressed as thiamethoxam) There are <u>also</u> separate clothianidin MRLs
tolyfluanid (sum)	Tolyfluanid (Sum of tolyfluanid and dimethylaminosulfotoluidide expressed as tolyfluanid)
triadimefon & triadimenol	Triadimefon and triademenol
vinclozolin (animal products)	Vinclozolin, iprodione, procymidone, sum of compounds and all metabolites containing the 3,5-dichloroaniline moiety expressed as 3,5-dichloroaniline This definition applies to animal products only
vinclozolin (sum)	Vinclozolin (sum of vinclozolin and all metabolites containing the 3,5-dichloroaniline moiety, expressed as vinclozolin) This definition applies to all foods except animal products

## Glossary

This is a 'standard' glossary which defines the key terms used in the PRiF reports. Not all the terms listed here are used in this particular report.

**97.5<sup>th</sup> percentile consumer:** Please refer to glossary entry for 'High level consumer'.

**Acceptable Daily Intake (ADI):** This is the amount of a chemical which can be consumed every day for a lifetime in the practical certainty, on the basis of all known facts, that no harm will result. It is expressed in milligrams of the chemical per kilogram of body weight of the consumer. The starting point for the derivation of the ADI is usually the 'no observed adverse effect level' (NOAEL) that has been observed in animal studies for toxicity. This is then divided by an uncertainty factor (most often 100) to allow for the possibility that animals may be less sensitive than humans and also to account for possible variation in sensitivity between individuals. The studies from which NOAELs and hence ADIs are derived take into account any impurities in the pesticide active substance as manufactured, and also any toxic breakdown products of the pesticide.

**Acetylcholine:** Acetylcholine is a neurotransmitter, a chemical that carries signals through the nervous system. See *cholinergic*

**Acetylcholinesterase:** This is an enzyme which degrades acetylcholine and is involved in the regulation of nerve impulses. Inhibition of this enzyme can interfere with this nerve transmission function. This is a short-term effect of concern with organophosphate and carbamate pesticides at levels above the ARfD.

**Acute Reference Dose (ARfD):** The definition of the ARfD is similar to that of the ADI, but it relates to the amount of a chemical that can be taken in at one meal or on one day without appreciable health risk to the consumer. It is normally derived by applying an appropriate uncertainty factor to the lowest NOAEL in studies that assess acute toxicity or developmental toxicity.

As a matter of policy, the EU does not use NOAELs from tests that involve deliberate administration of pesticides to humans to determine ADIs and ARfDs. However, where such data have been ethically and scientifically derived some authorities, e.g., the World Health Organization, do consider such data. Where human data are used there is usually less uncertainty in the resulting reference value compared to extrapolating from animal tests to humans, and a lower uncertainty factor (most often 10) is used to account for the variation in sensitivity between individuals.

The initial risk assessments in PRiF reports use the agreed EU reference values. However, where intakes are above the EU value and a reference value based on acceptable human data is available a refined assessment, which is a more appropriate indicator of the risk, is also reported.

**Analyte:** This is the name for the substance that the PRiF surveys look for and measure if present; it could be a pesticide itself or a product from a pesticide when it is degraded, or metabolised.

**COLEACP (Europe-Africa-Caribbean-Pacific Liaison Committee):** It aims to promote the competitive export of fresh fruit, vegetables, flowers and ornamental plants from the ACP. Its specialised information and advisory services are open to all ACP companies in the horticultural

export sector and are financed by the European Commission. It has two overriding objectives to enable ACP companies to comply with European food safety and traceability requirements and to consolidate the position of small-scale producers in the ACP horticultural export sector.

**Cholinergic:** In relation to the animal nervous system, processes and structures are cholinergic if they release or use acetylcholine.

**Cryogenic Milling:** Processing of commodities at very low temperatures can be achieved by milling/grinding pre-frozen samples in the presence of dry ice, a procedure known as 'cryogenic milling'.

**Extensions of Authorisations for Minor Use (EAMUs):** Users and authorisation holders of agricultural Plant Protection Products (PPP) may apply to have the authorisation of specific PPP's extended to cover uses additional to those authorised and shown on the manufacturer's product label. For many reasons, label recommendations of authorised pesticides do not cover the control of every problem which may arise. This is particularly true for crops that are grown on a comparatively small scale in the UK as well as for pests and diseases that occur less often, or which are new to the UK. As part of the process evidence on residues that would arise from the use is required, and consumer safety is evaluated and if necessary a specific MRL set. EAMU is pronounced "emu" these types of authorisations are also informally called "off labels".

**Genotoxicity:** Genotoxicity is the effect of substances (called genotoxins) which can alter or damage the genetic material (DNA, RNA or chromosomes) within a cell. Cells have the capacity to protect themselves from genotoxic effects by many repair processes and therefore many genotoxic events do not become evident as mutations. Where mutations occur, this can lead to cancer or effects that can be passed to unborn children (e.g., birth defects, inherited diseases).

**Good Agricultural Practice in the Use of Pesticides (GAP):** The nationally authorised safe uses of pesticides under conditions necessary for effective and reliable pest control (the way products should be used according to the statutory conditions of authorisation which are stated on the label). GAP encompasses a range of pesticide applications up to the highest authorised rates of use, applied in a manner which leaves a residue which is the smallest practicable. Authorised safe uses are determined at the national level and include nationally registered recommended uses, which take into account public and occupational health and environmental safety considerations. Actual conditions include any stage in the production, storage, transport, distribution and processing of food commodities and animal feed.

**High-level Consumer:** A term used in UK risk assessment calculations to describe the amount of food consumed by a person. In line with internationally agreed approaches, the PRiF uses the 97.5<sup>th</sup> percentile value, which is generally about three times the average amount consumed. This takes account of different eating patterns that may occur throughout the population.

**Human Data:** See under Acute Reference Dose

**In vitro:** a test performed *in vitro* "in the glass" means that it is performed outside of a living organism and usually involves isolated tissues, organs, or cells.

**In vivo:** live animal studies

**Import Tolerance:** an MRL set for imported products where the use of the active substance in a plant protection product on a commodity is not authorised in the European Community (EC) or an existing EC MRL is not sufficient to meet the needs of international trade. All import tolerances are assessed for consumer safety.

**Imported:** The tables in the reports record whether the sample was of UK origin, or imported. This can mean different things depending on the commodity. See also 'Origin'. The PRiF report the country from where the produce has been imported only if this is clear from the packaging or labelling.

**JMPR:** Joint FAO/WHO Meeting on Pesticide Residues, which conducts scientific evaluations of pesticide residues in food.

**LOD (Limit of Determination) and LOD MRLs:** The Limit of Determination (LOD) is the lowest concentration of a pesticide residue or contaminant that can be routinely identified and quantitatively measured in a specified food, agricultural commodity or animal feed with an acceptable degree of certainty by the method of analysis.

**LOD MRL (Maximum Residue Levels set at the LOD):** These are marked by a '\*'. For some pesticides and commodities insufficient trials data are available on which to set a maximum residue level or there may be no use of the pesticide on that crop. In these cases, the MRL may be set at a default level i.e., at the limit of determination (LOD) where analytical methods can reasonably detect the presence of the pesticide. **These MRLs are not based on Good Agricultural Practice (GAP).** Also, see under Reporting limit.

**Lowest Observed Adverse Effect Level (LOAEL):** The lowest concentration or amount of a substance, found by experiment or observation, which causes detectable adverse alteration of morphology, functional capacity, growth, development or life span of the target organism under defined conditions of exposure.

**Off Label:** See Extensions of Authorisations for Minor Use (EAMUs)

**Maximum Residue Level (MRL):** The maximum concentration of a pesticide residue (expressed as mg/kg) legally permitted in or on food commodities and animal feeds. MRLs are based on good agricultural practice data and residues in foods derived from commodities that comply with the respective MRLs are intended to be toxicologically acceptable.

MRLs are intended primarily as a check that GAP is being followed and to assist international trade in produce treated with pesticides. **MRLs are not in themselves 'safety limits'**, and exposure to residues in excess of an MRL does not automatically imply a hazard to health.

The MRLs applicable in the UK are now largely set under EC legislation.

Maximum Residue Levels (MRLs) reflect levels of pesticides that could occur in produce, which has been treated in accordance with good agricultural practice. Where pesticides do not give rise to readily detectable residues, or are not authorised for use on particular commodities, MRLs are set at the lowest level which can be identified in routine laboratory analysis. Thus, they provide a mechanism for statutory controls on pesticides in produce which is put into circulation and for monitoring correct use of these chemicals.

If no use of a pesticide on a crop is identified when MRLs are set the tolerance for that pesticide/crop combination is set at the limit of determination (effectively zero). Limit of determination MRL are marked by a '\*'.

MRLs are established under the Pesticides (Maximum Residue Levels in Crops, Food and Feeding Stuffs) (England and Wales) Regulations 1999 (as amended), the Pesticides (Maximum Residue Levels in Crops, Food and Feeding Stuffs) (Scotland) Regulations 2000 and the Pesticides (Maximum Residue Levels in Crops, Food and Feeding Stuffs) Regulations (Northern Ireland) 2002. These Regulations list all statutory MRLs established under UK national or EC

procedures. Today, virtually all these MRLs are set under an ongoing EC programme and the Regulations are amended periodically as levels are set for increasing numbers of pesticides.

There are a number of pesticides which do not yet have statutory MRLs. In the absence of such MRLs we advise suppliers to adhere to any appropriate levels established by the Codex Alimentarius Commission (CAC) a United Nations body established to promote global trading standards. Codex MRLs are not statutory but have been risk-assessed when set and provide a suitable standard in the absence of a statutory MRL.

MRLs may be extended to composite and processed products but levels are not specifically laid down in legislation. They are derived by calculation on an individual basis.

**Maximum Residue Levels set at the LOD (LOD MRL):** See LOD MRL. For some pesticides and commodities, insufficient trials data are available on which to set a maximum residue level or there may be no use of the pesticide on that crop. In these cases, the MRL may be set at a default level, i.e., at the limit of determination (LOD) where analytical methods can reasonably detect the presence of the pesticide. **These MRLs are not based on Good Agricultural Practice (GAP).**

**MRL exceedances:** When a residue is found at a level higher than that set for the MRL.

**MRL Exceedances and Relationship with the Acceptable Daily Intake (ADI):** Before permitting any use of a pesticide, a detailed assessment is made to ensure that residues in foods derived from commodities comply with MRLs and will not give rise to unacceptable risks to consumers. MRLs do take account of consumer safety aspects and, in effect, are set at levels below safety limits. However, MRLs must not be confused with safety limits, which are expressed in terms of the acceptable daily intake (ADI) of a particular pesticide residue from all sources. The ADI (expressed as mg/kg bw/day) is the amount of chemical that can be consumed every day of an individual's entire lifetime in the practical certainty, on the basis of all known facts, that no harm will result. See ADI for further information.

Whenever unexpectedly high or unusual residues occur during monitoring, the risk to consumers, from exposure to residues at the highest levels found, is assessed by comparison of predicted intakes with the ADI or ARfD as appropriate.

**No MRL:** For certain pesticides an MRL may not have been set.

**Metabolite:** A degradation or conversion product from a pesticide when it is metabolised.

**Multiple Residues:** In this report this term is used to describe when more than one pesticide is found in an individual food sample. It may have arisen because the crop was treated at different times with pesticides applied singularly, or when pesticides are applied as mixtures (several pesticides mixed in the spray tank at the same time), or the marketed pesticide product contains more than one pesticide or any combination of these three situations. Mixtures may be used in response to specific pest pressures and also as part of strategies to minimise pesticide resistance building up on pest populations.

**NEDI:** National Estimate of Daily Intake. An estimate of intake of pesticide in the diet over the long-term to compare to the ADI. The NEDI is based on median or mean residue levels and a high level consumption (97.5<sup>th</sup> percentile value) for the daily amounts of the food item consumed over the long-term. For further details on the calculation of NEDIs please refer to section 3 of the data requirements handbook using the following link: [The HSE Pesticide Website](#) then search for Consumer Exposure. Here you will find information and further links.

**NESTI:** National Estimate of Short-Term Intake. An estimate of peak intake of pesticide in the diet to compare to the ARfD. The NESTI is based on the highest residue found multiplied by a variability factor (see glossary description) and a high level consumption (97.5<sup>th</sup> percentile value) for the amount of the food item consumed over a single day. For further details on the calculation of NESTIs please refer to section 3 of the data requirements handbook using the following link: [The HSE Pesticide Website](#) then search for Consumer Exposure. Here you will find information and further links.

**Neurotoxicity:** Neurotoxicity is the effect of substances (called neurotoxins) which alter the normal working of an animal's nervous systems and/or damage the nervous tissue.

**No Observed Adverse Effect Level (NOAEL):** The greatest concentration or amount of a substance, found by experiment or observation, which causes no detectable adverse alteration of morphology, functional capacity, growth, development or life span of the target organism under defined conditions of exposure.

**Off Label:** See EAMUs

**Origin:** The brand name annex reports the origins of the samples tested. This can mean different things depending on the commodity. For example, butter is often labelled as 'UK origin'; however, the majority of it comes in bulk from New Zealand and is split into smaller blocks and packaged in the UK. Lettuce is a fresh produce and 'UK origin' usually means that it has been grown and packaged in the UK. Processed commodities such as cereal bars often contain multiple raw ingredients, each of which may come from a different source/origin. Therefore, the origin of the produce usually reflects the place where it was manufactured. The PRiF report the origin as stated on the packaging or labelling of the commodity concerned, unless other more accurate information is available to indicate that the origin is from elsewhere. Some products are listed as 'unknown origin' because the labelling does not give this information.

**Parent:** The chemical form of a pesticide as applied to plants, as opposed to metabolites and breakdown products.

**Percentile:** A percentile is a value that divides a sample of measurements at a specific point when they are listed in ascending order of magnitude. For example, the 97.5<sup>th</sup> percentile from a food consumption survey is a value that is equal to or more than 97.5% of the measurements and equal to or less than 2.5% of the measurements. So, in a sample of 40 daily food consumption values, the 97.5<sup>th</sup> percentile is equal to or more than 39 of the measurements. Such high percentile estimates of food consumption are used in risk assessments as they are more protective than using average consumption levels.

**Permitted Level (PL):** The permitted levels (expressed as mg/kg), in specific commodities, of some substances which can be classified as pesticides but are controlled under the Miscellaneous Food Additives Regulations 1995 (S.I. 1995 No. 3187).

**Pesticide:** A pesticide is any substance, preparation or organism prepared or used for destroying any pest. The majority of pesticides sought by the PRiF in its monitoring are those used to control pests in agricultural crops, although non-agricultural products may be included where there is a specific reason for doing so, e.g., where there are implications in terms of possible intakes of residues.

**Probabilistic Modelling:** The usual estimates of consumer exposure use single high values for both consumption amounts and residue levels. Whilst these are based on realistic UK dietary survey data and residue levels, they tend to overestimate most representative intakes. This is

because they do not take into account actual variations in both amounts consumed and residue levels. Probabilistic modelling is a technique that considers all the possible different combinations of consumption and residue levels. This provides information on the probability of particular intakes occurring.

**Rapid Alert System for Food and Feed (RASFF):** The European Commission's Rapid Alert System for Food and Feed (RASFF) allows member authorities (EU and EFTA member States) to quickly exchange information about measures taken when responding to risks detected in food or feed. This exchange of information helps authorities in countries inside the European single market to act more rapidly and in a coordinated way in response to a possible health threats caused by food or feed.

RASFFs notifications about pesticide residues are sent when a residue is over the MRL taking into account measurement uncertainty and a potential consumer risk has been identified. For pesticide residues in food traded in the single market this means when a risk assessment has identified that risk to people eating the food cannot be ruled out.

More information is available on the European Commission website at [RASFF - Food and Feed Safety Alerts](#).

**Relationship between GAP and MRLs:** The MRL can be defined as the maximum concentration of a pesticide residue (expressed as mg/kg) likely to **occur** in or on food commodities and animal feeds, after the use of the pesticide according to the GAP.

**Reporting Limit:** The reporting limit is the lowest level at which residues will be reported by a laboratory for a survey, as agreed in advance with the laboratory. It can be equal to or higher than the limit of quantification (sometimes also referred to as the limit of determination). The limit of quantification is the lowest concentration that has been validated to meet strict acceptance criteria and may vary slightly from laboratory to laboratory depending on the equipment available and operating procedures used. The reporting limit should be at or below the MRL. For a small number of pesticides e.g., monocrotophos, we are looking for the pesticide below the LOD MRL because we are specifically interested in prevalence in food due to the nature of the pesticide. In such cases, tests are performed in the laboratory to support the lower reporting limits by validating the method at lower limits. **'None were detected above the Set RL'**: This term is used in the Brand Name Annex, where no residues were found above their reporting limit.

**Residue:** Residues may be present in vegetable and animal products following the application(s) of a pesticide(s). They may not only include the pesticide that was applied but other degradation or reaction products and metabolites that may be of toxicological significance. The levels or amounts of residues present are expressed in milligrams of the chemical in a kilogram of crop/food/commodity (mg/kg), or parts per million.

**Risk Assessment:** A risk assessment is carried out when residues are found in foods to determine whether, at the levels found, they present a concern for consumer health or not. Consumer risk assessments are routinely conducted as part of the approval process for pesticides and are based on residue trials. Approval of a pesticide is only recommended when the consumer risk is acceptable.

**Safety Factor:** Values used in extrapolation from experimental studies in animals (usually 100) or humans (usually 10) to the population: for PRiF assessments this represents a value by which the NOAEL is divided to derive an ADI or ARfD. The value depends on the nature of the effect, the dose-response relationship, and the quality of the toxicological information available. The use of such a factor accounts for possible differences in susceptibility between the animal



species tested and humans, and for variation between different individuals in the population. The terms 'uncertainty factor' and 'assessment factor' are also sometimes used for this factor; the PRiF will use 'safety factor'.

**Sample:** The nature of all samples is as designated in the EC's 'sampling' Directive – 2002/63/EC. Examples are: apple – at least 10 apples weighing at least 1 kg; grapes – at least 5 bunches, weighing at least 2 kg.

**Technical Exceedances:** When an MRL has been set at the LOD because there have been no data to support a higher level. In the context of this report, 'technical exceedances' always relate to produce from third countries.

**Variability Factor:** A value that describes the variation in residue levels between the highest unit level and the average level in samples made up of many units. Internationally this is agreed to be the 97.5th percentile unit residue level divided by the average of the sum. The variability factor multiplied by the measured residue level from a composite sample (i.e., a sample made up by mixing several units before analysis) gives an estimate of the likely higher residue levels that may have occurred in individual units. These estimated higher levels are used in short-term risk assessments involving fruit and vegetables where consumers eat only a portion of a single item, e.g., melon, or a small number of units e.g., apples and potatoes.