



# Weekly Influenza and COVID-19 Surveillance graphs

UKHSA publishes a weekly national influenza and COVID-19 surveillance report which summarises the information from the surveillance systems which are used to monitor influenza, COVID-19 and other seasonal respiratory viruses in England.

Additional figures based on these surveillance systems are included in this slide set.

The figures presented in this slide set are based on data from week 35 (between 28 August and 04 September 2022).



# Contents

- 1) [COVID-19 Pandemic Overview](#)
- 2) [Confirmed COVID-19 episodes in England](#)
- 3) [Respiratory Datamart system \(England\)](#)
- 4) [Second generation surveillance system \(SGSS\)](#)
- 5) [Community surveillance](#)
- 6) [Surveillance in 'educational-age' cohorts](#)
- 7) [Secondary Care surveillance](#)
- 8) [Mortality surveillance](#)
- 9) [Possible reinfections in England](#)
- 10) [Co/secondary infections with COVID-19](#)




UK Health  
Security  
Agency

# COVID-19 Pandemic Overview




# Confirmed COVID-19 episodes tested under Pillar 1, by sample week, since week 5 2020






# Confirmed COVID-19 episodes tested under Pillar 2, by sample week, since week 5 2020






# Weekly overall hospital and ICU/HDU admission rates per 100,000 of new COVID-19 cases reported through SARI Watch, England since week 12 2020





# Number of deaths since week 10 2020 by week of death and time since laboratory confirmation of COVID-19, England



Since 1 April 2022, free universal symptomatic and asymptomatic testing for the general public in England is no longer available, as outlined in the plan for [living with COVID-19](#). Data should be interpreted in the context of this change to testing.



UK Health  
Security  
Agency

# Confirmed COVID-19 episodes in England






## Confirmed COVID-19 episodes in England

### Data Information

- From the week 32 report onwards, case rates have been updated to use the latest ONS population estimates for mid-2020. Previously case rates were calculated using the mid-2019 population estimates
- From 11 January 2022 the requirement for [confirmatory PCR testing in individuals who test positive using a lateral flow device was temporarily removed](#).
- Rates by ethnicity and IMD quantile will continue to be presented using the mid-2019 estimates, until the mid-2020 estimates become available.
- From 31 January 2022, UKHSA moved all COVID-19 case reporting in England to use a new episode-based definition which includes possible reinfections. Each infection episode is counted separately if there are at least 91 days between positive test results (PCR or LFD). Each infection episode begins with the earliest positive specimen date. Further information can be found on the [UK COVID-19 dashboard](#).
- Since 1 April 2022, free universal symptomatic and asymptomatic testing for the general public in England is no longer available, as outlined in the plan for [living with COVID-19](#). As such, there will be a reduction in the reporting of data obtained through Pillar 2 from April 2022 onwards. Data in this report should be interpreted in the context of this change to testing. [Public health guidance](#) remains in place for cases and their close contacts




## Confirmed COVID-19 episodes tested under Pillar 2, based on sample week with overall weekly PCR positivity for Pillar 2 (%)






## Weekly confirmed COVID-19 case rates per 100,000, by episode, tested under Pillar 2, by sex






## Weekly confirmed COVID-19 case rates per 100,000, by episode, tested under Pillar 2, by age group






## Weekly PCR positivity (%) of confirmed COVID-19 cases tested overall and by sex under Pillar 2






## Weekly PCR positivity (%) of confirmed COVID-19 cases tested under Pillar 2, by male and age group






## Weekly PCR positivity (%) of confirmed COVID-19 cases tested under Pillar 2, by female and age group






## Weekly confirmed COVID-19 case rates by episode, per 100,000 population (Pillar 2), by UKHSA centres and sample week








## Weekly PCR positivity of confirmed COVID-19 cases tested under Pillar 2 (%) by UKHSA centres and sample week






## Weekly incidence per 100,000 population by ethnicity (Pillar 2), England






# Weekly COVID-19 episodes tested under Pillar 1, per 100,000 population by age group and region, weeks 26 to 35






## Weekly COVID-19 episodes tested under Pillar 2, per 100,000 population by age group and region, weeks 26 to 35






## Weekly COVID-19 episodes tested under Pillar 1, per 100,000 population by ethnicity and region, weeks 26 to 35






## Weekly COVID-19 episodes tested under Pillar 2 per 100,000 population by ethnicity and region, weeks 26 to 35






## Weekly COVID-19 rate tested under Pillar 1, per 100,000 population by IMD quintile (1 being the most deprived and 5 being the least deprived)



\*Incidence rates have been calculated using the mid-2019 ONS population estimates




## Weekly COVID-19 rate tested under Pillar 2, per 100,000 population by IMD quintile (1 being the most deprived and 5 being the least deprived)








## Weekly rate of COVID-19 episodes per 100,000 population (Pillar 2), by upper-tier local authority, England (box shows enlarged map of London area)






# Cumulative rate of COVID-19 episodes per 100,000 population tested under Pillar 1, by upper-tier local authority, England (box shows enlarged map of London area)





## Cumulative rate of COVID-19 episodes per 100,000 population tested under Pillar 2, by upper-tier local authority, England (box shows enlarged map of London area)










UK Health  
Security  
Agency

# Respiratory Datamart system (England)




## Respiratory DataMart – Influenza subtypes











## Respiratory DataMart – Respiratory syncytial virus (RSV) weekly positivity by UKHSA region







## Respiratory DataMart – other respiratory viruses








# Second generation surveillance system (SGSS)

# SGSS reported Influenza A cases by region (all ages)




The presented figures are based on laboratory reports through SGSS. Testing and reporting procedures vary by virus, UKHSA centre and over time, including short-term trends in testing. Therefore

8 September 2020 comparisons should be done with caution.

Previously, this data was presented by report date however is now presented by specimen date.

## SGSS reported Influenza B cases by region (all ages)




The presented figures are based on laboratory reports through SGSS. Testing and reporting procedures vary by virus, UKHSA centre and over time, including short-term trends in testing. Therefore comparisons should be done with caution.

Previously, this data was presented by report date however is now presented by specimen date.




## SGSS reported RSV cases by region (all ages)



The presented figures are based on laboratory reports through SGSS. Testing and reporting procedures vary by virus, UKHSA centre and over time, including short-term trends in testing. Therefore comparisons should be done with caution. Previously, this data was presented by report date however is now presented by specimen date.




# SGSS reported Adenovirus cases by region (all ages)






# SGSS reported Parainfluenza cases by region (all ages)



The presented figures are based on laboratory reports through SGSS. Testing and reporting procedures vary by virus, UKHSA centre and over time, including short-term trends in testing. Therefore comparisons should be done with caution.




## SGSS reported Rhinovirus cases by region (all ages)



The presented figures are based on laboratory reports through SGSS. Testing and reporting procedures vary by virus, UKHSA centre and over time, including short-term trends in testing. Therefore comparisons should be done with caution.



## SGSS reported hMPV cases by region (all ages)



The presented figures are based on laboratory reports through SGSS. Testing and reporting procedures vary by virus, UKHSA centre and over time, including short-term trends in testing. Therefore comparisons should be done with caution.





UK Health  
Security  
Agency

# Community surveillance




## Acute respiratory infection (ARI) outbreaks linked to educational settings

### Data Information

We report on new acute respiratory infection (ARI) incidents reported to Health Protection Teams (HPTs) and entered on HPZone in the previous reporting week in educational settings by locality. The incidents captured on HPZone represent a subset of all ongoing clusters and outbreaks in England. A variety of arrangements are in place with local authorities and other stakeholders supporting HPTs, however data may not routinely be documented on HPZone. As a result, the number of outbreaks reported for some of the regions are underestimates



# Number of acute respiratory infection outbreaks reported to UKHSA by type of educational setting, England





# Number of acute respiratory infection outbreaks by type of educational setting, England

End of academic year total

Week 36 2020- 34 2021

PHE Centres	Cumulative number of suspected acute respiratory infection outbreaks by type of educational setting for the 2020/21 academic year Week 36 2020- 34 2021						
	Nursery	Primary School	Secondary School	Combined	Special Educational Needs (SEN) schools	College University	Total
<b>Total</b>	<b>846</b>	<b>2125</b>	<b>2122</b>	<b>40</b>	<b>666</b>	<b>268</b>	<b>6067</b>

Week 35 2022

Main table

PHE Centres	Cumulative number of suspected acute respiratory infection outbreaks by type of educational setting for the 2021/22 academic year from Week 35 2021						
	Nursery	Primary School	Secondary School	Combined	Special Educational Needs (SEN) schools	College University	Total
East Midlands Centre	72 (0)	59 (0)	30 (0)	14 (0)	170 (0)	6 (0)	<b>351 (0)</b>
East of England Centre	1 (0)	12 (0)	8 (0)	3 (0)	11 (0)	2 (0)	<b>37 (0)</b>
London Centre	370 (0)	1093 (0)	260 (0)	59 (0)	227 (0)	30 (0)	<b>2038 (0)</b>
North East Centre	0 (0)	2 (0)	0 (0)	0 (0)	3 (0)	0 (0)	<b>5 (0)</b>
North West Center	13 (0)	31 (0)	13 (0)	4 (0)	127 (0)	7 (0)	<b>195 (0)</b>
South East Centre	43 (0)	389 (0)	127 (0)	34 (0)	291 (0)	7 (0)	<b>891 (0)</b>
South West Centre	5 (0)	65 (0)	79 (0)	37 (0)	256 (0)	1 (0)	<b>441 (0)</b>
West Midlands Centre	19 (0)	74 (0)	52 (0)	7 (0)	142 (0)	6 (0)	<b>300 (0)</b>
Yorkshire & the Humber	17 (0)	36 (0)	27 (0)	5 (0)	84 (0)	0 (0)	<b>169 (0)</b>
<b>Total</b>	<b>540 (0)</b>	<b>1761 (0)</b>	<b>596 (0)</b>	<b>161 (0)</b>	<b>1306 (0)</b>	<b>59 (0)</b>	<b>4423 (0)</b>

\* Number of acute respiratory infection for the most recent week in brackets





UK Health  
Security  
Agency

# Secondary Care surveillance



## Weekly admission rates for hospital and ICU/HDU laboratory confirmed COVID-19 cases reported through SARI Watch, week 35




Source: UKHSA SARI-Watch (Severe Acute Respiratory Infection-Watch, formerly CHES).

\*Only NHS Acute trusts that have reported ≥1 day in the past week; excludes Specialist trusts. Acute NHS trusts (including Specialist trusts) reporting into SARI-Watch COVID-19 hospitalisation surveillance are typically around 100 per week. This was 87 for the hospitalisation (all levels of care) indicator in week 29 August 2022 to 04 September inclusive and 77 trusts for the ICU/HDU indicator. For the maps, as Specialist trusts are excluded, the number of trusts providing data on COVID-19 hospitalisations in week ending 04 September 2022 was 78 and 70 for ICU/HDU admissions for COVID-19.




# Age/sex pyramid of hospitalisations (all levels of care) for COVID-19, data from sentinel acute NHS trusts, England

(a) Peak of 2<sup>nd</sup> wave (week 53 2020 to week 3 2021) n= 6,359



Reporting trusts=22

(b) Most recent 4 weeks (week 32 2022 to 35 2022) n= 703




Reporting trusts=13



# Age/sex pyramid for admissions to ICU/HDU for COVID-19, mandatory


UK Health Security Agency  
case level data, acute NHS trusts, England

(a) Peak of 2<sup>nd</sup> wave (week 53 2020 to week 3 2021) n= 3,349



Reporting trusts=70

(b) Most recent 4 weeks (week 32 2022 to week 35 2022) n=109




Reporting trusts=29






# Laboratory confirmed admissions for COVID-19, to acute NHS trusts, by level of care and ethnicity

(a) Peak of 2<sup>nd</sup> wave (week 53 2020 to  
week 3 2021)



Reporting trusts  
Lower level of care=21  
ICU/HDU=68


(b) Most recent 4 weeks (week 32 2022  
to 35 2022)



Reporting trusts  
Lower level of care=13  
ICU/HDU=26




# Weekly COVID-19 hospitalisation rate per 100,000 trust catchment population by age group and region, weeks 26 to 35






## Hospital admission rate (excluding ICU/HDU) by ethnicity per 100,000 trust catchment population



Caveat: From week 24 (2021) the ethnicity analysis is based on a new method for assigning ethnicity, developed by UKHSA. The previous method used the most recent ethnicity recorded through linkage to Hospital Episode Statistics. However, this method led to unfeasibly high rates in the 'Other' ethnic group when applied to COVID-19 cases, hospitalisation or mortality. The new method uses the most frequent ethnicity recorded through linkage to Hospital Episode Statistics, unless the most frequent was 'Other' when the second most frequent was chosen.




## Rate of admission to ICU/HDU by ethnicity, per 100,000 trust catchment population



Caveat: From week (24 2021) the ethnicity analysis is based on a new method for assigning ethnicity, developed by UKHSA. The previous method used the most recent ethnicity recorded through linkage to Hospital Episode Statistics. However, this method led to unfeasibly high rates in the 'Other' ethnic group when applied to COVID-19 cases, hospitalisation or mortality. The new method uses the most frequent ethnicity recorded through linkage to Hospital Episode Statistics, unless the most frequent was 'Other' when the second most frequent was chosen.



# COVID-19 as primary reason for admission among SARS-CoV-2 positive patient by week of admission



## Notes

- 1) Case-level sentinel data from SARI-Watch, from week 35 2021 (commencing 30 August 2021) to week 33 2022 (ending 21 August 2022) inclusive
- 2) Total 27104 records in period of analysis, of which 35% (n=9498) had COVID-19 as primary reason for admission ('Yes').
- 3) SARS-CoV-2 patients with evidence of COVID-19 treatment but have 'No' or 'Unknown' for COVID-19 as primary reason for admission (n=924) are reassigned to COVID-19 as primary reason of admission ('Yes').
- 4) Reassignment increases COVID-19 as primary reason for admission ('Yes') from 9498 to 10422
- 5) 23% (6327/27104) of total records in this period have missing data on the 'Admission due to COVID-19' indicator – these are excluded from analysis
- 6) Caveats: London trusts under-represented and most recent weeks are subject to retrospective updates




UK Health  
Security  
Agency

# Mortality surveillance



# Cumulative mortality rate of COVID-19 cases per 100,000 population tested under Pillar 1 and 2 since the beginning of the pandemic by 28 day definition





UK Health  
Security  
Agency

# Co/secondary infections with COVID-19





## Co/secondary infections with COVID-19

Caveat - undertesting for other pathogens may result in an underestimate of co/secondary infection cases.

Preceding/co-/secondary infections refers to when a patient has a COVID-19 infection with one or more other pathogen (Please see Appendix 1 – Pre-/co-/secondary infection with COVID-19 definitions.)

- Preceding infection: COVID-19 acquired after another pathogen
- Co-infection: COVID-19 and other pathogen acquired at the same time
- Secondary infection: COVID-19 acquired before another pathogen

Numbers of pre-/co-/secondary infection remain low across UKHSA surveillance systems.

For patients with severe respiratory failure requiring Extra Corporeal Membrane Oxygenation (ECMO), analysis of data from five adult ECMO centres in England indicates that among patients with severe respiratory failure due to COVID-19, almost a third of these have co/secondary infections.

Published data analysis from pandemic wave 1 (W-1) indicates increased mortality associated with COVID-19 and [influenza](#), [key bacterial and fungal infections](#) and [invasive pneumococcal disease \(IPD\)](#) in comparison to patients without co/secondary infection.

[Data analysis](#) from W-1 indicates that *Aspergillus* and *candidemia* cases have increased risk of mortality in comparison to patients without co/secondary infection.



## Co/secondary infections among patients with severe respiratory failure requiring Extra Corporeal Membrane Oxygenation (ECMO)

Analysis is based on cumulative data from five adult ECMO centres in England. Data for the current and previous seasons are presented. Each season commences around October (ISO week 40) ending in September in the following year (ISO week 39).

Data for the current season (2021-22) is from 4 October 2021 to 15 May 2022 inclusive (week 40 2021 to week 19 2022). This period includes effects from the Delta and Omicron waves of the pandemic. The 2020-21 season is from 28 September 2020 to 3 October 2021 inclusive and includes effects from the Alpha and Delta waves. The 2019-20 season is from 30 September 2019 to 27 September 2020 inclusive and includes effects from the original Wuhan strain.

In the 2021-22 season, 41% (31/76) of ECMO patients admitted for severe respiratory failure due to laboratory confirmed COVID-19 had clinically significant co/secondary infections. In the previous season (2020-21) this proportion was 30% (134/402). In the 2019-20 season this proportion was 33% (79/236).

In all three seasons the majority of clinically significant co/secondary infections among respiratory failure COVID-19 cases comprised Gram-negative bacilli from the order Enterobacterales:

- 45% (14/31) in the current season 2021-22
- 32% (43/134) in 2020-21 and 46% (36/79) in 2019-20. The decrease in 2020-21 compared to 2019-20 reached borderline significance ( $p=0.057$ ). (In the last data assessment, the decrease over this period was statistically significant but since then there were further updates to the data from ECMO centres). No change was detected in other key pathogens between these two time periods.



UK Health  
Security  
Agency

# Surveillance of bacterial, fungal and viral infections, in COVID-19 patients in England, Jan 2020 – June 2022

HCAI, Fungal, AMR, AMU & Sepsis Division



# Updates

From 31 January 2022, UKHSA has changed the COVID-19 case definition to include multiple infection episodes. Reported co-/secondary/preceding infections in England now use the new definition, revising all cases back to the beginning of the pandemic.

The Unified Infection Dataset (UID) project has been extended to incorporate the Co- and Secondary infections with COVID-19 datasets

The following outputs included in this section have been produced via the UID, combining previously separate data pipelines

- Key HCAI bacterial and fungal specimens reported to SGSS and HCAI data capture system
- Respiratory viral specimens reported to SGSS and Respiratory Datamart
- Fungal specimens reported to mycology reference lab (MRL)

The Co- and secondary infections team have undertaken an extensive data validation exercise which has identified additional respiratory viral specimens from Respiratory Datamart and allowed us to make improvements to the methodologies. Preceding infections for all pipelines (other pathogen infections occurring before COVID-19 specimen) are now included. Please note, all cases since January 2020 have been revised in line with this validation.

Data are provisional and subject to change due to possible delayed reporting of microbiological samples



# Number of COVID-19 patient-episodes with bacterial, fungal or viral infections in COVID-19 patients diagnosed in England during wave 3\*, by infection type and timing of diagnosis

Bacterial/ fungal/ viral infection by specimen type	COVID-19 patient-episodes with bacterial/ fungal/ viral infection		Timing of bacterial/fungal/viral diagnosis in relation to COVID-19 diagnosis								
			Preceding infection			Coinfection			Secondary infection		
	n	% of COVID cases	n	% infections by site	% of COVID cases	n	% infections by site	% of COVID cases	n	% infections by site	% of COVID cases
Bacterial/fungal bloodstream & lower respiratory infection	255	<0.01	23	9.02	<0.01	9	3.53	<0.01	223	87.45	<0.01
Bacterial/fungal bloodstream infection	8,189	0.06	3,477	42.46	0.02	1,631	19.92	0.01	3,081	37.62	0.02
Bacterial/fungal lower respiratory infection	2,646	0.02	622	23.51	<0.01	250	9.45	<0.01	1,774	67.04	0.01
<i>Clostridioides difficile</i> infection	1,040	0.01	419	40.29	<0.01	102	9.81	<0.01	519	49.90	<0.01
Fungal respiratory/bloodstream infection (MRL)‡	198	<0.01	14	7.07	<0.01	3	1.52	<0.01	181	91.41	<0.01
Other respiratory virus infection	2,980	0.02	827	27.75	0.01	1,390	46.64	0.01	763	25.60	0.01
Any site†	15,345	0.10	5,394	35.15	0.04	3,386	22.07	0.02	6,565	42.78	0.04

## Key findings:

- 0.1% of COVID-19 patient-episodes had a bacterial, fungal or other respiratory viral infection detected in either the 28 days prior or following (60 days following for MRL) their COVID-19 diagnosis
- Prevalence in W3 lower than W2 and W1; however, patient-episodes of COVID-19 and another key infection are still higher in W3 than W1 (15,345 vs 4,636, respectively)
- Most infections with key organisms were categorised as secondary infections (42.78%).

Please see appendix 1 for pre-/co-/secondary infection definitions with SARS-CoV-2

Please note patients can have multiple COVID-19 infection-episodes, numbers here do not reflect the number of patients.


\*SARS-CoV2 specimen dates from 27 Apr 2021 to 22 May 2022 (N=14,778,196). Last updated 22 Jun 2022.

‡ Definition for secondary infection differs for MRL specimens - detection within 60 days

† includes the combination Bacterial/fungal bloodstream & *Clostridioides difficile* infection (12 preceding, 1 coinfection & 20 secondary), Bacterial/fungal bloodstream, lower respiratory & *Clostridioides difficile* infection (1 secondary), & Bacterial/fungal lower respiratory & *Clostridioides difficile* infection (3 secondary)



## Most frequent bacterial/fungal species in blood or lower respiratory tract specimens, by timing of diagnosis, in COVID-19 patients diagnosed in England during wave 3




### Key findings:

In wave 3, the most frequent bacterial/fungal organisms identified from blood specimens were *Escherichia coli*, *Staphylococcus aureus* and *Klebsiella pneumoniae* and from respiratory specimens were *Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Klebsiella pneumoniae*.



## Most frequent viral specimens, by timing of diagnosis, in COVID-19 patients diagnosed in England during wave 3




### Key findings:

In wave 3, the most frequent viral organisms identified from respiratory specimens were RSV, rhinovirus and influenza A.



## Most frequent fungal species (MRL), by timing of diagnosis, in COVID-19 patients diagnosed in England during wave 3



### Key findings:

In wave 3, the most frequent fungal organisms identified were *Aspergillus fumigatus* complex and *Candida albicans*.





Bacteria/Fungi	First Wave	Second Wave	Third Wave	Total Cases
	(30 Jan 2020 - 28 June 2020)	(29 June 2020 – 30 April 2021)	(1 May 2021 – 24 February 2022)	
<i>Aspergillus fumigatus</i> isolates (azole resistant)	46 (4)	120 (2)	137 (12)	303 (18)
Probable/Proven cases of CAPA*	15	38	44	97
<i>Candida</i> spp.: Candidemia	63	133	17	213
<i>Bordetella pertussis</i>	0	0	0	0
<i>Haemophilus influenzae</i>	3	2	0	5
<i>Neisseria meningitidis</i>	2	0	0	2
<i>Streptococcus pneumoniae</i>	40	45	14	99

\*COVID-19-associated pulmonary aspergillosis

Please note fungal data refers to secondary infections only. Mycology data contains results from Mycology reference laboratory data, Candidaemia is representative of deep infection. One case of osteomyelitis, one case of ventriculitis and one case of endocarditis was documented in wave two. Fungal data are also included in the overall numbers in slides 6-8 but have been stratified here with additional details. *Bordetella pertussis* co-infection is defined as +/- 28 d Culture/PCR (based on pertussis sample date), +/- 28 Serology/Oral fluid (anti-pertussis toxin Ig) (based on pertussis symptom onset date, excluding cases without onset date). *Haemophilus influenzae*, *Neisseria meningitidis* and *Streptococcus pneumoniae* co-infection is defined as +/- 2d. *Legionella*, *Mycoplasma* and gastrointestinal infection data not included. Please note, testing in W1 was not open to the community and therefore W1 cases are predominantly hospitalised patients vs. W2 and W3.

# Appendix 1: Pre-/co-/secondary infection definitions with COVID-19

The day pertains to the date of the sample collection that yielded a positive result. These definitions do not apply to persistent COVID-19 patients. Patients with persistent COVID-19 require independent clinical assessment.

Organism	Definition co-infection with SARS-CoV-2 †	Definition of infection pre-SARS-CoV-2 infection (other pathogen is primary infection) or Definition of post SARS-CoV-2 secondary infection (SARS-CoV-2 is primary infection)
Influenza A	+/- 1d	2-28d <sup>^</sup>
Influenza B	+/- 1d	2-28d <sup>^</sup>
RSV	+/- 1d	2-28d
Adenovirus	+/- 1d	2-28d
Enterovirus	+/- 1d	2-28d
Human metapneumovirus	+/- 1d	2-28d
Parainfluenza (any subtype)	+/- 1d	2-28d
Seasonal coronavirus	+/- 1d *	2-28d
Rhinovirus	+/- 1d	2-28d
Co-infections in ECMO patient (patients with most severe clinical respiratory signs)		
ECMO patients	Individual case review	Individual case review
Blood stream and respiratory infections (bacterial and fungal)		
<i>Achromobacter xylosoxidans</i>	+/- 1d	2-28d
<i>Acinetobacter</i> spp.,	+/- 1d	2-28d
<i>Aspergillus</i>	+/- 1d	2-28d (pre) 2-60d (post, continually hospitalised patients only)
<i>Bordetella pertussis</i>	+/- 28 d Culture/PCR (based on pertussis sample date) +/- 28 Serology/Oral fluid (anti-pertussis toxin Ig) (based on pertussis symptom onset date, excluding cases without onset date)	N/A (Pertussis presentation is often delayed)
<i>Burkholderia cepacia</i>	+/- 1d	2-28d
<i>Candida</i> spp	+/- 1d	2-28d (pre) 2-60d (post, continually hospitalised patients only)
<i>Chlamydia pneumoniae</i>	0-7d PCR	PCR within 14-28 d (8-13d PCR <sup>*</sup> )
<i>Enterobacter</i> spp.,	+/- 1d	2-28d
<i>Enterococcus</i> spp.	+/- 1d	2-28d
<i>E. coli</i>	+/- 1d	2-28d
<i>Haemophilus influenzae</i>	+/- 2d	3-28d

Continued overleaf

# Appendix 1 continued: Pre-/co-/secondary infection definitions with COVID-19

The day pertains to the date of the sample collection that yielded a positive result. These definitions do not apply to persistent COVID-19 patients. Patients with persistent COVID-19 require independent clinical assessment.

Organism	Definition co-infection with SARS-CoV-2 †	Definition of infection pre-SARS-CoV-2 infection (other pathogen is primary infection) or Definition of post SARS-CoV-2 secondary infection (SARS-CoV-2 is primary infection)
Blood stream and respiratory infections (bacterial and fungal)		
<i>Klebsiella</i> spp.	+/- 1d	2-28d
<i>Legionella pneumophila</i> /species	Individual case review	Individual case review
<i>Mycoplasma pneumoniae</i>	0-7d PCR, IgM serology 0-21d <16y	PCR within 14-28 d (8-13d PCR*)
<i>Neisseria meningitidis</i>	+/- 2d	3-28d
<i>Pseudomonas</i> spp.,	+/- 1d	2-28d
<i>Serratia</i> spp.,	+/- 1d	2-28d
<i>Staphylococcus aureus</i>	+/- 1d	2-28d
Coag-neg <i>Staphylococcus</i> (S. <i>haemolyticus</i> )	+/- 1d	2-28d
<i>Stenotrophomonas</i> spp., (S. <i>maltophilia</i> )	+/- 1d	2-28d
<i>Streptococcus</i> spp. ‡	+/- 1d	2-28d
<i>Streptococcus pneumoniae</i>	+/- 2d	3-28d
Tuberculosis		
<i>Mycobacterium tuberculosis</i>	Individual case review	Individual case review
Pathogens of the immunocompromised (eg HIV)		
HIV	Individual case review	Individual case review
Gastrointestinal infections		
Listeria	0-5d *	Individual case review
Campylobacter	0-5d *	Individual case review
Shiga toxin-producing <i>E. coli</i> (STEC)	0-5d *	Individual case review
Norovirus	0-5d *	Individual case review
Salmonella	0-5d *	Individual case review
Shigella	0-5d *	Individual case review
Anaerobes		
<i>C. difficile</i>	+/- 1d	2-28d
<i>Bacteroides</i> sp. ( <i>B. fragilis</i> and non- <i>fragilis</i> <i>Bacteroides</i> )	+/- 1d	2-28d

See next slides for notes

# Appendix 1 continued: Pre-/co-/secondary infection definitions with COVID-19

## Notes

† From SARS-CoV-2 first detection date. Not including multiple episodes of SARS-CoV-2 per patient.

\* Additional data check required. (Resistance is not detailed, data for MERS is not currently available).

^ Definition post- SARS-CoV-2 secondary infection (SARS-CoV-2 is primary infection). This has been extended from prior 14d secondary infection definition for influenza used by PHE to account for disparities in testing throughout the 28d period after SARS-CoV-2 detection.

‡ Streptococcus species includes the following groups and species:

Group	Species/other names
Anginosus Group	<i>Streptococcus anginosus</i> ; <i>Streptococcus constellatus</i> ( <i>Streptococcus constellatus</i> subspecies <i>constellatus</i> <i>Streptococcus constellatus</i> subspecies <i>pharynges</i> ); <i>Streptococcus</i> Group F; <i>Streptococcus intermedius</i> ; <i>Streptococcus milleri</i> group; <i>Streptococcus sinensis</i>
Bovis Group	<i>Streptococcus alactolyticus</i> ; <i>Streptococcus bovis</i> untyped; <i>Streptococcus equinus</i> ; <i>Streptococcus gallolyticus</i> subspecies <i>gallolyticus</i> ( <i>Streptococcus bovis</i> biotype I); <i>Streptococcus infantarius</i> ( <i>Streptococcus infantarius</i> sp <i>infantarius</i> ; <i>Streptococcus bovis</i> biotype II); <i>Streptococcus lutetiensis</i> ; <i>Streptococcus infantarius</i> subspecies <i>coli</i> ( <i>Streptococcus bovis</i> biotype II); <i>Streptococcus pasteurianus</i> ( <i>Streptococcus bovis</i> biotype II)
Closely Related Genera	<i>Abiotrophia</i> spp.; <i>Aerococcus</i> spp.; <i>Faklamia</i> spp.; <i>Gemella</i> spp.; <i>Globicatella sanguinis</i> ; <i>Granulicatella</i> spp.; <i>Leuconostoc</i> spp.; <i>Pedicoccus</i> spp.; <i>Peptostreptococcus</i> spp.
Mitis Group	<i>Streptococcus cristatus</i> ; <i>Streptococcus mitior</i> ; <i>Streptococcus mitis</i> ; <i>Streptococcus oralis</i> ; <i>Streptococcus pseudopneumoniae</i> ; <i>Streptococcus infantis</i> ; <i>Streptococcus peroris</i>
Mutans Group	<i>Streptococcus mutans</i> ; <i>Streptococcus sobrinus</i>
Other streptococci (including but not limited to)	Anaerobic streptococcus; <i>Streptococcus acidominimus</i> ; <i>Streptococcus</i> spp., other named/not fully identified; <i>Streptococcus suis</i> ; <i>Streptococcus uberis</i>
Salivarius Group	<i>Streptococcus vestibularis</i> ; <i>Streptococcus thermophilus</i>
Sanguinis Group	<i>Streptococcus gordonii</i> ; <i>Streptococcus massiliensis</i> ; <i>Streptococcus parasanguinis</i> ; <i>Streptococcus sanguinis</i>
<i>Streptococcus</i> Group A	Group A; <i>Streptococcus pyogenes</i> ; <i>Streptococcus dysgalactiae</i> subspecies <i>equisimilis</i>
<i>Streptococcus</i> Group B	Group B; <i>Streptococcus agalactiae</i>
<i>Streptococcus</i> Group C	Group C; <i>Streptococcus dysgalactiae</i> subspecies <i>equisimilis</i> ; <i>Streptococcus equi</i> subspecies <i>zooepidemicus</i>
<i>Streptococcus</i> Group G	Group G; <i>Streptococcus canis</i> ; <i>Streptococcus dysgalactiae</i> subspecies <i>equisimilis</i>