

| Year of issue | Country | PL number | Shelf life | Storage conditions |
|---------------|-----------------------|---------------|---|---|
| 2019 | PALESTINIAN AUTHORITY | PL 44124/0024 | 3 years. After dilution: the physicochemical stability in glucose 5 %, sodium chloride 0.9 % solution, Ringer's solution or Ringer's acetate solution has been demonstrated for 24 hours at 25°C. From a microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user. | Store in a refrigerator (2°C – 8°C). Keep the ampoules in outer carton in order to protect from light. For storage conditions after dilution of the medicinal product, see section 6.3. |
| 2020 | HONG KONG | PL 16853/0021 | 3 years. | For prolonged periods store between 2°C and 8°C. Protect from light. Syntometrine may be stored up to 25°C for 2 months when protected from light, but must then be discarded. |
| 2020 | VIETNAM | PL 16853/0021 | 3 years | For prolonged periods store between 2°C and 8°C. Protect from light. Syntometrine may be stored |

| | | | | |
|------|-----------|---------------|---------|--|
| | | | | up to 25°C for 2 months when protected from light, but must then be discarded. |
| 2020 | MALAYSIA | PL 16853/0021 | 3 years | For prolonged periods store between 2 ^o and 8°C. Protect from light. Syntometrine may be stored up to 25°C for 2 months when protected from light, but must then be discarded. |
| 2020 | SINGAPORE | PL 16853/0021 | 3 years | For prolonged periods store between 2 ^o and 8°C. Protect from light. Syntometrine may be stored up to 25°C for 2 months when protected from light, but must then be discarded. |
| 2020 | VIETNAM | PL 44124/0024 | 3 years | After dilution: the physicochemical stability in glucose 5 %, sodium chloride 0.9 % solution, Ringer's solution or Ringer's acetate solution has been demonstrated for 24 hours at 25°C. From a microbiological point of view, the product should be used |

| | | | | |
|------|-------------------|---------------|-------------|---|
| | | | | immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user. |
| 2021 | BRUNEI DARUSSALAM | PL 16853/0021 | 3 years | For prolonged periods store between 2 ^o and 8°C. Protect from light. Syntometrine may be stored up to 25°C for 2 months when protected from light, but must then be discarded |
| 2021 | SAUDI ARABIA | PL 16853/0021 | 3 years | For prolonged periods store between 2 ^o and 8°C. Protect from light. Syntometrine may be stored up to 25°C for 2 months when protected from light, but must then be discarded. |
| 2021 | VIETNAM | PL 16853/0021 | 3 years | For prolonged periods store between 2 ^o and 8°C. Protect from light. Syntometrine may be stored up to 25°C for 2 months when protected from light, but must then be discarded |
| 2021 | SAUDI ARABIA | PL 40369/0006 | Three years | Store in a refrigerator (2°C - 8°C). May be |

| | | | | |
|------|--------------|---------------|-------------|--|
| | | | | stored below 25°C for 6 months, but then discarded. |
| 2021 | SAUDI ARABIA | PL 40369/0005 | Three years | Store in a refrigerator (2°C - 8°C). May be stored below 25°C for 6 months, but then discarded |
| 2022 | ALGERIA | PL 40369/0005 | Three years | Store in a refrigerator (2°C - 8°C). May be stored below 25°C for 6 months, but then discarded |
| 2023 | IRAN | PL 40369/0006 | Three years | Oxytocin should not be infused via the same apparatus as blood or plasma, because the peptide linkages are rapidly inactivated by oxytocin-inactivating enzymes. Oxytocin is incompatible with solutions containing sodium metabisulphite as a stabiliser. |
| | | | | |