

Title: Community water fluoridation expansion in north east England IA No: 9617 RPC Reference No: Lead department or agency: Department of Health & Social Care Other departments or agencies:	Impact Assessment (IA)
	Date: 4 March 2024
	Stage: Consultation
	Source of intervention: Domestic
	Type of measure: Other
	Contact for enquiries: waterfluoridationconsultation@dhsc.gov.uk
Summary: Intervention and Options	RPC Opinion: Not applicable

Cost of Preferred (or more likely) Option (in 2019 prices)			
Total Net Present Social Value	Business Net Present Value	Net cost to business per year	Business Impact Target Status
£315m	0	0	

What is the problem under consideration? Why is government action or intervention necessary?

Tooth decay is a significant yet largely preventable public health problem, affecting all stages of life; it is the most common oral disease in children, with significant health inequalities, can cause pain and difficulties eating and sleeping. It is progressive if left untreated but can be prevented by fluoride and minimising the amount and frequency of sugar consumption. The government proposes using powers in the Health & Care Act 2022 to expand community water fluoridation in the north-east region of England, increasing the levels of fluoride in the water supply, if needed, to levels needed to improve dental health. Legislation requires a public consultation before a final decision is made.

What are the policy objectives of the action or intervention and the intended effects?

There is strong scientific evidence that water fluoridation schemes can help reduce tooth decay and is a safe and effective public health intervention¹. Around 6 million people in England are already covered by agreements to add fluoride to their drinking water supplies, including parts of the north-east region. This proposal would expand existing community water fluoridation to an additional 1.6 million people. Implementation is intended to improve oral health of adults and children in the north-east region, reduce inequalities in oral health in the north-east region of England, and reduce impacts of poor oral health on dental services and the NHS in the north east region of England.

What policy options have been considered, including any alternatives to regulation? Please justify preferred option (further details in Evidence Base)

Impacts are compared to a 'do nothing' option where existing areas are fluoridated, and others are not. Existing costs continue and additional costs or risks of expansion will not be incurred. Alternative policy options considered include access to dental services and targeted oral health improvement programmes. In July 2022 the government announced a package of reforms to meet oral health need and increase access to dental care. In February 2024, the government and NHS England published a national plan for the recovery and reform of NHS dentistry services. Expansion of community water fluoridation in the north-east region is the preferred option. Water fluoridation is complementary to other interventions and does not rely on behaviour change, which is particularly important for children and vulnerable groups. The north east region is the first area under consultation as it has existing feasibility studies, large inequalities in oral health, and the water company has experience of water fluoridation.

Is this measure likely to impact on international trade and investment?	No
--	----

¹ See for example health monitoring reports 2014, 2018 and 2022 at [Water fluoridation: health monitoring report for England 2022 - GOV.UK \(www.gov.uk\)](https://www.gov.uk/government/reports/2022/water-fluoridation-health-monitoring-report-for-england-2022)

Are any of these organisations in scope?	Micro No	Small No	Medium No	Large Yes
What is the CO ₂ equivalent change in greenhouse gas emissions? (Million tonnes CO ₂ equivalent)	Traded: N/A		Non-traded: N/A	
Will the policy be reviewed? It will be reviewed. If applicable, set review date: Not applicable				

I have read the Impact Assessment and I am satisfied that, given the available evidence, it represents a reasonable view of the likely costs, benefits and impact of the leading options.

Signed by the responsible

Andrea Leadsem

Date: 4 March 2024

Description: Do Nothing

FULL ECONOMIC ASSESSMENT

Price Base Year	PV Base Year	Time Period	Net Benefit (Present Value (PV)) (£m)		
			Low:	High:	Best Estimate: 0
COSTS (£m)	Total Transition (Constant Price) Years		Average Annual (excl. Transition)		Total Cost (Present Value)
Low	Optional		Optional		Optional
High	Optional		Optional		Optional
Best Estimate					0
Description and scale of key monetised costs by 'main affected groups'					
These are defined to be 0.					
Other key non-monetised costs by 'main affected groups'					
These are defined to be 0.					
BENEFITS (£m)	Total Transition (Constant Price) Years		Average Annual (excl. Transition)		Total Benefit (Present Value)
Low					
High					
Best Estimate					0
Description and scale of key monetised benefits by 'main affected groups'					
These are defined to be 0.					
Other key non-monetised benefits by 'main affected groups'					
These are defined to be 0.					
Key assumptions/sensitivities/risks					Discount rate
We assume that no new water fluoridation schemes are introduced under the 'do nothing' option.					

BUSINESS ASSESSMENT (Option 1)

Direct impact on business (Equivalent Annual)			Score for Business Impact Target (qualifying provisions only) £m:
Costs: 0	Benefits: 0	Net: 0	

Description: Expansion of community water fluoridation in the north east region

FULL ECONOMIC ASSESSMENT

Price Base Year 2019	PV Base Year 2019	Time Period Years 40	Net Benefit (Present Value (PV)) (£m)		
			Low: £282m	High: £349m	Best Estimate: £315m

COSTS (£m)	Total Transition (Constant Price) Years	Average Annual (excl. Transition)	Total Cost (Present Value)
Low			Social cost: £261m (Financial cost: £44m)
High			Social cost: £328m (Financial cost: £56m)
Best Estimate			Social cost: £294m (Financial cost: £50m)

Description and scale of key monetised costs by 'main affected groups'

Expanding water fluoridation incurs discounted capital costs, estimated to be approximately £139m social cost (£26m financial), and operational costs, revenue associated with day to day running and maintenance, estimated to be approximately £1.1M per year. These are not direct costs to business; The Secretary of State is responsible for reimbursing water companies for reasonable costs associated with water fluoridation schemes, managed through an ongoing business as usual programme.

Other key non-monetised costs by 'main affected groups'

None

BENEFITS (£m)	Total Transition (Constant Price) Years	Average Annual (excl. Transition) (Constant Price)	Total Benefit (Present Value)
Low			
High			
Best Estimate			£610m

Description and scale of key monetised benefits by 'main affected groups'

In England water fluoridation schemes aim for a concentration of 1 mg per litre as strong scientific evidence has shown this to be the optimal level of fluoride for a temperate climate to reduce tooth decay. Reducing caries is expected to lead to dental healthcare cost savings from avoided fillings and extractions, productivity gains from workdays missed and quality of life benefits to the individual; an economic model monetises these over a 40 year time horizon.

Other key non-monetised benefits by 'main affected groups'

Other non-monetised benefits may include benefits accrued beyond the 40 year time horizon. The likely benefit associated with partial exposure of fluoridated water over the life course. The impact from missed education, avoided out of pocket expenses from paying for extractions or fillings (where applicable) and travelling to and from dental practices and forgone need for further dental healthcare such as root canals, dentures and bridges.

Key assumptions/sensitivities/risks**Discount rate (%)**

3.5/1.5%, (after 30 years 3.0/1.29%)

We assume no population growth within the north east region, and no change to dental caries over time. Any benefits accrued by individuals moving into the area would be equally and oppositely offset by individuals moving out of the area. Some model inputs rely on England data and we assume these hold for the north east region.

BUSINESS ASSESSMENT (Option 2)

Direct impact on business (Equivalent Annual)			Score for Business Impact Target (qualifying provisions only) £m:
Costs: £0	Benefits:	Net:	

Contents

Impact Assessment (IA)	1
Summary: Intervention and Options	1
Summary: Analysis & Evidence Policy Option 1	3
Summary: Analysis & Evidence Policy Option 2	4
Evidence Base	7
Problem under consideration.....	7
Tooth decay in England	7
Tooth decay in the north east region.....	8
Policy objective.....	8
Policy options	9
Do nothing option.....	9
Alternative options to reduce tooth decay	10
Summary and preferred option with description of implementation plan	11
Costs and benefits.....	11
Summary table of costs and benefits.....	11
Overview of economic model.....	12
Costs.....	13
Benefits.....	15
Summary of Analysis & Sensitivity Analysis	21
Risks.....	22
Specific Impact Tests	23
Impact on small and micro businesses	23
Equalities	23
Rural proofing	23
Competition assessment.....	23
Carbon assessment and sustainable development specific impact test / environmental impact	24
Human rights	24
Justice impact test	24
Potential trade implications of measure	24
Monitoring and Evaluation	24
Annex A: Modelling Approach	24
A1. Cohort approach.....	24
A2. DMFTs.....	26
A3. Dental healthcare savings	27
A4. Productivity gains	29
A5. Direct health benefit.....	30

Evidence Base

Problem under consideration

1. Tooth decay is a significant, yet largely preventable, public health problem in England. It is caused by consuming sugary food and drinks and lack of availability to fluoride. It affects people at all stages of life and is the most common oral disease in children. It can result in pain and cause people difficulties eating, sleeping and socialising. Tooth decay is a progressive disease which, if left untreated, can cause severe pain and infection.

Tooth decay in England

2. In 2022, 24% of 5 year olds in England had experienced tooth decay.¹ Those in the most deprived 20% of areas of the country were 2.5 times as likely to have experience of dental decay as those in the least deprived 20% of areas (Figure 1). There were also inequalities in the prevalence of experience of dental decay by ethnic group, which was significantly higher in the Other ethnic group (45%) and the Asian or Asian British ethnic group (38%) than for other groups.




Figure 1. Prevalence of experience of dental decay in 5 year olds in England, 2022 by national IMD 2019 quintiles

3. Tooth decay can have a detrimental effect on children’s learning and development as pain and infections from decayed teeth can result in school absences. In 2019, 6% of children aged under 16 in England had time off nursery or school in the last 6 months because of problems with their teeth, mouth or gums.²
4. In 2021/22, tooth extraction episode rates were more than 3 times higher in the most deprived 20% of areas in the country (329 per 100,000 population of 0 to 19 year olds) than the least deprived 20% of areas (95 per 100,000 population)³. Parents may also have to take time off work.
5. Tooth decay also has a considerable impact on the NHS and it is still the most common reason for hospital admission in children aged between 6 to 10 years. The costs to the NHS of hospital admissions for tooth extractions in children aged 0 to 19 years have been estimated based on the latest NHS national cost collection data. The costs were £81.0 million for all tooth extractions and £50.9 million for extractions due to tooth decay in the financial year 2021 to 2022.³

Tooth decay in the north east region

- During 2021 to 2022, the number of Finished Consultant Episodes for children and adolescents aged 0-19 for hospital tooth extractions, with caries as primary diagnosis per 100,000 was 369 per 100,000 population in the north east region compared to 378 per 100,000 in Yorkshire and The Humber region, the highest rate in the country and 205 per 100,000 for England³.
- In 2022, the proportion of teeth with experience of dentinal decay that had been extracted in 5 year olds was highest in the north east region (13.4%), compared to 6.4% across England overall (see Figure 2).




Figure 2. Proportion of teeth with experience of dentinal decay that have been extracted in 5 year olds in England by region, 2022

- The 2009 Adult Dental Health Survey⁴ showed information on adult dental health in England, and by region in England; north east region estimates are presented alongside the range across England regions for comparison:
 - 30% of adults in England and 34% of adults in the north east region had decayed teeth. Regionally this ranged from 21% in the south east coast to 39% in the west midlands.
 - 10% of adults with one or more natural teeth in the north east region had excellent oral health (defined as 21 or more teeth, 18 or more sound and untreated teeth, no active decay at any site, no periodontal pocketing or loss of attachment above 4mm, and no plaque or calculus); in England, this varied from 4% in the west midlands to 20% in the east of England.
 - 9% of adults in England and 8% of adults with teeth in the north east region reported feeling current pain related to their teeth. Regionally this ranged from 7% in the south east coast region to 11% in the north west.

Policy objective

- The policy objectives are to reduce tooth decay in children and adults in the north east region, to reduce oral health inequalities in the north east region of England and to reduce pressures on dental services and the NHS in the north east region. This is based on existing oral health inequalities, local support and the availability of an operable and efficient scheme

proposal. In this instance, oral health is defined in terms of tooth decay. The indicators of success will be reduced tooth decay amongst the population.

Policy options

Do nothing option

10. The 'do nothing' option for this policy is the baseline with which the preferred option expanding water fluoridation in the north east region is compared. In the 'do nothing' scenario, no new water fluoridation schemes are introduced in the north east region. There would be no costs or benefits in this scenario. Government actions to improve access to NHS dentistry, reduce sugar consumption would continue and have a positive effect on improving oral health. Other locally led oral health improvement work would continue based on local decision making.

Expanding community water fluoridation

11. The preferred policy option is to expand water fluoridation schemes in the north east region. There are already water fluoridation schemes in operation in some areas of the north east region, but the preferred option would be to expand them to a wider area.

12. Fluoride is a naturally occurring mineral found in soil, food and drink and also in drinking water supplies, in varying amounts. In some parts of England the level of fluoride in the public water supply already reaches the target concentration of water fluoridation schemes (one milligram per litre (1mg/l)), as a result of the geology of the area. In other areas the fluoride concentration has been adjusted to reach this level as part of a fluoridation scheme. In England water fluoridation schemes aim for a concentration of 1 mg per litre as this is the optimal level of fluoride to reduce tooth decay.

13. No new schemes have been implemented for over 40 years. From 2013 local authorities had the responsibility, through the Water Industry Act 1991, to propose and consult on new fluoridation schemes and variations to or termination of existing schemes. However, local authorities reported difficulties with that process. In particular, local authority boundaries were not coterminous with water flows which requires the involvement of several authorities in these schemes and made development of arrangements more complex. The Health and Care Act 2022 transferred to the Secretary of State of Health and Social Care the power to directly introduce, vary or terminate water fluoridation. Changes to water fluoridation schemes are subject to statutory consultation procedures.

14. The proposal to expand water fluoridation should be seen as a complementary strategy and not a substitute for other effective methods of increasing fluoride use. Water fluoridation would help to achieve the policy objective, as it would be a universal intervention to improve health for a whole area. It does not rely on behaviour change which is particularly important for children and vulnerable groups.

15. An alternative policy option would be to expand water fluoridation schemes to cover different areas, more or all of England. This is not favoured at the present time and the north east region has been prioritised as the first area under consultation to introduce new, or vary existing, water fluoridation arrangements using powers in the Health and Care Act 2022, based on several factors: in the north east region, necessary feasibility studies were well developed under the previous legislative framework for the whole area; they were also close to going out to public consultation when the legislation changed; there are large inequalities in oral health across the north east region; and the water company supplying the area has experience in water fluoridation.

Alternative options to reduce tooth decay

16. In 2014, Public Health England published an evidence-informed toolkit for local authorities to inform commissioning better oral health for children and young people.⁵ This outlined a range of potential interventions that could be introduced to improve oral health:
- **Interventions to support consistent evidence informed oral health information**, such as oral health training for the wider professional workforce and integrating oral health into targeted home visits by health and care workers.
 - **Community based preventive services**, such as targeted community-based fluoride varnish programmes and targeted provision of toothbrushes and tooth paste (i.e. postal or through health visitors).
 - **Supportive environments**, such as supervised tooth brushing, healthy food and drink policies, and fluoridation of public water supplies.
 - **Community action**, such as targeted peer support groups or school/community food co-operatives.
 - **Healthy public policy**, such as infant feeding policies.
17. Initiatives at a national and local level are already in place, addressing tooth decay from these different perspectives on the issue. A combination of interventions is required to help prevent tooth decay and improve oral health.
18. **Brushing teeth twice daily with fluoride toothpaste** also helps to prevent tooth decay. A 2013 national survey indicated that around 8 in 10 children aged 5 to 15 in England brushed their teeth twice or more a day according to parental reports⁶. Tooth brushing twice daily among 15 year olds eligible for free school meals (FSM) was lower than among their peers not eligible for FSM.⁷⁷ Expert opinion suggests that children need to be helped and supervised by an adult, when brushing, until at least 7 years of age. Supervised tooth brushing in early years settings is one way of improving oral health. However, it would not achieve the policy objective of improving oral health for both adults and children in the north east region. Furthermore, it requires behaviour change by the individual, which may be more achievable for some than others. Water fluoridation is a preventative policy which improves oral health without behaviour change and would be complementary to other locally commissioned interventions that increase fluoride use.
19. **Healthy food & drink schemes:** Sugar consumption is the main cause of tooth decay. Measures taken to reduce sugar consumption will have a positive effect on improving children's oral health. The case for change and strategic framework of the Major Conditions Strategy, published August 2023, included a commitment to continue to work with stakeholders and industry to reduce sugar, salt and calories in food, including for baby food and drink. This includes businesses continuing to work towards delivering the 20% sugar reduction target that make the biggest contributions to intake of sugar for children aged up to 18 years.
20. The most recent progress report showed that reductions in sugar were made in all food categories, with the highest reductions being in breakfast cereals (14.9%) and yogurt and fromage frais (13.5%) between 2015 and 2020. The overall reduction was 3.5% when weighted for the highest selling products. The difference between this figure and the reductions in cereals and yogurts occurs because of the increased sales of higher sugar products such as chocolate confectionary. Levels of sugar in drinks subject to the Soft Drinks Industry Levy (SDIL) reduced by 46% over the same period. Further data on juices and milk-based drinks, and drinks subject to the Soft Drinks Industry Levy (SDIL), will be published in 2024. Water fluoridation can mitigate the impact of sugar consumption and subsequent tooth decay.

21. **Facilitating access to dental services:** In July 2022, the government announced a package of reforms to improve access to NHS dentistry. This outlined steps to meet oral health need and increase access to dental care. The changes implemented include improvements to ensure dentists are remunerated more fairly for more complex work. In February 2024, the government and NHS England published a national plan for the recovery and reform of NHS dentistry services.
22. From 1 April 2023 responsibility for commissioning primary care dentistry to meet the needs of the local population has been delegated to all integrated care boards (ICBs) across England. ICBs are responsible for having local processes in place to involve patient groups, and for undertaking oral health needs assessments, to identify areas of need and determine the priorities for investment.
23. The NHS Long Term Workforce Plan, published on 30 June 2023, sets out the steps the National Health Service and its partners need to take to deliver an NHS workforce that meets the changing needs of the population over the next 15 years. These include a 40% increase to dentistry undergraduate training places by 2031/32.
24. Water fluoridation provides a population level intervention to reduce tooth decay that does not rely on accessing dental services.

Summary and preferred option with description of implementation plan

25. In summary, whilst alternative options to expanding water fluoridation exist which may improve oral health, these are considered complimentary rather than a substitute. Water fluoridation has unique qualities which will help address the policy objective, particularly for vulnerable groups. Other programmes and policies are already in place or could be in place via local authority commissioning structures and water fluoridation schemes would complement these initiatives. The north east region has been chosen as existing feasibility studies on water fluoridation have been carried out, the water company has experience of water fluoridation and there are large inequalities in oral health. Therefore, this impact assessment considers the costs, benefits and risks of expanding water fluoridation in the north east region, compared to a do nothing scenario. Costs and benefits of alternative options are not included.
26. Implementation would proceed following public consultation and assuming authority to proceed is in place. This would include development of necessary and appropriate fluoridation legal agreements and/or variations to existing agreements with Northumbrian Water Ltd. Development would include settling final planning, capital and operating requirements. This would involve transition to the current work on existing water fluoridation schemes and the Secretary of State's responsibilities for reimbursing water companies for the reasonable capital and revenue costs of these schemes.

Costs and benefits

Summary table of costs and benefits

		Do nothing	Expansion of water fluoridation
Costs	Capital costs	£0	£139m (social; £26m financial)
	Operational costs	£0	£1.1m per year (financial)
	Total costs (social, discounted)	£0	£294m
Benefits	Financial benefits (dental healthcare cost savings & productivity gains)	£0	£586m

Quality of life benefits to the individual	£0	£24m
Total benefits (social, discounted)	£0	£610m

Note: None of the above constitute direct or indirect costs to business. Costs are met by Secretary of State.

Overview of economic model

27. The Office for Health Improvement and Disparities (OHID) has developed an economic model (the “WF model”) to quantify the costs and benefits of the proposed expansion of the water fluoridation programme across the north east region of England. Further detail on the methodology is set out in Annex A: Modelling Approach. Here, we include a high-level overview of the modelling approach. Flow charts are included throughout this section; they are a high level representation of the model’s functionality.
28. Water fluoridation leads to reduced caries in both deciduous (baby) and permanent (adult) teeth. For each cohort (year group) the WF model first estimates the expected reduction in caries for individuals. This is then used to assess the magnitude of health benefit to the individual as well as cost savings, from a reduction in dental resource use (fillings and extractions) and lost workplace productivity. The cost savings are then compared to the expected cost of delivering the expansion of water fluoridation to understand its value for money over time.

Cohort modelling approach

29. The WF model uses a cohort-based modelling approach which follows fixed groups of individuals over time. Each year group forms its own cohort with a new cohort entering the model each year. The cohorts are composed of a group of representative or ‘average’ individuals in the north east region of England in terms of their cumulative decayed, missing or filled teeth (dmft/DMFT), which is the outcome metric used to measure carious teeth. This is discussed further in Annex A: Modelling Approach.
30. The model uses a 40 year time horizonⁱ. The Green Book⁸ indicates that infrastructure projects, like new buildings, should be appraised over a 60 year period, whereas refurbishment of existing buildings is considered over 30 years. Given the new, or changes to existing, infrastructure required to implement water fluoridation, a 40 year appraisal period has been selected to appraise both costs and benefits. Rather than estimating a cohort’s annual benefits until death, benefits and costs are capped at 40 years post introduction of the programme. This is considered a reasonable approach, as it negates the need to draw in assumptions pertaining to average cohort life expectancy which would have added an additional layer of uncertainty. Capping benefits at 40 years rather than modelling death rates ensures our value for money estimates remain a conservative reflection of the programme’s potential true value.
31. Due to the cohort modelling approach taken, the WF model only estimates benefits for individuals born into the cohorts once water fluoridation is already in place. The WF model does not account for benefits that might be accrued by individuals who are already living when water fluoridation is commenced, due to uncertainty around the evidence related to the benefits of partial exposure. The potential benefits for these individuals, such as children and adults who already live in the area before it is expanded, are not quantified. As a result, the WF model produces a potentially conservative estimate of monetised benefits and they could be larger if a larger group than the cohorts stand to benefit from expansion of water fluoridation.

ⁱ The 40 year time horizon dictates that, for example, the 40th cohort to enter the deciduous teeth model (born 40 years from now), will only receive 1 year of benefits.

Core assumptions

32. Assumptions underpinning the analysis of costs and benefits for the expansion of water fluoridation are discussed throughout this Impact Assessment and in detail in 'Annex A: Modelling Approach'; sources for assumptions are discussed in detail in the annex. Core assumptions to note include:

- No population growth within the north east regionⁱⁱ.
- Any benefits accrued by individuals moving into the area would be equally and oppositely offset by individuals moving out of the area.
- Some inputs to the model rely on data for England; we assume these estimates hold for the north east region, for example the effect size from fluoride on dmft.
- As noted above, the WF model bases analysis of benefits on reduced caries, measured through dmft/DMFT. We do not use historic trends in dmft/DMFT to predict future baseline dmft/DMFT values over our 40 year time horizon, rather the baseline dmft/DMFT values observed in the control population are fixed over time. This reflects the fact we do not know if average dmft/ DMFT values in the population will increase, decrease or stay the same over the next 40 years. Data from the oral health survey of 5 year old children indicates that levels of dental decay in the north east region remained at similar levels between 2017 and 2022²⁷.
- In line with the Green Book⁸ a 3.5% discount rate has been applied to our financial costs and benefits, with a 3.0% rate used after year 30. A 1.5% discount rate is applied to health impacts, with a 1.29% rate used after year 30.

Evidence

33. As well as published systematic reviews, this impact assessment and the WF model have made use of evidence from statutory health monitoring reports. There is a legal obligation to monitor health impacts of water fluoridation schemes every four years. The health monitoring report for England 2022 looked at three dental health outcomes: experience and severity of tooth decay in 5 year olds and hospital admissions for removing teeth due to decay in children and young people.

34. The model also uses evidence from National Dental Epidemiology Programme surveys, for example the oral health survey of 5 year old children takes place every 2 years to collect information of children who attend mainstream, state-funded schools across England.

35. Prior to consultation we convened external experts focussed on water fluoridation and economic modelling to test appropriate outcome measures and available data sources to model impacts for children and adults.

Costs

36. The costs associated with the delivery of the expansion of water fluoridation consist of:

- Capital costs such as the costs associated with the machinery and equipment needed to operate the fluoridation plant, and
- Operational costs such as the revenue costs associated with the day to day running and maintenance.

ⁱⁱ Holding population growth constant over the 40 year time horizon, appears appropriate in the context of the ONS population estimates for 0-4 year olds in the North East region. The ONS estimates that our population of interest will steadily decline from 2019 to 2030 (falling to approx. 90% of its baseline 2019 value) before reversing this trend, steadily growing at a more gradual rate. By 2043 the last year of estimated ONS data the population has not returned to its 2019 baseline, but the trend remains shallowly upward. In light of these estimates and the paucity of data for the remaining 20 years, 'no population grow' was adjudged a responsible simplifying assumption.

37. Neither capital nor operational costs represent direct costs to business. Before proposing a new fluoridation arrangement, the Secretary of State is required to consult the relevant water company on whether this would be “operable and efficient” (s87(11) of the Water Industry Act 1991). We have consulted Northumbrian Water Ltd (NWL) that the proposed scheme would be operable and efficient. The Secretary of State is also responsible for reimbursing water companies for reasonable capital and operating costs associated with water fluoridation schemes.
38. Recent inflationary pressures have been factored into the cost estimates, as NWL provided cost estimates for 2023 which would have accounted for inflationary elements specific to their cost base between 2019-2023. We have used the GDP deflator to put this into 2019 prices.
39. In the ‘do nothing’ scenario, there will be no new costs incurred as water provision will continue as normal, without fluoridation being expanded to new areas. For the expansion scenario, the estimated costs incurred by DHSC when expanding water fluoridation to more areas in the north east region are discussed below.

Capital costs

40. Capital costs represent the plants and associated equipment required to implement the proposed scheme. These estimates have been developed in consultation with Northumbrian Water Ltd during feasibility work. These set up costs would be incurred in years one and 20 of the appraisal period as it is assumed that equipment would need to be replaced after 20 years.
41. The capital cost for the lifetime of the expansion is estimated to be approximately £34 million (undiscounted).
42. In line with Green Book supplementary guidance recommendations to address optimism bias in the estimate of costs, we have uplifted the capital costs by a factor of 1.1 to account for any potential bias in the estimate. This is a conservative measure that helps to ensure we do not underestimate the capital costs and thus overstate the value of the programme. To account for potential bias in the estimates we have applied the 1.1 cost multiplierⁱⁱⁱ.

Operational costs

43. Operating costs have also been identified through consultation with Northumbrian Water Ltd. These operating costs are associated with the maintenance and administration of the water treatment plant on a day-to-day basis, such as payroll, overhead costs, as well as chemicals and maintenance expenses. Operational costs are estimated to be approximately £1.1m per year.

Social value of the cost estimates

44. Total discounted costs of the programme are estimated to be £50m over 40 years of the programme. In line with the Department impact assessment guidance the cost impact is then converted to QALYs at a rate of £15,000 per QALY, the QALYs are then discounted before they are remonetised at the social value of a QALY; £70,000 resulting in present value total cost estimate of £294m, from the social perspective.

Total discounted costs

45. In summary, total financial and social cost estimates over the 40 years of the programme are estimated to be £50m and £294m respectively.

Uncertainties associated with costs

ⁱⁱⁱ Supporting public service transformation: cost benefit analysis guidance for local partnerships - GOV.UK (www.gov.uk)

46. Cost estimates have been produced by Northumbrian Water Limited using their cost estimation system using actual project data. Given potential uncertainty with the costs, we also consider the NPV where financial costs are 30% higher (£327m social, £56m financial) and 30% lower (£261m social, £44m financial); the chosen higher and lower range (30%) has been provided by NWL.

Benefits

47. The World Health Organization recommends a maximum level of 1.5 milligrams per litre of water (mg/l)⁹. This is also the regulatory upper limit for fluoride set out in the Water Supply (Water Quality) regulations for England (monitored by the Drinking Water Inspectorate) and applies both to fluoridation schemes and fluoride naturally present in water. In England water fluoridation schemes aim for a concentration of 1 mg per litre as strong scientific evidence has shown this to be the optimal level of fluoride to reduce tooth decay, in both deciduous and adult teeth. In England, the water supply (water quality) regulations for England¹⁰ applies. Evidence from multiple sources¹¹ suggest that the addition of fluoride (1mg of fluoride per litre of water (mg/L)) to drinking water reduces caries in both deciduous and permanent teeth. A reduction in caries because of water fluoridation expansion can be expected to lead to several types of benefit, cost savings to society and quality of life benefits to the individual and their family, dental healthcare cost savings and productivity gains. These and an overview of the WF model methodology are discussed below. Further detail on the methodology is included Annex A: Modelling Approach.

48. The benefits set out below occur because of a reduction in caries; the outcomes used to measure dental caries are 'decayed missing or filled teeth' (dmft for deciduous teeth and DMFT for permanent teeth). The benefits are based on the observed difference in dmft/DMFT values in populations exposed to fluoridated water (treatment population) relative to a characteristically similar population not exposed to fluoridated water (control population); monitoring data²³ in England indicates that water fluoridation is associated with an observed ~20% reduction in caries at the age of 5¹².

49. To calculate the difference in average cumulative dmft/DMFT values for the control and treatment population, observed values are used where data are available^{iv} and linear approximation is used between ages where appropriate; for ages 12 onward for the treatment population, observed values are not available and instead the preventative fraction (27% reduction in DMFTs due to water fluoridation^v) is applied, based on Griffin et al¹³. More information about dmft/DMFT is found in Annex A: Modelling Approach.

Dental healthcare cost savings

50. The expansion of water fluoridation is expected to result in dental healthcare cost savings associated with a reduced need for tooth extractions and fillings. Avoided dmft/DMFT drive dental healthcare cost savings because these need to be extracted, filled or in the case of deciduous teeth, natural loss of teeth. Both extractions and fillings represent a direct cost to the health care system, as they require dental time and resource to treat.

51. The methodology for estimating potential cost savings from avoided dental healthcare is set out in Figure 3 and Figure 4 for deciduous and permanent teeth respectively. For deciduous teeth, we calculate the proportion of teeth with decay that are extracted, filled, or left untreated for 0 to 5 year olds. All extractions for 0 to 5 year olds are assumed to occur in secondary care (treatment in a hospital setting), and all fillings occur in primary care (dentist and community dental services), so from there we can estimate the avoided cost of extractions and fillings. For teeth with decay that are left untreated some will be fall out

^{iv} From the Water Fluoridation Health Monitoring Report for England (2014) and the Adult Dental Health Survey (2009)

^v Sensitivity analysis has been conducted, in order to understand what the minimum effect size would need to be for the programme to breakeven at the expiration of the 40 year time horizon

(exfoliate) naturally and therefore bear no dental healthcare costs. Some will not exfoliate naturally over time and will require extraction between the age of 6 and 10; for these procedures, we estimate the costs of secondary and primary care. Taken together this gives the total possible avoided extraction and filling cost for deciduous teeth.

52. For cost savings associated with avoided decay in permanent teeth, we estimate the proportion of decayed teeth that are extracted and that are filled; the associated costs of treating these health issues, either through primary or secondary care, are calculated and summed to produce a total avoided dental healthcare cost for decay to permanent teeth (see Figure 4).
53. Finally, the cost of the additional fillings and extractions under non fluoridation conditions, will inevitably displace dental health services that would have otherwise been provided to patients; this is the opportunity cost of the proposal. Following current DH guidance, the opportunity cost is calculated at one Quality Adjusted Life Years (QALY) per £15,000. The stream of QALYs foregone is then discounted at a rate of 1.5% per year (1.29% after 30 years). The displaced QALYs are then re-monetised at the value of £70,000 a QALY; representing the social value of a QALY.




Figure 3. Methodology for cost savings associated with avoided decay in deciduous teeth (per cohort, receiving full 10 years of benefit)




Figure 4. Methodology for cost savings associated with avoided decay in permanent teeth for the first cohort

Productivity gains

54. There may be productivity gains from a decrease in lost time at work due to a reduced need for fillings/extractions or caring for children requiring fillings/extractions.
55. We calculate the potential productivity gains for deciduous teeth and permanent teeth separately. For both, we first estimate the proportion of decayed teeth that are extracted or filled. For extractions we calculate the assumed school or work absence due to the extraction itself and assume that some school or work will also be missed while waiting for treatment. For fillings, we also estimate some school or work absence due to the filling procedure. With a total number of school or work days lost due to extractions and fillings, we are then able to estimate the total value of lost work days by multiplying for the average daily wage in the north east region. For deciduous teeth, our modelling factors in the proportion of carers who are employed and the impact on their working days; for permanent teeth, our modelling factors in the proportion of the adult population who are employed.




Figure 5. Methodology for productivity gains associated with avoided decay in deciduous teeth

56. Taken together, the total benefit accrued from dental healthcare cost savings and productivity gains of avoided decay to deciduous and permanent teeth is £586m (social perspective) over 40 years of the programme.

Quality of life benefits to the individual

57. A direct health benefit may occur for the individual, in the form of improved quality of life (QoL) due to the avoidance of caries, measured through dmft/DMFT, they would not have otherwise experienced in the absence of water fluoridation. Tooth decay can cause problems with eating, sleeping, communication and socialising.¹⁴

58. The methodology to calculate the direct health benefits that might arise from water fluoridation is set out in Figure 6. It follows the same approach to that set out in the Global burden of disease study which calculates disability adjusted life years associated with tooth decay³².

59. To estimate these benefits, avoided decay in deciduous and permanent teeth is considered separately, but the overarching methodology as set out in Figure 6 is generalised. The total number of caries that could be avoided are calculated (as measured through dmft/DMFT) we then estimate the number of avoided caries which would have resulted in mild vs severe pain.
60. These are then multiplied by the Disability Adjusted Life Years loss for mild and severe pain respectively. Disability Adjusted Life Years averted are calculated by multiplying the proportion of the year spent with symptomatic caries by the disability weight of systematic caries estimated from the Global Burden of Disease study (GBD)³². For individuals suffering from mild symptoms, they are estimated to experience pain for one hour per day over the course of one year. For individuals suffering from severe symptoms, they are estimated to be in pain for ~55 days for permanent teeth and ~27 days for deciduous teeth (as evidenced in the GBD study); for the remainder of the year they are assumed to experience pain for one hour per day.
61. Disability Adjusted Life Years are converted to Quality Adjusted Life Years gained; this is then multiplied by the social value of a Quality Adjusted Life Years to give a total social cost saving for direct health impacts from avoided decay in deciduous and permanent teeth.




Figure 6. Methodology for calculating direct health benefits from avoided decay in deciduous and permanent teeth

62. Taken together, applying the Green Book £70,000 social value of a QALY to the estimated 343.16 QALYs gained (discounted) from the programme, results in an estimated total discounted cost saving of £24m associated with direct health impacts from avoided decay to deciduous and permanent teeth, over 40 years of the programme.

Non-monetised benefits

63. **Benefits beyond the age of 45:** The model observes a 40 year time horizon for cohorts beginning at birth with deciduous teeth and 5 years of age with permanent teeth. This means that benefits are cut off at the maximum age of 45 years. This cut off is arbitrary and represents a conservative approach to estimating the time horizon over which benefits

accrue. It is likely that, assuming water fluoridation continues beyond the 40 year appraisal period, it continues to have an effect on DMFT values beyond 45 years of age. However, the monetised benefit of this effect is not included in the model as it would require drawing into our analysis survival/death related data and assumptions. Extending the model's time horizon to include benefits until an estimated death date for each cohort would inflate the estimated benefit water fluoridation expansion would be expected to deliver but with the addition of increased uncertainty associated with our estimates.

64. **Cost of lost education:** The model estimates the number of lost nursery or school days due to fillings and extractions in children. This parameter value is used to calculate the number of workdays lost due to the caring responsibilities of the child's parent/guardian. The time lost at work has been monetised using the wage rate. However, we do not attempt to monetise the value of the lost days at school for children undergoing extractions or fillings. Although the model quantitatively captures the number of lost nursery/school days per cohort, we do not have a robust estimate of the monetary value of lost school days deemed appropriate to apply within our model. Broader evidence suggests that, generally, the higher the percentage of sessions missed across key stages at KS2 and KS4, the lower the level of attainment at the end of the key stage^{vi}.
65. **Out of pocket expenses:** The avoidance of travel cost associated with getting to and from dental appointments, as well as the cost of dental treatment for those who pay for it, represents a likely cost saving we have not attempted to include in our model. Water fluoridation is expected to result in avoided fillings and extractions, and by extension number of trips to an individual's dental practice. The reduced number of trips to the dental practice reduces the individual's out of pocket expense, such as those related to fuel for personal transportation, public transport costs or lost time due to time spend traveling to the appointment. It has not been possible to quantify these potential cost savings due to lack of necessary transportation data.
66. **Family health benefits:** The model focuses on the quality of life benefits to the individual. However, there may also be benefit from avoided tooth decay to their family including parents who may experience health impacts like lost sleep due to their child suffering from tooth ache.
67. **Dental practice capacity:** It is likely that the averted dmft/DMFT and the associated knock-on effect on reduced need for dental appointment would free up capacity within the dental practice. The freed-up capacity may result in reduced wait times, so we might assume that people being seen more quickly by their dentist would result in caries being picked up at an earlier, possibly reversible stage. Again, we have not attempted to monetise this effect because it would require a high degree of assumption to offset the absence of data.
68. **Benefits of partial exposure to water fluoridation:** The model assumes only children born into the proposed expansion area after the commencement will receive total benefits associated with water fluoridation. The model assumes a child born before the start of the programme will not receive any benefits associated with reduced caries in deciduous teeth, even if they are exposed to water fluoridation for a large proportion of their childhood. This reflects a conservative approach. The model also assumes that partial longitudinal exposure results in no benefits. These are highly conservative assumptions. In addition, as we assume permanent teeth erupt between the ages of 5 to 12, only children who turn 5 after the start of expansion will receive benefits to their permanent teeth. For example, a child who is 4 at the time of the commencement of expansion will begin to receive benefits

^{vi} [The link between absence and attainment at KS2 and KS4, Academic year 2018/19 – Explore education statistics – GOV.UK \(explore-education-statistics.service.gov.uk\)](https://www.gov.uk/explore-education-statistics)

accruing to their adult teeth when they turn 5, because their permanent teeth are fully exposed to water fluoridation from the point of eruption.

69. **Second order and third order health effects:** The reduction in fillings or extractions represent the first order health effect. Following an extraction or filling, in some instances, there can be second or third order health events which would also be avoided. These include:

- a small proportion of forgone extractions/filling which would have resulted in infection and further complications requiring follow up dental or medical treatment
- foregone need for root canals including the cost associated with crowning the tooth
- forgone need for dentures and bridges following an extraction.

We have not attempted to include these costs in the model, due to insufficient/quality of the data. Excluding these costs ensures that the model represents a highly conservative estimate of the true value.

70. **Dental treatment under general anaesthesia** presents a small but real risk of life-threatening complications for children and carries significant morbidity for children undergoing this procedure¹⁵.

Summary of benefits

71. Overall, all benefits (dental healthcare cost savings, productivity gains and quality of life benefits to the individuals) are estimated to amount to £610m (social perspective) over 40 years.

Summary of Analysis & Sensitivity Analysis

72. Assuming that capital costs are incurred fully in years one and 20 of the scheme, the social benefit cost ratios are set out in the table below. This suggests that the programme would breakeven after 19 years based on social benefit cost ratios, as noted in Figure 7.

Table 1. Social cost benefit ratios, with capital costs accounted for in year 1 and 20

Time horizon (post introduction of the water fluoridation expansion)	Social benefit cost ratios
10 years	0.20
20 years	1.07
30 years	1.45
40 years	2.07




Figure 7. Accrual of costs and benefits over time

73. As a form of sensitivity analysis, we wanted to understand how sensitive our results were to changes in the size of the effect of water fluoridation on adult teeth. In our analysis we use the parameter value of 27% for the reduction in DMFTs due to fluoridated water (Griffin et al.¹³). The WF model estimates that the minimum effect size needed in permanent teeth for the water fluoridation programme to exactly breakeven by the end of the 40 year period was 12.1%.

Risks

Fluorosis

74. There is a well-established adverse association between levels of fluoride in water and the prevalence of dental fluorosis. Dental fluorosis is one of several different conditions that can affect the appearance of teeth. In England it is usually seen as paper white flecks or fine white lines, but it can vary in appearance from barely visible white lines to patches which may be of aesthetic concern. The risk period for the development of dental fluorosis in permanent (adult) teeth is when the teeth are growing in the jaws. Dental fluorosis cannot develop after teeth are formed. The first 2 to 3 years of life are generally accepted to be the period of highest susceptibility for fluorosis affecting the front teeth. Water fluoridation schemes in England aim to achieve a level of 1mg of fluoride per litre of water, equivalent to one part per million of water. This level was chosen to reduce tooth decay and reduces the risk of severe dental fluorosis.

75. The impact of milder forms of fluorosis on measured quality of life (using the Oral Health Related Quality of Life scale) is certainly less than that of tooth decay and may be non-existent or even positive.¹⁶¹⁷ A positive effect on quality of life may seem counterintuitive but may be explained by the fact that the white flecking of enamel associated with very mild fluorosis can give the impression of having teeth that are whiter than average¹⁸. More severe dental fluorosis can cause brown staining and pitting of teeth but is generally seen in those countries with very high naturally occurring levels of fluoride in groundwater rather than in areas with community water fluoridation schemes¹⁹. Dental fluorosis can also occur in the absence of water fluoridation, through ingestion of other sources of fluoride during tooth formation, particularly toothpaste and other fluoride supplements.

76. A study reporting on fluorosis prevalence and severity in England was commissioned to inform the 2018 health monitoring report. The population under examination was drawn from 4 cities; Newcastle upon Tyne and Birmingham, which are fluoridated, and Liverpool and Manchester that are not fluoridated)²⁰. The prevalence of any positive score for fluorosis was greater in Newcastle and Birmingham compared to Manchester and Liverpool. Fluorosis of a severity that might be of aesthetic concern affected 10.3% of children from the fluoridated cities and 2.2% in the non-fluoridated cities. There was, however, no difference in satisfaction with dental appearance between the 2 groups. Further research in adults in England has suggested that the aesthetic impact of fluorosis diminishes with age²¹.

General health risks

77. The Water Fluoridation Health monitoring report for England 2022 supports earlier findings and wider evidence that water fluoridation, at levels recommended in the UK, is a safe and effective public health measure to reduce dental caries and inequalities in dental health. The Water Fluoridation Health Monitoring Working Group continues to review evidence and will publish a further report in 2026.

Specific Impact Tests

Impact on small and micro businesses

78. Northumbrian Water Limited is the main business affected by the proposal to expanding water fluoridation in the north east region. They are considered a large business based on the number of employees. Costs for implementing the expansion would be incurred by government and not by the business. Therefore, we do not anticipate disproportionate burdens on small and micro business because of the proposals.

Equalities

79. A Public Sector Equalities Duty (PSED) assessment has been published separately alongside this as part of the consultation package. Adding fluoride to the water is a population level intervention that reaches everyone who lives there and uses mains water irrespective of their protected characteristics. Like many other non-communicable diseases there are significant inequalities related to deprivation in the levels of tooth decay around the country. Oral health inequalities were investigated in 2021 and for most of the protected characteristics there was no available evidence, or inconsistent evidence, on associations between oral health, care services and the protected characteristic. The PSED assesses a positive likely effect overall in the region with some narrowing inequalities on the basis of age, disability and race.

Rural proofing

80. Where the proposed scheme is operable, we do not anticipate a disproportionate impact of the policy on rural communities. Water fluoridation schemes will apply to both rural and urban areas. Some rural areas may not receive water fluoridation as it would not be operable and efficient to do so. These areas are clearly presented in the consultation package.

Competition assessment

81. We do not anticipate that the expansion of water fluoridation will impact competition. It should not directly or indirectly limit the number or range of suppliers, nor should it limit the ability of suppliers to compete or reduce suppliers' incentives to compete vigorously.

Carbon assessment and sustainable development specific impact test / environmental impact

82. We do not anticipate a disproportionate impact of the policy on rural communities; water fluoridation schemes will apply to both rural and urban areas.

Human rights

83. Expansion of water fluoridation is based on primary legislation passed by Parliament. We do not consider there to be any negative impact on human rights. The consultation seeks views on any missed evidence on ethical arguments.

Justice impact test

84. We do not anticipate an impact on the justice system from expanding water fluoridation in the north east region.

Potential trade implications of measure

85. We do not anticipate an impact on trade from expanding water fluoridation in the north east region.

Monitoring and Evaluation

86. The Secretary of State for Health and Social Care, has a legal duty to monitor the effects of water fluoridation schemes on health and report on it every 4 years. The next report is due in March 2026. This report is unlikely to report a significant change to population dental health if decision to expand water fluoridation in the north east region is taken, as there will be a long lead in time to fluoridation commencing. The current and previous water fluoridation health monitoring reports have taken a national view comparing areas with fluoride to those without fluoride in the water. Future health monitoring reports will consider the health effects of fluoride on people living in areas covered by water fluoridation schemes.

87. As part of the annual National Dental Epidemiology Programme, the Office for Health Improvement and Disparities coordinates dental surveys which describe levels of oral health in specific population cohorts. These annual surveys look at the oral health of five year old children biennially. This programme supports the collection, analysis and dissemination of reliable and robust information on the oral health needs of local populations. These surveys report the data at different geographical levels, from lower tier local authority level up to regional and national level.

88. The results of the biennial oral health surveys of five year old children in England inform the Public Health Outcomes Framework indicators²².

89. Any evaluation requirements needed in addition to the above monitoring processes will be considered during policy development and the consultation process.

Annex A: Modelling Approach

A1. Cohort approach

A1.1 Cohort specification

90. The WF model is formed of two overlapping cohort-based models, a deciduous teeth model and a permanent teeth model. The cohorts enter the deciduous teeth model at birth on an annual basis, exiting at 10 years of age or at the end of the model's 40 year time horizon, whichever comes first. Individuals enter the permanent teeth model at 5 years of age and

exit at the elapse of the 40 year time horizon. The overlapping nature of the cohort models dictate that between the ages of 5 to 10 years the individual is part of a cohort within both the deciduous and permanent teeth models.

91. The WF model assumes 0 to 5 year olds have only deciduous teeth, children between the ages of 5 and 10 are assumed to have a mix of deciduous and permanent teeth and after 10 years of age children have only permanent teeth. Consequently, water fluoridation benefits to deciduous teeth accrue from birth to 10 years. Benefits to permanent teeth accrue between the ages 5 up to a maximum of 45. Only the first cohort receives 40 year of benefit due to the time horizon cut off. This assumes that after the start of the programme children aged 5 to 10 exposed to water fluoridation receive benefits to both their deciduous teeth as well as to their emerging permanent teeth. There is no double counting of benefits between the ages of 5 and 10, because we use two entirely separate health outcomes to reflect the decay to permanent and deciduous teeth separately (see Figure 8 **Error! Reference source not found.**).




Figure 8. Permanent and deciduous model, and when different ages enter the model

A1.2 Cohort dynamics

92. Benefits associated with deciduous teeth are accrued from 0 to 10 years of age. The first cohort, those born the year expansion begins, will receive 10 years of benefits starting in year 1 to illustrate from 2023/24. Given the 40 year time horizon over which costs and benefits are observed the 31st cohort i.e. those individuals born in 2054/2055 receive benefits to their deciduous teeth from the age of 1 to 9 years and not in year 10, as year 10 would occur beyond the 'cut off' dictated by the 40 year time horizon.
93. For permanent teeth a similar approach is applied, however given the nature of deciduous and permanent dentation, benefits to permanent teeth do not begin until the individual turns 5 years old, unlike deciduous teeth. However, annual benefits are not capped by the emergence of a new set of teeth, this dictates that annual benefits accrue for each cohort until the expiration of the 40 year time horizon. As a worked example to illustrate, individuals turning 5 in 2023/24 when expansion begins, will receive annual benefits for 40 years (until they turn 45 years of age). The second permanent dentation cohort, those, aged 4 at the beginning of the programme, receive no annual benefit to their permanent teeth in year 1. Their annual benefit associated with permanent dentation begins in year two when they turn 5. Therefore they receive 39 years of annual benefit and so on until the 40th cohort who turn 5 years of age in the 40th year of the programme, receive just 1 year of benefit before the cut off imparted by the 40 year time horizon.

A2. DMFTs

94. Water fluoridation reduces dental caries. The outcomes used to measure dental caries are 'decayed missing or filled teeth' (DMFT) for permanent teeth and 'decayed missing or filled teeth' (dmft) for deciduous teeth.
95. DMFT is an index score ranging from 0-32 based on the case for permanent teeth which are decayed missing or filled. It is a cumulative measure with a value of 0, which means no teeth are decayed missing or filled and a value of 32 constituting all teeth being missing filled or decaying. A value of 7 would reflect the fact the individual had 7 teeth which are decayed, missing or filled.




Figure 9. Cumulative DMFT values (permanent teeth) for the first cohort of individuals

96. The benefits of water fluoridation are therefore based on the observed difference in dmft/DMFT values in populations exposed to fluoridated water (treatment population) relative to a characteristically similar population not exposed to fluoridated water (control population). For example, the Water Fluoridation Health Monitoring Report for England, 2018²³ shows that after controlling for other factors, 5 year olds in water fluoridated areas exhibit on average 0.81 dmfts per child relative to 1.01 in the population of 5 year olds not exposed to fluoridated water. This constitutes an observed ~20% reduction in caries at the age of 5 due to water fluoridation.
97. Figure 9 above shows the average cumulative number of DMFTs per individual over a 40-year time horizon from the age of 5-45 (permanent teeth). The purple line represents the average number of cumulative DMFT we would expect to see per individual in the north east region under non-fluoridated conditions (control) vs the green line, which is the average number of cumulative DMFT per individual following the introduction of water fluoridation (treatment). The average cumulative DMFT values for the control population (purple line) are observed at 12, 20, 30, 40 and 50 years of age^{vii}. To estimate the values for the intermediate ages, for example ages 13 through 19, we linearly approximate the values between the 2 time points.
98. To estimate the average cumulative DMFT value for the treatment population, which is represented by the green line, for ages 5 to 12 years we use the observed values from the Water Fluoridation Health Monitoring Report for England 2014²³. To estimate the average cumulative DMFT values for the treatment population from the age of 12 onward we apply

^{vii} DMFT rates at 40 years and 50 years were used to derive the estimated DMFT value for a 45 year old

the preventative fraction (% reduction in DMFT due to water fluoridation) from Griffin et al^{viii}, to the control population DMFT value. The cumulative DMFT values for the control population from the age of 12 onward are calculations based on data from the Adult Dental Health Survey 2009, England²⁴.

99. Figure 10 presents the control vs treatment cumulative dmft values for deciduous teeth (dmft) for 0 to 5 year olds. For dmft the same broad approach is applied however, instead of estimating the values for the treatment population by applying the preventative fraction to the control values, the Health Monitoring Report for England 2018²³ provides observed average values for both the treatment and control population groups, negating the need to estimate treatment dmft through application of a preventative fraction to control group values.




Figure 10. Deciduous teeth - cumulative dmft values for the first cohort of individuals (treatment vs control)

A3. Dental healthcare savings

100. Dental healthcare savings occur due to avoided caries, leading to avoided extractions and fillings for deciduous and permanent teeth. A flow chart outlining the methodology for calculating dental healthcare savings due to avoided caries can be seen in Figure 3 and Figure 4.

Deciduous teeth

101. **To estimate the cost of avoided extraction of deciduous teeth (0 to 5 years):**

- The number of avoided dmft per cohort per 6 months is multiplied by the proportion of teeth with decay experience that are extracted (10% is based on data from PHE's oral health survey of 5 year old children in 2014/15²⁵).
- This is then multiplied by the proportion of extractions occurring in secondary care (0 to 5 years) which is assumed to be 100%^{ix}; this gives us the number of avoided extractions occurring in a hospital settings per cohort per 6 months.

^{viii} [Effectiveness of fluoride in preventing caries in adults - PubMed \(nih.gov\)](#)

^{ix} Based on expert opinion

- This value is then multiplied by the cost of extraction in secondary care (£836, from the 2020/21 National Tariff Payment System²⁶) to give the avoided extraction cost for deciduous teeth, per cohort per 6 months from ages 0 to 5.
- As assumed that 100% of extractions for this age group occur in secondary care, we do not need to consider extractions at the dental practice for the 0 to 5 population.

102. **To estimate the cost of avoided extractions of deciduous teeth (6 to 10 years):**

- The number of avoided dmft per cohort per 6 months is used as a starting point. To calculate the number of these avoided dmft which go untreated by the age of 5, as we know that 10% of avoided dmft are extracted and 12% are filled, based on data from PHE's oral health survey of 5 year old children in 2014/15²⁷, this dictates that 78% of these avoided dmft go untreated from years 0 to 5. Multiplying the avoided dmft per cohort per 6 months by 78% gives us the number of avoided dmft going untreated per cohort per 6 months.
- To calculate what proportion of this untreated decay did not exfoliate naturally over time, we take the number of avoided dmft which go untreated and multiply by the probability of treatment before teeth exfoliated naturally (16 %, estimated through the PHE Oral Health Return on Investment tool²⁸) to give us the number of avoided dmft which went untreated yet did not exfoliate naturally and therefore now require extraction or filling between the years 6 to 10.
- This value is then multiplied by the probability that the tooth is extracted following no initial treatment or natural exfoliation (74%, estimated through the PHE Oral Health Return on Investment tool²⁸) to give the number of avoided cases of decay that would have gone untreated and resulted in an extraction.
- Half of these avoided extractions (50%, based on Kay et al., 2018²⁹) for the 6 to 10 year old population would have occurred in a hospital setting at a cost of £836, with the remaining 50 % taking place in the dental practice at a cost of £90.09^{xii}. Summing the avoided extraction cost in secondary care with the avoided extraction cost in the dental practice gives us the total avoided extraction cost for deciduous teeth in 6 to 10 year olds.

103. **The avoided filling cost for deciduous teeth** is calculated in a similar way to extractions, as outlined above. The only difference is that rather than a portion of fillings taking place in secondary care, 100% of fillings are carried out in the dental practice for all ages.

Permanent teeth

104. To calculate the cost of avoided extraction of permanent teeth:

- The number of avoided DMFT per cohort per 6 months is multiplied by the proportion of teeth with decay experience that are extracted (46% calculated using data from the Adult Dental Health Survey²⁴) and then multiplied again by the proportion of missing teeth which are missing due to dental extraction (90%^x). This gives us the number of avoided extractions in permanent teeth per cohort per 6 months.
- To estimate the cost of secondary care extractions, this value is multiplied by the proportion of extractions in secondary care (for adults; 10%^{xi}), and then multiplied again by the cost of extraction in secondary care (£836²⁶).

^x Expert opinion

^{xi} Expert opinion

- To estimate the cost of dental practice extractions, 90% are assumed to occur in the dental practice since 10% occur in a secondary setting. The number of extractions is multiplied by 90%, and then multiplied by the cost of extraction in primary care (£90.09^{xii}).

105. For the avoided filling cost for permanent teeth, the approach to modelling fillings mirrors the approach to calculating avoided extraction costs in permanent teeth, with the only difference being that 100% of fillings occur at the dental practice^{xiii}, at a cost of £90.09^{xii}; 0% of fillings occur in a hospital setting and therefore they do not need to be considered within this calculation.

A4. Productivity gains

106. Productivity gains occur due to avoided caries, leading to avoided missed school/work due to extractions. A flow chart outlining the methodology for calculating dental healthcare savings due to avoided caries can be seen in Figure 5.

Deciduous teeth

107. To calculate the total lost school days due to extractions in deciduous teeth:

- To calculate lost school days due to the extraction itself, the number of avoided extractions are multiplied by the assumed number school/work absence days due to an extraction' (3.5 days^{xiv}).
- To calculate the number of school/work days also lost for those experiencing pain in the build up to the extraction, the number of avoided extractions are multiplied by the proportion of individuals missing school/work due to dental issues during wait for tooth extraction (26%, based on data from Goodwin et al.³⁰), and then multiplied by 3.1 days, the estimated number of days absent from school/work due to pain, while waiting for treatment (Goodwin et al.³⁰).

108. To calculate the number of lost days at school due to fillings, the total number of avoided fillings are multiplied by 0.5 days, the assumed number of days absent from school/work due to a filling^{xv}.

109. To calculate the productivity gains related to caries in deciduous teeth:

- The total number of school/work days absent (due to the extraction itself, due to experiencing pain in the build up to the extraction, and due to fillings) is multiplied by the proportion of carers who are employed' (41%, Goodwin et al.³⁰) to arrive at a value for total number of lost days at work avoided.
- This value is then multiplied by the average daily wage (£124.60, based on the ONS Employment and Labour Market data for the north east region³¹) to give us the total value of lost workdays avoided in relation to deciduous teeth.

Permanent teeth

110. The calculation for lost working days avoided due to caries in permanent teeth follows the same method as for deciduous teeth with two key differences:

- We do not attempt to convert lost school days to lost workdays for individuals aged 12 to 18; the rationale here is that the 12 to 18 year old do not necessarily

^{xii} Expert opinion

^{xiii} Expert opinion

^{xiv} Expert opinion was elicited as part of the development of the PHE ROI tool, 4 days was judged appropriate for children. In order to be conservative, we have used an average value of 2.5 days for both children and adults, reflecting half a day for the procedure as well as two working/school days for recovery.

^{xv} Expert opinion

require someone at home to provide care, and consequently do not require a parent or guardian to take time of off work to look after them.

- Rather than multiplying the total lost school days avoided by the proportion of carers who are employed (41%), we multiply the avoided lost work days by the proportion of the adult population who are employed (75%, based on ONS Employment and Labour Market data for England³¹).

A5. Direct health benefit

Deciduous teeth

111. To calculate the direct health benefit accruing to deciduous teeth, we start with the total number of avoided caries (per cohort per 6 months).

To estimate the DALY loss associated with mild pain:

112. We take the averted caries value and multiply it by the proportion of decay resulting in mild pain (41%, calculated from the Global Burden of Disease (GBD) study³²). This calculation gives us the avoided caries which would have led to mild pain. DALY loss from a case of mild tooth ache for deciduous teeth (0.0004) is calculated by multiplying the proportion of the year spend in mild pain for deciduous teeth (0.04) by an estimate of symptomatic disability weight for dental caries (both calculated from the GBD study³²; 0.01). Avoided caries which would have led to mild pain is then multiplied by the DALY loss from a case of mild tooth ache for deciduous teeth (0.0004) to give us the DALY loss associated with mild pain.

To estimate the DALY loss associated with severe pain:

113. We take the avoided number of caries (per cohort per 6 months) and multiply it by the proportion of decay resulting in severe pain (19%; calculated from the GBD study³²). This gives us the avoided caries which would have led to severe pain. DALY loss from a case of severe tooth ache in deciduous teeth (0.001) is calculated by multiplying the proportion of a year spent in severe pain for deciduous teeth (0.11; calculated from the GBD study³²) is multiplied by an estimate of symptomatic disability weight for dental caries (calculated from the GBD study³²; 0.01). Avoided caries which would have led to severe pain is then multiplied by the DALY loss from a case of severe tooth ache for deciduous teeth (0.0011) to give us the DALY loss associated with severe pain.

114. With 41% of caries resulting in mild pain and 19% leading to severe pain, the remaining 40 % of caries are observed as asymptomatic. Given the asymptomatic nature of these caries, there is no health effect to be calculated from averting this caries. Consequently, the total DALYs averted per cohort per 6 months is derived from the aggregation of the DALYs associated with mild and severe pain (paragraphs 0 and 0).

115. The DALYs averted are converted to QALYs gained using the conversion factor set out in Sassi 2006³³ (1.37 for deciduous teeth and 0.66 for permanent teeth) which gives us the annual QALY gain per cohort per year. We then monetise this value by multiplying by the social value of a QALY (£70,000), before aggregating across all cohorts for each calendar year.

Permanent teeth

116. The method used to calculate the health benefit accruing to avoided caries in permanent teeth uses the same methodology as outlined above for deciduous teeth. However, it should be noted that two of the parameter values used to calculate the health benefit differ for permanent teeth relative to deciduous teeth:

- The DALY loss for a case of severe tooth ache is higher for permanent compared to deciduous teeth.

- The DALY to QALY conversion factor used for deciduous teeth (1.37) differs to the conversion factor used for permanent teeth (0.66)³³. This is because the factor is a function of age of disease onset; for deciduous teeth we have used the conversion factor associated with 5 year olds, for permanent teeth, we have used the conversion factor associated with 25 year olds.

-
- ¹ Office for Health Improvement & Disparities (2023). *Oral health survey of 5 year old children 2022*. Available at: <https://www.gov.uk/government/statistics/oral-health-survey-of-5-year-old-children-2022> [Accessed 25 September 2022].
- ² NHS Digital (2021). *Health Survey for England 2019: Supplementary analysis of dental health*. Available at: <https://digital.nhs.uk/data-and-information/publications/statistical/health-survey-for-england/2019/health-survey-for-england-2019-supplementary-analysis-on-dental-health> [Accessed 25 September 2023].
- ³ Office for Health Improvement & Disparities (2023). *Hospital tooth extractions in 0 to 19 year olds 2022*. Available from: <https://www.gov.uk/government/statistics/hospital-tooth-extractions-in-0-to-19-year-olds-2022/hospital-tooth-extractions-in-0-to-19-year-olds-2022> [Accessed 25 September 2023].
- ⁴ NHS Digital (2011). *Adult dental health survey 2009*. Available at: <https://digital.nhs.uk/data-and-information/publications/statistical/adult-dental-health-survey/adult-dental-health-survey-2009-summary-report-and-thematic-series> [Accessed 27 November 2023].
- ⁵ Public Health England (2014). *Local authorities improving oral health: commissioning better oral health for children and young people*. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/321503/CBOHMaindocumentJUNE2014.pdf [Accessed 25 September 2023].
- ⁶ Health & Social Care Information Centre (2013). *Children's Dental Health Survey 2013 – Country specific report: England*. Available from: <https://files.digital.nhs.uk/publicationimport/pub17xxx/pub17137/cdhs2013-england-report.pdf> [Accessed 25 September 2023].
- ⁷ Public Health England (2021). *Inequalities in oral health in England*. Available at: https://assets.publishing.service.gov.uk/media/6051f994d3bf7f0453f7b9a9/Inequalities_in_oral_health_in_England.pdf [Accessed 27 November 2023].
- ⁸ HM Treasury (2022). *The Green Book*. Available at: <https://www.gov.uk/government/publications/the-green-book-appraisal-and-evaluation-in-central-government/the-green-book-2020> [Accessed 7 September 2022].
- ⁹ World Health Organization (2017). *Guidelines for drinking-water quality, 4th edition, incorporating the 1st addendum*. Available at: [Guidelines for drinking-water quality, 4th edition, incorporating the 1st addendum \(who.int\)](https://www.who.int/publications/m/item/guidelines-for-drinking-water-quality-4th-edition-incorporating-the-1st-addendum) [Accessed 26 September].
- ¹⁰ The Water Supply (Water Quality) Regulations 2016 (2016). Available at: [The Water Supply \(Water Quality\) Regulations 2016 \(legislation.gov.uk\)](https://www.legislation.gov.uk/uksi/2016/1626/contents/made) [Accessed 26 September 2023].
- ¹¹ See for example [Water fluoridation: health monitoring report for England 2022 - GOV.UK \(www.gov.uk\)](https://www.gov.uk/government/publications/water-fluoridation-health-monitoring-report-for-england-2022); [Australian National Health and Medical Research Council 2017](https://www.nhmrc.gov.au/about-us/research-and-evidence/fluoridation) ; [Royal Society of New Zealand 2021](https://www.nzdr.co.nz/fluoridation); [Irish Health Research Board 2022](https://www.hse.ie/eng/fluoridation); [Canadian Agency for Drugs and Technologies in Health 2019](https://www.hc-sc.gc.ca/health/fluoridation); [Canadian Agency for Drugs and Technologies in Health 2019](https://www.hc-sc.gc.ca/health/fluoridation)
- ¹² Public Health England (2014). *Water fluoridation: health monitoring report for England 2014*. Available at: <https://www.gov.uk/government/publications/water-fluoridation-health-monitoring-report-for-england-2014> [Accessed 12 September 2023].
- ¹³ Griffin, O., Regnier, E., Griffin, P. M. & Huntley, V. (2007). Effectiveness of fluoride in preventing caries in adults. *J Dent Res*, 86 (5), pp. 410-415.
- ¹⁴ Office for Health Improvement & Disparities (2022). *Child oral health: Applying All Our Health*. Available at: <https://www.gov.uk/government/publications/child-oral-health-applying-all-our-health/child-oral-health-applying-all-our-health> [Accessed 7 September 2022].
- ¹⁵ Public Health England (2017) <https://www.gov.uk/government/publications/health-matters-child-dental-health/health-matters-child-dental-health> [accessed 12 October 2023]
- ¹⁶ Do, L. G. & Spencer, A. (2007). Oral health-related quality of life of children by dental caries and fluorosis experience. *J Public Health Dent*, 67 (3), pp. 132-139.
- ¹⁷ Chankanka, O., Levy, S. M., Warren, J. J. & Chalmers, J. M. (2010). A literature review of aesthetic perceptions of dental fluorosis and relationship with psychosocial aspects/oral health-related quality of life. *Community Dentistry and Oral Epidemiology*, 28 (2), pp. 97-109.
- ¹⁸ Hawley, G. M., Ellwood, R. P. & Davies, R. M. (1996). Dental caries, fluorosis and the cosmetic implications of different TF scores in 14-year-old adolescents. *Community Dental Health*, 13 (4), pp. 189-192.
- ¹⁹ Public Health England (2016) *Improving oral health: a community water fluoridation toolkit for local authorities* <https://www.gov.uk/government/publications/improving-oral-health-community-water-fluoridation-toolkit#:~:text=This%20publication%20was%20withdrawn%20on,toolkit%20is%20no%20longer%20current> [accessed 12 September 2023]

-
- ²⁰ Pretty, I. A., Boothman, N., Morris, J., MacKay, L., Lui, Z., McGrady, M. & Goodwin, M. (2016). Prevalence and severity of dental fluorosis in four English cities. *Community Dental Health*, 33 (4), pp. 292-296.
- ²¹ Macey, R., Tickle, M., MacKay, L., McGrady, M., & Pretty, I. A. (2018). A comparison of dental fluorosis in adult populations with and without lifetime exposure to water fluoridation. *Community Dentistry and Oral Epidemiology*, 46(6), 608-614.
- ²² Office for Health Improvement & Disparities (n.d.). *Public health outcomes framework*. Available at: <https://fingertips.phe.org.uk/profile/public-health-outcomes-framework/data#page/1/gid/1000044/pat/159/par/K02000001/ati/15/are/E92000001/yr/3/cid/4/tbm/1> [Accessed 3 October 2023].
- ²³ Public Health England (2014). *Water fluoridation: health monitoring report for England 2014*. Available at: <https://www.gov.uk/government/publications/water-fluoridation-health-monitoring-report-for-england-2014> [Accessed 12 September 2023].
- ²⁴ NHS Digital (2011). *Adult Dental Health Survey 2009*. Available at: <https://digital.nhs.uk/data-and-information/publications/statistical/adult-dental-health-survey> [Accessed 12 September 2023].
- ²⁵ Public Health England (2016). *Oral health survey of 5 year old children 2015*. Available at: <https://www.gov.uk/government/statistics/oral-health-survey-of-5-year-old-children-2014-to-2015> [Accessed 12 September 2023].
- ²⁶ NHS England (2021). *Nationl Tariff Payment System: national prices and prices for blended payments*. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/560972/Rapid_review_ROI_oral_health_5_year_old.pdf [Accessed 12 September 2023].
- ²⁷ Public Health England (2016). *Oral health survey of 5 year old children 2015*. Available at: <https://www.gov.uk/government/statistics/oral-health-survey-of-5-year-old-children-2014-to-2015> [Accessed 12 September 2023].
- ²⁸ Public Health England (2016). *Improving the oral health of children: cost effective commissioning*. Available at: [Improving the oral health of children: cost effective commissioning - GOV.UK \(www.gov.uk\)](https://www.gov.uk/government/publications/improving-the-oral-health-of-children-cost-effective-commissioning) [Accessed 12 September 2023].
- ²⁹ Kay, E., Owen, L., Taylor, M., Claxton, L. & Sheppard, L. (2018). The use of cost-utility analysis for the evaluation of caries prevention: an exploratory case study of two community-based public health interventions in a high-risk population in the UK. *Community Dent Health*, 35 (1), pp. 30-36.
- ³⁰ Goodwin, M., Sanders, C., Davies, G., Walsh, T. & Pretty, I. A. (2015). Issues arising following a referral and subsequent wait for extraction under general anaesthetic impact on children. *BMC Oral Health*, 15 (3). Available at: [Issues arising following a referral and subsequent wait for extraction under general anaesthetic: impact on children | BMC Oral Health | Full Text \(biomedcentral.com\)](https://doi.org/10.1186/s12902-015-0030-9) [Accessed 12 September 2023].
- ³¹ Office for National Statistics (n.d.). *Earnings and Working Hours*. Available at: [Earnings and working hours - Office for National Statistics \(ons.gov.uk\)](https://www.ons.gov.uk/economic-issues/income-wealth-and-living-conditions/earnings-and-working-hours) [Accessed 12 September 2023].
- ³² Kassebaum, N. J., Smith, A. G. C., Bernabé et al. (2017). Global, Regional, and National Prevalence, Incidence, and Disability-Adjusted Life Years for Oral Conditions for 195 Countries, 1990–2015: A Systematic Analysis for the Global Burden of Diseases, Injuries, and Risk Factors. *Journal of Dental Research*, 96 (4), pp. 380-387.
- ³³ Sassi, F. (2006). Calculating QALYs, comparing QALY and DALY calculations. *Health Policy and Planning*, 21 (5), pp. 402-408.