



UK Health
Security
Agency

**National influenza and
COVID-19 surveillance report**
Week 20 report
(up to week 19 2024 data)
16 May 2024

Contents

Contents.....	2
Executive summary.....	4
User feedback.....	6
Laboratory surveillance.....	7
Respiratory DataMart system (England).....	7
Confirmed COVID-19 cases (England).....	13
Microbiological surveillance.....	16
SARS-CoV-2 variants.....	16
Influenza virus characterisation.....	16
Influenza antiviral susceptibility.....	18
Community surveillance.....	19
SIREN healthcare cohort study.....	19
Winter COVID-19 Infection study.....	19
Acute respiratory infection incidents (ARI).....	19
FluSurvey.....	22
Google search queries.....	22
Flu Detector.....	23
Syndromic surveillance.....	24
Primary care surveillance.....	28
RCGP Clinical Indicators (England).....	28
RCGP sentinel swabbing scheme in England.....	29
Secondary care surveillance.....	33
Influenza, SARI Watch.....	33
COVID-19, SARI Watch.....	40
ECMO, SARI Watch.....	46
RSV admissions, SARI Watch.....	46
COVID-19 vaccination.....	47
COVID-19 vaccine uptake in England.....	47
Mortality surveillance.....	48

COVID-19 deaths	48
All-cause mortality assessment (England).....	48
International update	49
Global COVID-19 update.....	49
Global influenza update.....	49
Other respiratory viruses	49
Related links	50
About the UK Health Security Agency	51

For additional information including regional data on COVID-19 and other respiratory viruses, COVID-19 in educational settings, co- and secondary infections with COVID-19 and other data supplementary to this report, please refer to the [accompanying graph pack](#).

For additional information regarding data source please refer to [sources of surveillance data for influenza, COVID-19 and other respiratory viruses](#).

Executive summary

This report summarises the information from the surveillance systems which are used to monitor COVID-19 (caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)), influenza, and diseases caused by seasonal respiratory viruses in England. References to COVID-19 represent the disease name and SARS-CoV-2 represent the virus name. The report is based on data from week 19 of 2024 (between 6 May and 12 May 2024).

Please note that after the week 21 report (covering data up to week 20), this surveillance report will be moving to a condensed summer report and will then be released every two weeks.

Overall

In week 19, influenza activity decreased slightly across most indicators. COVID-19 activity increased across some indicators, though hospitalisations remained stable. Respiratory syncytial virus (RSV) activity fluctuated at low interseasonal levels.

Influenza

Through Respiratory DataMart, influenza positivity decreased slightly to 2.0% in week 19 compared with 2.7% in the previous week.

Through primary care surveillance, consultations for influenza-like-illness (ILI) decreased to 1.7 per 100,000 in week 19 compared with 2.8 per 100,000 in the previous week and remained within the baseline activity level range. The influenza positivity in GP sentinel swabbing increased to 3.9% in week 18.

There were 4 confirmed influenza acute respiratory infection (ARI) incidents reported in week 18, a slight decrease compared with the previous week (5 ARI).

Overall, the hospitalisation rate for influenza decreased and was at low levels at 0.40 per 100,000 compared with 0.58 per 100,000 in the previous week. The latest rate was within the baseline range (less than 1.57 per 100,000). The overall ICU or HDU rate for influenza was low and stable at 0.02 per 100,000 compared with 0.02 in the previous week and was within the baseline impact range.

Emergency department (ED) attendances for ILI decreased.

This season we have seen a combination of influenza A(H1N1) and A(H3N2). There has also been a small amount of influenza B activity later in the season.

COVID-19

Through Respiratory DataMart, SARS-CoV-2 increased slightly to 9.0% compared with 8.2% in the previous week.

Pillar 1 COVID-19 positivity increased overall and within all age groups and most regions in week 19, although total weekly cases decreased compared with the previous week.

The number of reported SARS-CoV-2 confirmed ARI incidents in week 19 decreased slightly compared with the previous week. There were 12 SARS-CoV-2 confirmed ARI incidents reported in week 19 in England, a decrease from 20 in the previous week.

Overall, COVID-19 hospitalisations stabilised to 3.13 per 100,000 compared with 3.21 per 100,000 in the previous week. Hospitalisations were highest in the aged 85 years and over age group. COVID-19 ICU admissions remained low and stable at 0.08 per 100,000 in week 19.

Respiratory Syncytial Virus (RSV)

Through Respiratory DataMart, RSV positivity remained low at 0.3%, with the highest positivity in those aged under 5 years at 0.7%. ED attendances for acute bronchiolitis decreased nationally.

Other viruses

Adenovirus positivity remained low at 2.8%, with the highest positivity in those aged under 5 years at 7.5%. Human metapneumovirus (hMPV) positivity remained low at 2.1%, with the highest positivity in those aged under 5 years at 6.1%. Parainfluenza positivity decreased slightly to 4.3%, with the highest positivity in those aged under 5 years at 6.1%. Rhinovirus positivity decreased slightly to 12.2% overall, with the highest positivity in those aged between 5 and 14 years at 29.7%.

User feedback

As part of our ongoing commitment to continuous improvement, we are asking for feedback on the National weekly influenza and COVID-19 surveillance report through the [survey accessible below](#). The purpose of this survey is to deepen our understanding of how readers engage with the report, highlighting areas readers find valuable and pinpointing areas for enhancement. The insights obtained from this survey will play a pivotal role in shaping the direction of future report development. The survey will be open until the end of the weekly reporting season.

Scan this QR code using a mobile device:

[National weekly influenza and
COVID-19 surveillance report
2023/24 Feedback Survey](#)



Laboratory surveillance

Respiratory DataMart system (England)

In week 19, data is based on reporting from 12 out of the 16 sentinel laboratories.

In week 19, 4,046 respiratory specimens reported through the Respiratory DataMart System were tested for influenza. There were 81 positive samples for influenza; 28 influenza A(not subtyped), 21 influenza A(H3N2), 5 influenza A(H1N1)pdm09, and 27 influenza B (Figure 3). Overall, influenza positivity decreased slightly to 2.0% in week 19 compared with 2.7% in the previous week.

In week 19, 4,529 respiratory specimens reported through the Respiratory DataMart System were tested for SARS-CoV-2. There were 406 positive samples for SARS-CoV-2 with an overall positivity of 9.0%, which increased slightly compared with 8.2% in the previous week. The highest positivity was seen in adults aged over 65 years at 12.4%.

RSV positivity remained low at 0.3%, with the highest positivity in those aged under 5 years at 0.7%.

Adenovirus positivity remained low at 2.8%, with the highest positivity in those aged under 5 years at 7.5%.

Human metapneumovirus (hMPV) positivity remained low at 2.1%, with the highest positivity in those aged under 5 years at 6.1%.

Parainfluenza positivity decreased slightly to 4.3%, with the highest positivity in those aged under 5 years at 6.1%.

Rhinovirus positivity decreased slightly to 12.2% overall, with the highest positivity in those aged between 5 and 14 years at 29.7%.

DataMart data is provisional and subject to retrospective updates.

Figure 1a. Respiratory DataMart weekly positivity (%) for influenza, SARS-CoV-2, RSV and rhinovirus, England




Figure 1b. Respiratory DataMart weekly positivity (%) for adenovirus, hMPV and parainfluenza, England





Figure 2. Respiratory DataMart weekly positivity (%) for influenza by year, England [note 1]



[note 1] Data from seasons 2020 to 2021 and 2021 to 2022 has been removed as there was low activity throughout these seasons.

Figure 3. Respiratory DataMart samples positive for influenza by type and subtype, England




Figure 5. Respiratory DataMart weekly positivity (%) for SARS-CoV-2 by year, England




Figure 6. Respiratory DataMart weekly positivity (%) for SARS-CoV-2 by age, England




Figure 7. Respiratory DataMart weekly positivity (%) for RSV by year, England





Figure 8. Respiratory DataMart weekly positivity (%) for RSV by age, England



Confirmed COVID-19 cases (England)


As of 9 am on 14 May 2024, there were a total of 2,151 Pillar 1 cases in week 19, a 7.0% decrease from the previous week.

COVID-19 polymerase chain reaction (PCR) positivity for Pillar 1 increased in week 19, with a weekly mean positivity rate of 9.8% compared with 8.6% in the previous week.

Pillar 1 positivity rates were highest in those aged over 85 years at a weekly mean positivity rate of 15.7% (increased from 15.1% among those aged over 85 years in week 18). They were highest in East Midlands at a weekly mean positivity rate of 11.8% (this increased from 8.9% in East Midlands in week 18).


Data notes: Changes to testing policies over time may affect positivity rates and incidence rates and should be interpreted accordingly. COVID-19 case reporting in England uses an episode-based definition which includes possible reinfections, each infection episode is counted separately if there are at least 91 days between positive test results (PCR or rapid lateral flow device). Each infection episode begins with the earliest positive specimen date.

Figure 9. Confirmed COVID-19 episodes tested under Pillar 1, based on sample date with overall 7-day rolling average PCR positivity for Pillar 1 (%) [note 2]



Age


Figure 10. 7-day rolling average PCR positivity (%) of confirmed COVID-19 cases tested under Pillar 1 by age group [note 2]



[note 2] The highlighted line corresponds to the age group in the subplot title, grey lines correspond to all other age groups.

Geography


Figure 11. 7-day rolling average PCR positivity (%) of confirmed COVID-19 cases tested under Pillar 1 by UKHSA region [note 3]



[note 3] The highlighted line corresponds to the UKHSA region in the subplot title, grey lines correspond to all other regions.

Ethnicity

Figure 12. Weekly incidence of confirmed COVID-19 cases per 100,000 population by ethnicity (Pillar 1), England [note 4]



[note 4] The highlighted line corresponds to the ethnicity in the subplot title, grey lines correspond to all other ethnicities.

Microbiological surveillance

SARS-CoV-2 variants

Due to processing changes for this dataset, updates to this section of the report are currently temporarily paused from 8 April 2024. Further information on future publications will be provided when available.

Influenza virus characterisation

UKHSA characterises the properties of influenza viruses through one or more tests, including genome sequencing (genetic analysis) and haemagglutination inhibition (HI) assays (antigenic analysis). This data is used to compare how similar the currently circulating influenza viruses are to the strains included in seasonal influenza vaccines, and to monitor for changes in circulating influenza viruses. The interpretation of genetic and antigenic data sources is complex due to a number of factors, for example, not all viruses can be cultivated in sufficient quantity for antigenic characterisation, so that viruses with sequence information may not be able to be antigenically characterised as well. Occasionally, this can lead to a biased view of the properties of circulating viruses, as the viruses which can be recovered and analysed antigenically, may not be fully representative of majority variants, and genetic characterisation data does not always predict the antigenic characterisation.

As of week 19 of 2024, the UKHSA Respiratory Virus Unit (RVU) has genetically characterised 927 influenza A(H3N2) viruses, which had been detected this season (since week 40), with 925 of these belonging in genetic subclade 3C.2a1b.2a.2 in the 2a.3a.1 subgroup. 2 A(H3N2) viruses belonging to subgroup 2a.3a were characterised. The Northern Hemisphere 2023/24 influenza A(H3N2) vaccine strain (an A/Darwin/9/2021-like virus) also belongs in genetic subclade 3C.2a1b.2a.2.

In the same period, 1,151 influenza A(H1N1)pdm09 viruses have been characterised. Sequencing of the haemagglutinin (HA) gene shows that 992 belong in genetic subgroup 6B.1A.5a.2a and 159 in subgroup 6B.1A.5a.2a.1. The Northern Hemisphere 2023/24 influenza A(H1N1)pdm09 vaccine strain (an A/Victoria/4897/2022 (H1N1)pdm09-like virus) also belongs in genetic subclade 6B.1A.5a, within the 6B.1A.5a.2a.1 cluster.

Since week 40, 544 influenza B/Victoria lineage viruses have been genetically characterised belonging in clade V1A.3a.2. Of these, 510 viruses belonged in genetic subclade C.5 and one in C.2. The Northern Hemisphere 2023/24 influenza B/Victoria lineage vaccine strain (a B/Austria/1359417/2021-like virus) also belongs in this V1A.3a.2 clade.

Different lineages may dominate during the season, and a close watch will be kept on the proportion of different viruses circulating to assist with the evaluation of vaccine effectiveness.

The RVU has confirmed by genome sequencing the detection of live attenuated influenza vaccine (LAIV) viruses in 5 influenza A positive samples and in 8 influenza B positive samples. These have been collected since week 40, from children aged 2 to 16 years.

One influenza A(H1N2)v virus has been genetically characterised belonging in clade 1B.1.1. This is an unusual detection of a variant H1N2 (H1N2v) virus in a human clinical sample. The HA and NA genes as well as internal gene segments from the A(H1N2)v detection show a very close relationship to contemporary 1B.1.1 swine influenza A viruses from the UK.

Table 1. Number of influenza viruses characterised by genetic and antigenic analysis at the UKHSA Respiratory Virus Unit since week 40 of 2023

(Sub)type	Genetic group	Number sequenced
A(H3N2)	3C.2a1b.2a.2a.3a.1	925
	3C.2a1b.2a.2a.3	2
	Total	927
A(H1N1)pdm09	6B.1A.5a.2a	992
	6B.1A.5a.2a.1	159
	Total	1,151
B/Victoria-lineage	V1A3a.2 / C.5	510
	V1A3a.2 / C.2	1
	Total	511
A(H1N2)v	1B.1.1	1
	Total	1

Influenza antiviral susceptibility

Influenza positive samples are genome sequenced and screened for mutations in the virus neuraminidase (NA) and the cap-dependent endonuclease (PA) genes known to confer neuraminidase inhibitor or baloxavir resistance, respectively. The samples tested are routinely obtained for surveillance purposes, but diagnostic testing of patients suspected to be infected with antiviral-resistant virus is also performed.

Influenza virus sequences from samples collected between week 40 of 2023 and week 19 of 2024 have been analysed.

Analysis of 899 A(H3N2) viruses found no viruses with known markers of resistance to neuraminidase inhibitors. Analysis of 1,148 A(H1N1)pdm09 by sequencing found 6 oseltamivir resistant viruses taken from 5 patients:

- patient 1: 2 samples with a H275Y amino acid substitution. Immune compromised adult patient known to have received oseltamivir treatment
- patient 2: 1 sample with a H275Y amino acid substitution. Adult patient with a COPD exacerbation known to have received oseltamivir treatment
- patient 3: 1 sample with a H275Y amino acid substitution. Immune compromised adult patient known to have received oseltamivir treatment
- patient 4: 1 sample with a D199E amino acid substitution. Immune compromised adult patient known to have received oseltamivir treatment
- patient 5: 1 sample with a H275Y amino acid substitution. Adult patient referred for resistance testing, clinical details pending

Analysis of 503 influenza B NA sequences found no evidence of known markers of resistance to neuraminidase inhibitors. One influenza B virus with a I221T mutation, previously known to result in reduced susceptibility to Oseltamivir was detected.

No viruses with known markers of resistance to baloxavir marboxil were detected in 799 A(H3N2), 942 A(H1N1)pdm09 and 439 influenza B PA sequences analysed.

Table 2. Antiviral susceptibility of influenza positive samples tested at UKHSA-RVU

(Sub)type	Neuraminidase inhibitors: susceptible	Neuraminidase inhibitors: reduced susceptibility	Baloxavir: susceptible	Baloxavir: reduced susceptibility
A(H3N2)	899	0	799	0
A(H1N1)pdm09	1,142	6	942	0
B/Victoria-lineage	503	1	439	0

Community surveillance

SIREN healthcare cohort study

Testing in SIREN has been paused from 28 April 2024.

Winter COVID-19 Infection study

The Winter COVID-19 Infection Survey has now ended, with a final publication on 14 March 2024. Published reports can be accessed through the [Office for National Statistics \(ONS webpage\)](#) and [UKHSA webpage](#).

Acute respiratory infection incidents (ARI)

Here we present data on ARI incidents in different settings that are reported to UKHSA Health Protection Teams (HPTs).

There were 29 new ARI incidents reported in week 19 in England. In the latest week, these included:

- 21 incidents reported from care homes, of which 6 were laboratory confirmed for SARS-CoV-2, 2 for influenza A(not subtyped), 1 for influenza (no subtype information available), 1 for RSV, and 1 for hMPV
- 7 incidents reported from hospitals, of which 5 were laboratory confirmed for SARS-CoV-2, 1 for influenza A(not subtyped), and 1 for parainfluenza
- 1 incident reported from other settings, which was laboratory confirmed for SARS-CoV-2

Please note that data back to week 40 was retrospectively updated following an improvement in the method to assign incidents to an identified pathogen using reports from health protection teams.

Figure 13. Number of ARI incidents by setting, England




Figure 14. Number of ARI incidents in all settings by virus type, England




Figure 15. Number of ARI incidents in care homes by virus type, England





Figure 16. Number of ARI incidents in educational settings by virus type, England



FluSurvey


As of week 18 of 2024, community surveillance using FluSurvey has been paused and will resume during the 2024 to 2025 season. No further data for the 2023 to 2024 season will be included in this report.

Google search queries

This is a web-based syndromic surveillance system which uses daily search query frequency statistics obtained from the Google Health Trends API (Application Programming Interface). This model focuses on search queries about COVID-19 symptoms as well as generic queries about 'coronavirus' (for example 'COVID-19'). The search query frequency time series is weighted based on symptom frequency as reported in other data sources. Frequency of searches for symptoms is compared with a baseline calculated from historical daily data. [Further information on this model](#) is available online.

During week 19, the overall and media-debiasing weighted Google search scores increased compared with the previous week (Figure 17).

Figure 17. Normalised Google search score for COVID-19 symptoms, with weighted score for media-debiasing and historical trend, England




Flu Detector

Flu Detector is a web-based model which assesses internet-based search queries for influenza-like illness (ILI) in the general population.

Daily ILI rate estimates are based on uniformly averaged search query frequencies for a week-long period (including the current day and the 6 days before it).

For week 19, the daily ILI query rate remained stable and was below baseline activity (Figure 18).

Figure 18. Daily estimated ILI Google search query rates per 100,000 population, England




Syndromic surveillance

During week 19, NHS 111 calls and online assessments for acute respiratory infections (ARI) decreased slightly. GP in-hours consultations for influenza-like-illness remained stable. GP out-of-hours daily contacts for ARI and influenza-like-illness decreased slightly. ED attendances for ARI decreased, including in those aged between 5 and 14 years. ED attendances for COVID-19-like illness remained stable nationally. ED attendances for acute bronchiolitis and influenza-like-illness decreased.

Please note that recent updates to the NHS Pathways clinical tool used by NHS 111 have affected levels of certain syndromic indicators. As a result of these changes a selection of existing NHS 111 syndromic respiratory indicators have been removed from this report and replaced with generic 'acute respiratory infections' calls and assessments indicators.


For further information on syndromic surveillance please see the [Syndromic surveillance: weekly summaries](#).

Figure 19a. Daily ED attendances for acute respiratory infection nationally, England [note 5]




[note 5] The solid black line is a 7-day moving average adjusted for holidays. The solid green line is the daily attendances. The black dotted line is the baseline. The grey columns show weekends and bank holidays.

Figure 19b. Daily ED attendances for acute respiratory infection by age group, England [note 6]




[note 6] The scales may vary in each graph to enable trend comparison. The black line is the 7-day moving average adjusted for bank holidays.

Figure 20a. Daily ED attendances for influenza-like illness nationally, England [note 5]




See [note 5] as above.

Figure 20b. Daily ED attendances for influenza-like illness by age group, England [note 6]




See [note 6] as above.

Figure 21a. Daily ED attendances for acute bronchiolitis nationally, England [note 5]



See [note 5] as above.

Figure 21b. Daily ED attendances for acute bronchiolitis by age group, England [note 6]




See [note 6] as above.

Primary care surveillance

RCGP Clinical Indicators (England)

The weekly ILI consultation rate through the Royal College of General Practitioners (RCGP) surveillance decreased to 1.7 per 100,000 registered population in participating GP practices in week 19 compared with 2.8 per 100,000 in the previous week. This was within baseline activity levels (less than 10.25 per 100,000) (Figure 22). By age group, the highest rates were seen in those aged between 15 and 44 years (2.4 per 100,000), followed by those aged between 45 and 64 years (1.8 per 100,000). The lower respiratory tract infections (LRTI) consultation rate decreased to 83.1 per 100,000 in week 19 compared with 95.5 per 100,000 in the previous week. Due to technical issues with the data suppliers the number of practices included in this week’s report is lower than normal.

Figure 22. RCGP ILI consultation rates, all ages, England



Moving Epidemic Method (MEM) thresholds are based on data from the 2015 to 2016 season to the 2022 to 2023 season. Please note the 2020 to 2021 and 2021 to 2022 seasons have been removed due to low activity throughout these seasons.

RCGP sentinel swabbing scheme in England

Starting from week 51, testing for enterovirus and rhinovirus have been delayed.


Based on the date samples were taken, in week 18 of 2024 (week commencing 29 April 2024) 458 samples were taken through the GP sentinel swabbing scheme in England of which 43 samples tested positive (Figure 23). Among all positive samples, 18.6% were positive for other seasonal coronaviruses, 16.3% were positive for hMPV, 30.2% were positive for SARS-CoV-2, 4.7% were positive for adenovirus, 27.9% for influenza and 2.3% were positive for RSV (Figure 24). Note there is a very small number of samples with an untyped influenza A result; this result occurs when subtyping fails due to a low viral load from the specimen. Due to the number of samples which have not yet been categorised, data should be interpreted with caution when compared with previous weeks. There were no available results for week 19. The proportion of detections among all positive samples is not calculated when the number of samples with a result is fewer than 50.

Among all samples which had a known test result, in week 18, positivity for SARS-CoV-2 was 4.2%, positivity for influenza was 3.9%, and positivity for RSV was 0.3% (Figure 25). Due to the number of samples which have not yet been categorised, data should be interpreted with caution when compared with previous weeks.

In previous reports, Figures 23, 24 and 25 were produced based on the date samples were received in the reference laboratory. From 23 November 2023 (week 47 report) these figures have been updated to be based on the date samples were taken.


From 27 November 2023, swabbing was temporarily increased in the Yorkshire and Humber region in response to the [identification of a case of influenza A\(H1N2\)v](#). This may lead to an over-representation of the Yorkshire and Humber region.

Figure 23. Number of samples tested for SARS-CoV-2, influenza, and other respiratory viruses in England by week, GP sentinel swabbing [note 7]



[note 7] Unknown category corresponds to samples with no result yet.


Figure 24. Proportion of detections of SARS-CoV-2, influenza, and other respiratory viral strains among virologically positive respiratory surveillance samples in England by week, GP sentinel swabbing scheme [note 8] [note 9]



[note 8] From week 51 data contains a substantial reduction of test results for enterovirus and rhinovirus due to a delay in testing for these pathogens.

[note 9] Data from the most recent weeks are not shown on this graph due to reporting delays.

Figure 25. Weekly positivity (%) for COVID-19, influenza and RSV in England, GP sentinel swabbing [note 9]



See [\[note 9\]](#) as above.

Secondary care surveillance


Influenza, SARI Watch

Surveillance of influenza hospitalisations to all levels of care is based on data from a small sentinel network of acute NHS trusts in England. Surveillance of admissions to ICU or HDU for influenza is mandatory with data required from all acute NHS trusts in England. Please note that the SARI Watch rates for 2023 to 2024 use the latest trust catchment population. For consistency the rates have been updated back to October 2020. The population denominator reflects changes in trust reconfiguration, hospital admission activity and population estimates.

In week 19 (ending 12 May 2024), the overall weekly hospital admission rate for influenza decreased to 0.40 per 100,000 compared with 0.58 per 100,000 in the previous week. The rate in the latest week was within the baseline range (less than 1.57 per 100,000). There were 35 new hospital admissions for influenza (21 influenza A(not subtyped), 2 influenza A(H1N1)pdm09, 3 influenza A(H3N2), and 9 influenza B).

In week 19, the overall ICU or HDU rate for influenza was low and stable at 0.02 per 100,000 compared with 0.02 in the previous week. The rate in the latest week remained within the baseline impact range (less than 0.11 per 100,000). There were 6 new case reports of an ICU or HDU admission for influenza in week 19 (5 influenza A(not subtyped) and 1 influenza A(H1N1)pdm09).

Figure 26. Weekly overall influenza hospital admission rates per 100,000 trust catchment population with MEM thresholds, reported through SARI Watch sentinel surveillance, England




MEM thresholds are based on data from the 2015 to 2016 season to the 2022 to 2023 season.

Please note the 2020 to 2021 and 2021 to 2022 seasons have been removed due to low activity throughout these seasons.

Influenza hospital admission rate is based on 20 sentinel NHS trusts for week 19.


SARI Watch data is provisional and subject to retrospective updates.

Figure 27. Weekly influenza hospital admissions by influenza type, reported through SARI Watch sentinel surveillance, England [note 10]



[note 10] Number of influenza hospital admissions based on sentinel NHS trusts.


Figure 28. Weekly hospital admission rate by UKHSA region for new influenza reported through SARI Watch sentinel surveillance [note 3] [note 11]



[note 11] Rates in some regions may not include all influenza surveillance sentinel trust sites from week to week. This may lead to variation in regional representation hence caution is required in interpreting the weekly data by region.

See [note 3] as above.

Figure 29a. Weekly hospital admission rate by age group for new influenza reported through SARI Watch sentinel surveillance - fixed y-axis [note 2]



See [note 2] as above.

Figure 29b. Weekly hospital admission rate by age group for new influenza reported through SARI Watch sentinel surveillance - adjusted y-axis





Figure 30. Weekly overall influenza ICU or HDU admission rates per 100,000 trust catchment population with MEM thresholds, reported through SARI Watch mandatory surveillance, England



MEM thresholds are based on data from the 2015 to 2016 to the 2022 to 2023 seasons. Please note the 2020 to 2021 and 2021 to 2022 seasons have been removed due to low activity throughout these seasons.

Influenza ICU or HDU admission rate is based on 90 NHS trusts for week 19.

SARI Watch data is provisional and subject to retrospective updates.

Figure 31. Weekly influenza ICU or HDU admissions by influenza type, reported through SARI Watch mandatory surveillance, England






Figure 32. Weekly ICU or HDU admission rate by UKHSA region for new influenza, reported through SARI Watch mandatory surveillance [note 3]




See [note 3] as above.

Figure 33a. Weekly ICU or HDU admission rate by age group for new influenza cases, reported through SARI Watch mandatory surveillance - fixed y-axis [note 2]



See [note 2] as above.

Figure 33b. Weekly ICU or HDU admission rate by age group for new influenza cases, reported through SARI Watch mandatory surveillance - adjusted y-axis




COVID-19, SARI Watch

Surveillance of COVID-19 hospitalisations to all levels of care and surveillance of admissions to ICU or HDU for COVID-19 are both mandatory with data required from all acute NHS trusts in England. Please note that the SARI Watch rates for 2023 to 2024 use the latest trust catchment population. For consistency the rates have been updated back to October 2020.

In week 19 (ending 12 May 2024), the overall weekly hospital admission rate for COVID-19 stabilised to 3.13 per 100,000 compared with 3.21 per 100,000 in the previous week. By UKHSA region, the highest hospital admission rate for COVID-19 was observed in the West Midlands (increased to 4.22 per 100,000 from 3.75 per 100,000 in the previous week, with a mixed picture in the remaining regions). By age group, the highest hospital admission rate for confirmed COVID-19 continued to be in those aged 85 years and over and decreasing to 31.41 per 100,000 compared with 34.03 per 100,000 in week 18. There was also a decrease in those aged between 75 to 84 years. There was a mixed picture in the remaining age groups although the direction of changes were small.

In week 19 (ending 12 May 2024), the overall weekly ICU or HDU admission rate for COVID-19 was very low and remained stable at 0.08 per 100,000, similar to the previous week. Note that with very low rates in critical care, small random fluctuations may occur. Note that ICU or HDU admission rates may represent a lag from admission to hospital to an ICU or HDU ward. The ICU or HDU admission rate for COVID-19 by UKHSA centre or by age group is currently fluctuating at low levels due to low underlying numbers.


Figure 34a. Weekly overall COVID-19 hospital admission rates per 100,000 trust catchment population, reported through SARI Watch mandatory surveillance, England



COVID-19 hospital admission rate is based on 88 NHS trusts for week 19.


SARI Watch data is provisional and subject to retrospective updates.

Figure 34b. Weekly percentage of hospitalisations with COVID-19 as primary reason, reported through SARI Watch sentinel surveillance, England




Data on proportions of hospitalisations primarily due to COVID-19 is based on returns from a smaller number of participating trusts in sentinel surveillance and may not be representative of all acute NHS trusts.

Figure 35. Weekly hospital admission rate by UKHSA region for new COVID-19 positive cases, reported through SARI Watch mandatory surveillance [note 3]



See [note 3] as above.

Figure 36a. Weekly hospital admission rate by age group for new COVID-19 positive cases reported through SARI Watch mandatory surveillance - fixed y-axis [note 2]



See [note 2] as above.

Figure 36b. Weekly hospital admission rate by age group for new COVID-19 positive cases reported through SARI Watch mandatory surveillance - adjusted y-axis






Figure 37. Weekly overall COVID-19 ICU or HDU admission rates per 100,000 trust catchment population, reported through SARI Watch mandatory surveillance, England



COVID-19 ICU or HDU admission rate is based on 73 NHS trusts for week 19.


SARI Watch data is provisional and subject to retrospective updates.

Figure 38. Weekly ICU or HDU admission rate by UKHSA region for new COVID-19 positive cases reported through SARI Watch mandatory surveillance [note 3]




See [\[note 3\]](#) as above.

Figure 39a. Weekly ICU or HDU admission rate by age group for new COVID-19 positive cases reported through SARI Watch mandatory surveillance - fixed y-axis [note 2]



See [note 2] as above.

Figure 39b. Weekly ICU or HDU admission rate by age group for new COVID-19 positive cases reported through SARI Watch mandatory surveillance adjusted y-axis




ECMO, SARI Watch

There were 5 extra corporeal membrane oxygenation (ECMO) admissions reported in week 19 from the 7 Severe Respiratory Failure (SRF) centres in the UK.

One admission was due to a suspected ARI, one was an ARI due to *S. aureus*, one was an ARI due to *L. pneumophila*, and 2 were due to non-infection.

Please note that the other group includes other viral, bacterial or fungal ARI, suspected ARI, non-infection (such as asthma, primary cardiac and trauma) and sepsis of non-respiratory origin.

Figure 40. Laboratory confirmed ECMO admissions in adults (COVID-19, influenza and non-COVID-19 confirmed) to Severe Respiratory Failure centres in the UK



SARI Watch data is provisional and subject to retrospective updates.

RSV admissions, SARI Watch

Week 16 of 2024 was the last week of RSV SARI Watch surveillance for the season.

Retrospective updates for data up to week 16 will be included [in the supplementary datafile](#) in the next weeks.

COVID-19 vaccination

COVID-19 vaccine uptake in England

Cumulative vaccination data up to week 19 2024 (week ending 12 May 2024) was extracted from the Immunisation Information System (formally National Immunisation Management Service). Age is calculated as age on 30 September 2024. From 15 April 2024, data are extracted on a Monday with data capped to the previous Sunday.


Data is provisional and subject to change following further validation checks. Any changes to historic figures will be reflected in the most recent publication.

Spring 2024 campaign

The spring 2024 data reported below covers any dose administered from 15 April 2024 provided there are at least 20 days from any previous dose. Eligible groups for the spring campaign are defined in the COVID-19 healthcare guidance [Green Book](#).

By the end of week 19 2024, (week ending 12 May 2024) 42.0% (2,429,207 out of 5,784,781 of all people aged 75 and over who are living and resident in England had received a vaccine dose in the spring 2024 campaign (Figure 41).

Figure 41. Cumulative weekly COVID-19 vaccine uptake in those who are living and resident in England vaccinated with a Spring 2024 dose since 15 April 2024 by age group.



Mortality surveillance

COVID-19 deaths

For further information on COVID-19 related deaths in England please see the [COVID-19 dashboard for deaths](#).

All-cause mortality assessment (England)

For further information on all-cause mortality in England please see the [Excess mortality within England: post-pandemic method report](#) which uses ONS death registration data, the [all-cause mortality surveillance report](#) which uses the European mortality monitoring (EuroMOMO) model to identify weeks with higher than expected mortality and the [ONS all-cause excess mortality report](#).

International update

Global COVID-19 update

For further information on the global COVID-19 situation please see the [World Health Organization \(WHO\) COVID-19 situation reports](#).

Global influenza update

For further information on the global influenza situation please see the [World Health Organization \(WHO\) Influenza update](#).

Influenza in Europe

For further information on influenza in Europe please see the [European Respiratory Virus Surveillance Summary weekly update](#).

Influenza in North America

For further information on influenza in the United States of America please see the [Centre for Disease Control weekly influenza surveillance report](#). For further information on influenza in Canada please see the [Public Health Agency weekly influenza report](#).

Influenza in Australia

For further information on influenza in Australia, please see the [Australian Influenza Surveillance Report and Activity Updates](#).

Other respiratory viruses

Avian influenza and other zoonotic influenza

For further information, please see the [Latest WHO update on 28 March 2024](#) and the [Latest UKHSA avian influenza technical briefing 14 July 2023](#).

Middle East respiratory syndrome coronavirus (MERS-CoV)

For further information please see the [WHO Disease Outbreak News Reports](#) and the [WHO publishes monthly updates](#).

[Further information on management and guidance of possible cases](#) is available online. The latest highlights that risk of widespread transmission of MERS-CoV remains very low.

Related links

[Previous national COVID-19 reports](#)

[Previous weekly influenza reports](#)

[Annual influenza reports](#)

[COVID-19 vaccine surveillance reports](#)

[Previous COVID-19 vaccine surveillance reports](#)

[Public Health England \(PHE\) monitoring of the effectiveness of COVID-19 vaccination](#)

[Investigation of SARS-CoV-2 variants of concern: technical briefings](#)

[Sources of surveillance data for influenza, COVID-19 and other respiratory viruses](#)

[RCGP Virology Dashboard](#)

UKHSA has delegated authority, on behalf of the Secretary of State, to process Patient Confidential Data under Regulation 3 The Health Service (Control of Patient Information) Regulations 2002

Regulation 3 makes provision for the processing of patient information for the recognition, control and prevention of communicable disease and other risks to public health.

About the UK Health Security Agency

UKHSA is responsible for protecting every member of every community from the impact of infectious diseases, chemical, biological, radiological and nuclear incidents and other health threats. We provide intellectual, scientific and operational leadership at national and local level, as well as on the global stage, to make the nation health secure.

The [UK Health Security Agency](#) is an executive agency, sponsored by the [Department of Health and Social Care](#).

© Crown copyright 2024

Prepared by: Immunisation and Vaccine Preventable Diseases Division
For queries relating to this document, please contact: respdsr.enquiries@ukhsa.gov.uk

Published: 16 May 2024
Publication reference: GOV-16694



You may re-use this information (excluding logos) free of charge in any format or medium, under the terms of the Open Government Licence v3.0. To view this licence, visit [OGL](#). Where we have identified any third-party copyright information you will need to obtain permission from the copyright holders concerned.



UKHSA supports the UN
Sustainable Development Goals

