



Animal &
Plant Health
Agency

United Kingdom Variety List Trials: Trial Procedures for Official Examination of Value for Cultivation and Use (VCU) Harvest 2024

Sugar Beet

Appendices

May 2023

Contents

Changes	2
Appendix 1 – Approved Trial Organisers/ Operators for Sugar Beet	2
Appendix 2 – Seed Treatment Products for Use on VL Trials – to be advised.....	3
Appendix 3 – Seed Delivery Deadline Dates	4
Appendix 4 – VCU Growing Trial Operators and Trial Locations for Sugar Beet	5
Appendix 5 – Control Varieties for VCU Assessments for Sugar Beet	6
Appendix 6 – Fungicide Programme for Sugar Beet	7
Appendix 7 – Dates by which Records should be sent to the Trial Design and Data Handling Organiser	8
Appendix 8 – % Assessment Keys for Sugar Beet Diseases.....	9
Leaf diseases.....	9
Infection Disease Severity Description	9
Appendix 9 – Growth Stage Key of Beet.....	10
Appendix 10 – Control of Groundkeepers from ALS tolerant plots in subsequent years. ..	11

Changes

Appendix 1 – Table changed; Trial Inspection Organiser updated from TBC to NIAB

Appendix 7 – Changed “Organiser” to “Operator”

Appendix 1 – Approved Trial Organisers/ Operators for Sugar Beet

Activity	Organisers / Operators Responsible
Trials Organiser	BSPB
Trial Design and Data Handling Operator	NIAB
Growing Trial Operators	See Appendix 4
Seed Handling Operator	See Appendix 4
Seed Treatment Operator	Applicants
DUS Operator	NIAB
Trial inspection Operator	NIAB
Technical Validation Operator	NIAB
Harvest Trial Operator	See Appendix 4
Quality Testing Operator	See Appendix 4
Data Review and Standard Setting Operator	NIAB

Appendix 2 – Seed Treatment Products for Use on VL Trials – to be advised

The following procedure is carried out to pellet and treat sugar beet seed for trials

Seed shall be treated with the following PPPs at the registered rates applicable and in accordance with the manufacturer's label recommendation.

- a) **Tachigaren 70 WP** To be applied as a surface coating to the pellet at **20 g product per unit** / 100,000 seeds
 - i. hymexazol total 20g commercial product = **14g ai/unit**; FRAC 32
 - ii. The hymexazol used shall be MAPP no: 17977

- b) **Force ST** To be applied as a surface coating to the pellet at **50ml product per unit** / 100,000 seeds
 - i. 200g tefluthrin per litre = **10g ai/unit**; Group 3A IRAC
 - ii. The Force SB used shall be MAPP no: 19042

Appendix 3 – Seed Delivery Deadline Dates

VCU seed must be delivered to the Seed Handling Operator by: **1st February**

Appendix 4 – VCU Growing Trial Operators and Trial Locations for Sugar Beet

Growing Trial Operators/Seed Handling Operators

Growing Trial Operator	Seed Handling Operator (If not Trial Operator)	Location of Trial	Harvest Operator	Quality Test Operator
NIAB	Trial Operator	North Walsham	BBRO	British Sugar
BBRO	NIAB	Rougham	BBRO	British Sugar
SESVANDERHAVE	NIAB	Caythorpe	SESVANDERHAVE	SESVANDERHAVE
SESVANDERHAVE	NIAB	Dawsmere	SESVANDERHAVE	SESVANDERHAVE
SESVANDERHAVE	NIAB	Heacham	SESVANDERHAVE	SESVANDERHAVE
KWS	NIAB	Swaffham Prior	KWS	KWS

Appendix 5 – Control Varieties for VCU Assessments for Sugar Beet

Appendix 6 – Fungicide Programme for Sugar Beet

Fungicides should be applied at or before the level of infection in the most susceptible variety reaches 5% in the older leaves. The programme should follow best local practice and a second application if necessary.

Appendix 7 – Dates by which Records should be sent to the Trial Design and Data Handling Operator

Record	Latest date of receipt
Site data part 1 (including site sketch)	Within 5 days of drilling trial
Site data part 2 plus diary	Within 5 days of harvesting the trial
Harvest date	Within 2 days of harvest
Confirmation of Trial Layout	Within 5 days of drilling the trial
Plot records	Within 5 days of record being taken
Harvest records	Within 5 days of harvesting the trial

Appendix 8 – % Assessment Keys for Sugar Beet Diseases

Leaf diseases

1. Examine leaves in 3 areas of each plot
2. Include all necrosis and chlorosis attributable to disease to be assessed
3. Estimate % infection using the description below, interpolating values if necessary
4. Record the average % infection from the 3 areas

Infection Disease Severity Description

Code	Description of severity
0	No infection observed
0.1	Older leaves with a trace of infection, other leaves uninfected.
1	Older leaves with up to 10 % infection, other leaves largely uninfected.
5	Older leaves with up to 25 % infection, middle aged leaves with a trace of infection.
10	Older and middle-aged leaves with up to 25 % infection, young leaves largely uninfected.
25	Leaves of all ages appear 50 % infected 50 % green on average
50	Leaves of all ages appear more infected than green on average
75	Very little green tissues left.
100	No green tissue left

Appendix 9 – Growth Stage Key of Beet

Beet Meier et al., 1993

Phenological growth stages and BBCH-identification keys of beet (*Beta vulgaris* L. ssp. *vulgaris*)

Code Description

Principal growth stage 0: Germination

- 00 Dry seed
- 01 Beginning of imbibition: seeds begin to take up water
- 03 Seed imbibition complete (pellet cracked)
- 05 Radicle emerged from seed (pellet)
- 07 Shoot emerged from seed (pellet)
- 09 Emergence: shoot emerges through soil surface

Principal growth stage 1: Leaf development (youth stage)

- 10 First leaf visible (pinhead-size): cotyledons horizontally unfolded
- 11 First pair of leaves visible, not yet unfolded (pea-size)
- 12 2 leaves (first pair of leaves) unfolded
- 14 4 leaves (2nd pair of leaves) unfolded
- 15 5 leaves unfolded
- Stages continuous till . . .
- 19 9 and more leaves unfolded

Principal growth stage 3: Rosette growth (crop cover)

- 31 Beginning of crop cover: leaves cover 10% of ground
- 32 Leaves cover 20% of ground
- 33 Leaves cover 30% of ground
- 34 Leaves cover 40% of ground
- 35 Leaves cover 50% of ground
- 36 Leaves cover 60% of ground
- 37 Leaves cover 70% of ground
- 38 Leaves cover 80% of ground
- 39 Crop cover complete: leaves cover 90% of ground

Principal growth stage 4: Development of harvestable vegetative plant parts Beet root

- 49 Beet root has reached harvestable size

Appendix 10 – Control of Groundkeepers from ALS tolerant plots in subsequent years.

Groundkeepers from ALS tolerant plots need to be controlled in the following crop. Groundkeepers can be produced through either harvest losses or non-germinated seed. Whilst good gleaning at harvest will minimize the risk of groundkeepers in the following year the following control measures should be adopted. Measure 3 should be regarded as obligatory.

1. Ploughing post-harvest. This will bury and kill any groundkeepers that may have been present. Your host grower should be made aware of this.
2. Using a non-ALS herbicide. In the following crop a non-ALS herbicide that will kill any beet present including ALS-tolerant beets. Your host grower should be made aware of this.
3. Inspecting the trial field. This must be done in the following season for the presence of groundkeepers. If any ALS-tolerant groundkeepers are present they must be removed, normally by hand. The location of the ALS plots should be recorded and these specific areas can be targeted for inspection. A diary entry should be made of inspection date, findings and control measures taken.



© Crown copyright 2023

You may re-use this information (excluding logos) free of charge in any format or medium, under the terms of the Open Government Licence v.3. To view this licence visit www.nationalarchives.gov.uk/doc/open-government-licence/version/3/ or email PSI@nationalarchives.gsi.gov.uk

This publication is available at www.gov.uk/government/publications

Any enquiries regarding this publication should be sent to us at

webmaster@apha.gov.uk

www.gov.uk/apha

The Animal and Plant Health Agency (APHA) is an executive agency of the Department for Environment, Food & Rural Affairs, and also works on behalf of the Scottish Government and Welsh Government.