



Which pertussis-containing vaccine should be given?

Given to pregnant women in each pregnancy

ADACEL (Sanofi)

Low dose diphtheria, tetanus and acellular pertussis vaccine (Tdap)



Boostrix-IPV or REPEVAX should be given to pregnant women if they have a known severe latex allergy or where ADACEL is not available and to obtain it would result in a delay in vaccination.

Given to babies at 8, 12 and 16 weeks

Infanrix hexa (GSK) or Vaxelis (Sanofi) Hexavalent vaccine DTaP/IPV/Hib/HepB

Diphtheria, tetanus, acellular pertussis, polio, Haemophilus influenzae type b (Hib) and hepatitis B




Infanrix hexa and Vaxelis should also be offered to children under 10 years of age who have not completed their primary immunisation course.



Given as a pre-school booster vaccine at age 3 years 4 months

Boostrix-IPV (GSK)

Low dose diphtheria, tetanus, acellular pertussis and polio vaccine (dTaP/IPV)



REPEVAX (Sanofi)

Low dose diphtheria, tetanus, acellular pertussis and polio vaccine (Tdap/IPV)



Either Boostrix-IPV or REPEVAX can be given as the pre-school booster vaccine (routinely at 3 years and 4 months and to children up to 10 years of age who have not received it). ADACEL does not contain polio and should therefore not be offered as the pre-school booster vaccine. Pertussis-containing vaccine is not routinely offered after 10 years of age, other than to pregnant women and eligible healthcare workers.

