

Great Britain and Northern Ireland Variety Lists / UK Plant Breeders' Rights Technical Protocol for Official Examination of Distinctness, Uniformity and Stability (DUS)

Turnip

Brassica rapa L. var. rapa (L.) Thell.

November 2024

Changes

- Updated title from United Kingdom to Great Britain and Northern Ireland
- Updated Plant Breeders' Rights to UK Plant Breeders' Rights
- Updated species latin name
- Updated dates relevant dates in document to 2024
- Section A
 - o Section 3.3
 - Updated language
 - Section 3.5
 - Removed fax no
 - o Section 4
 - Updated language
 - Section 7.1
 - Updated UPOV TGP from 8/4 to 8/5 and updated date
 - Updated UPOV TGP from TG/37/10 to 11 and updated date
- Section B
 - Section 4.1
 - Updated language
 - Section 5.1
 - Updated Seed Gazette to Gov website
 - Updated language
 - Section 6.1
 - Updated language
 - Section 7.1
 - Updated seed numbers
 - Section 7.2
 - Updated seed numbers
- Section C
 - Section 5.6
 - Added (thinned to) to Plant spacing
 - Section 6.3
 - Updated language
 - Section 6.5
 - Updated language
 - Section 7.2
 - Updated language
- Section D
 - o Section 5
 - Added "C Special Test" to type of observation of characteristics to cover character 1 and 2
 - Section 5.2
 - Updated characteristics
 - Section 5.3
 - Updated characteristics
- Section E
 - Section 6.2
 - Updated language
- Section G
 - Section 2.1
 - Removed latin name for species
 - Section 6.9
 - Updated scoring table

Contents

| Section A – General Information | 1 |
|---|----|
| 1 Purpose | 1 |
| 2 Scope | 1 |
| 3 Responsibilities | 1 |
| 4 Non-Compliance with the Protocol | 2 |
| 5 Responsibility for GM Releases | 2 |
| 6 Procedures for GM Varieties | 2 |
| 7 Associated Documents | 3 |
| Section B – Application Requirements | 4 |
| 1 Purpose | 4 |
| 2 Scope | 4 |
| 3 Responsibilities | 4 |
| 4 Receipt of Applications | 4 |
| 5 Receipt of Seed | 4 |
| 6 Seed Quality Requirements | 5 |
| 7 Seed Quantity | 5 |
| 8 Labelling Requirements, Including Provisions for GM Varieties | 6 |
| Section C – Growing Test Procedures | 7 |
| 1 Purpose | 7 |
| 2 Scope | 7 |
| 3 Responsibilities | 7 |
| 4 Reference Varieties | 7 |
| 5 Design of Tests | 7 |
| 6 Records and Recording | 9 |
| 7 Communications with the Applicant | 10 |

| Section D – Summary of DUS Characteristics to be Assessed, Method of Assessment Standards Applied | |
|---|----|
| 1 Purpose | 11 |
| 2 Scope | 11 |
| 3 Responsibilities | 11 |
| 4 Organisation | 11 |
| 5 DUS Characteristics to be Assessed | 11 |
| 5.2 Turnip Characteristics Routinely Recorded in DUS Tests | 13 |
| 5.3 Previously Approved Characteristics Not Routinely Recorded in DUS Tests | 19 |
| Section E – Reference Seed Stock Maintenance and VCU Seed Stock Authentication Procedures, where applicable | |
| 1 Purpose | 21 |
| 2 Scope | 21 |
| 3 Responsibilities | 21 |
| 4 Procedures for Reference Seed Stock Maintenance | 21 |
| 5 Procedures for VCU Seed Stock Authentication | 22 |
| 6 Procedures for the Inclusion of New Common Knowledge Varieties into the Reference Collection | |
| 7 Release of Reference Samples for Authorised Purposes | 23 |
| Section F – Procedures for Assessment of New Additional DUS Characters | 24 |
| 1 Purpose | 24 |
| 2 Scope | 24 |
| 3 Responsibilities | 24 |
| 4 Reference Varieties | 24 |
| 5 Procedures | 24 |
| Section G – Procedures for DUS Decisions | 25 |
| 1 Purpose | 25 |

| | 2 Scope | . 25 |
|---|---|------|
| | 3 Responsibilities | . 25 |
| | 4 Reference Varieties | . 25 |
| | 5 Distinctness | . 25 |
| | 6 Uniformity | . 26 |
| | 7 Stability | . 28 |
| | 8 DUS Report and Variety Description | . 29 |
| Α | Appendix 1 – Reference Collection Varieties | .30 |
| | 1 Variety Listing and Plant Breeders Rights | .30 |

Section A – General Information

1 Purpose

1.1 This Protocol sets out the procedures for conducting tests and assessments in relation to official examinations of DUS, maintenance of reference stocks of varieties of Turnip entered for Variety List (VL) Trials and Plant Breeders' Rights (PBR).

2 Scope

- 2.1 These procedures apply to all varieties of Turnip (*Brassica rapa* L. Turnip). Special procedures and responsibilities for Genetically Modified (GM) varieties are set out in Sections A5 and A6.
- 2.2 Except where specified in this protocol or authorised by the Plant Breeders' Rights Office for the UK, Animal and Plant Health Agency (APHA); only Variety List candidates, Plant Breeders' Rights candidates, candidates for Foreign Authorities and the reference varieties may be incorporated in the DUS tests.

3 Responsibilities

- 3.1 The growing tests and assessments in this protocol are carried out under the responsibility of the Secretary of State for Environment, Food and Rural Affairs, Scottish Ministers, Welsh Ministers and the Minister for Agriculture, Environment and Rural Affairs in Northern Ireland (the National Authorities).
- 3.2 They are supervised, on behalf of the National Authorities, by officials of the Testing Authorities: APHA; the Scottish Government (SG); the Department of Agriculture, Environment and Rural Affairs (DAERA); and the Welsh Government (WG).
- 3.3 This protocol is authorised by the Plant Variety and Seeds Committee (PVSC). It cannot be amended without its approval. Requests and suggestions for amendment of the protocol should be put in writing to APHA Plant Varieties and Seeds, either directly or via the Test Centre.
- 3.4 The procedures are administered by:

Plant Variety Rights Office for the UK Animal and Plant Health Agency Eastbrook Shaftesbury Road Cambridge

CB2 8DR Email: pvs.helpdesk@apha.gov.uk

3.5 Test Centre

The DUS growing tests and assessments in this protocol are co-ordinated and carried out by:

Vegetable DUS Test Centre

SASA

Roddinglaw Road

Edinburgh Tel No 0131-244 8890

EH12 9FJ Email vegvarietytesting@sasa.gov.scot

3.6 The Test Centre is responsible for providing the appropriate facilities.

4 Non-Compliance with the Protocol

4.1 Where the protocol uses the word "must" for any action, then failure to carry out this action will result in non-compliance. Where non-compliance occurs or there are concerns regarding the validity of any data or tests this must be reported to APHA. Where this protocol uses the word "should" for any action this is the method to be followed unless there are clear reasons which can be justified by the Test Centre.

5 Responsibility for GM Releases

5.1 GM Release Consent Holders are responsible for GM releases. All parties involved in DUS work operating under a GM Release Consent must adhere to the instructions of the Release Consent where necessary, to comply with the relevant consent conditions. Where DUS protocol non-compliance occurs, this must be reported to the consent holder and the Test Centre who will notify APHA.

6 Procedures for GM Varieties

- 6.1 Applicants intending to enter GM candidates must consult APHA, well in advance of their application, about specific requirements under GM regulations.
- 6.2 The Test Centre must ensure that no test or trial sites are planted with GM candidates and/or varieties until APHA has given the specific clearances.

7 Associated Documents

7.1 The following documents are associated with this protocol

| Reference | Title |
|--------------------------------|--|
| UPOV TG/1/3 | General Introduction to the Examination of Distinctness, Uniformity and Stability and the Development of Harmonised Descriptions of New Varieties of Plants (19.04.2002). |
| UPOV TGP/8/5 | Trial Design and Techniques Used in the Examination of Distinctness, Uniformity and Stability (28.10.2022). |
| UPOV TGP/9/2 | Examining Distinctness (29.10.2015). |
| UPOV TGP/10/2 | Examining Uniformity (01.11.2019). |
| UPOV TG/37/11 | Guidelines for the Conduct of Tests for Distinctness, Uniformity and Stability, Turnip (<i>Brassica rapa</i> L. var. <i>rapa</i> L.) 23.09.2022. |
| UPOV TWC/26/14 | An Adjustment to the COYD Method When Varieties are Grouped Within the DUS Trial. 01.08.2008. |
| GB and NI Variety Lists | The Seeds (National Lists of Varieties) Regulations 2001 (as amended) and The Seeds (Variety Lists) Regulations (Northern Ireland) 2020. |
| Plant Varieties Act 1997 | Plant Breeders' Rights Regulations 1998 and Plant Varieties Act 1997 |
| Plant Breeders' Rights 2019 | The Plant Breeders' Rights (Amendment etc.) (EU Exit) Regulations 2019 as amended by The Animal Health, Invasive Alien Species, Plant Breeders' Rights and Seeds (Amendment etc.) (EU Exit) Regulations 2019 and The Plant Breeders' Rights (Amendment) (EU Exit) Regulations 2020 |

Section B – Application Requirements

1 Purpose

1.1 The purpose of this section is to identify the specific DUS requirements for Variety List and/or Plant Breeders' Rights applications, as appropriate.

2 Scope

2.1 These procedures apply to all applications.

3 Responsibilities

3.1 The applicants are responsible for ensuring that these procedures are complied with.

4 Receipt of Applications

- 4.1 The latest date for receipt of applications for Variety List and/or for Plant Breeders' Rights, is stated on the GOV website https://www.gov.uk/national-lists-of-agricultural-and-vegetable-crops. Applications received after these dates may be considered for inclusion in the current year's tests and trials on a case-by-case basis.
- 4.2 The procedures for the submission of Variety Listing and/or Plant Breeders' Rights applications, Technical Questionnaires (TQs) and for payment of administration fees can be obtained from APHA at the address shown in Section A or on the GOV.UK website https://www.gov.uk/national-lists-of-agricultural-and-vegetable-crops
 - 4.3 Applicants should note in the TQ, submitted with the application, any additional characteristics which may require examinations that are listed in the DUS characteristics section D, 5.2 or 5.3 (an additional fee may be required).

5 Receipt of Seed

5.1 The latest date for receipt of seed is stated on the GOV website and is set administratively by APHA. In the absence of exceptional circumstances, seed submissions received after this date will be refused. Instructions for the delivery of seed will be made available to applicants by APHA https://www.gov.uk/national-lists-of-agricultural-and-vegetable-crops.

6 Seed Quality Requirements

- 6.1 The seed must satisfy the certification requirements for Basic Seed as laid down in the seed marketing legislation of the National Authorities.
- 6.2 The seed must not be chemically treated. Seed treatment, if required, will be undertaken by the Test Centre. The chemicals applied and rates of application will be determined by the Test Centre.

7 Seed Quantity

7.1 First Test Cycle

3,000 seeds or 10,000 seeds

7.2 Second Test cycle

7,000 seeds if 3,000 seeds were provided in year 1 No

seed if 10,000 seeds were provided in year 1

7.3 Shortfall in Seed Quantities

Where insufficient seed is available in the first instance a further stock must be supplied in the following year which will be authenticated against the original submission. An additional charge may be applied. This must be agreed in advance with APHA and the test centre.

8 Labelling Requirements, Including Provisions for GM Varieties

8.1 Applicants must clearly label their seed, inside and outside the bag, with the following information:

- Applicant
- AFP number (if known)
- Breeder's Reference number or name
- · Quantity of seed

8.2 All packages of GM material must be labelled clearly as "GMO" or "Genetically Modified Organism".

Section C – Growing Test Procedures

1 Purpose

1.1 The purpose of this section is to provide details of the procedures used in the growing tests for DUS analysis.

2 Scope

2.1 These procedures apply to all varieties of Turnip.

3 Responsibilities

- 3.1 The Test Centre is responsible for conducting these procedures.
- 3.2 The Test Centre will be responsible for ensuring that no material supplied to them is used for any other purpose than the conduct of these procedures or the release of reference samples for authorised purposes. (See Section E7).

4 Reference Varieties

- 4.1 The principles governing the selection of reference varieties are set out in Appendix 1.
- 4.2 Seed of reference varieties will be supplied by the Test Centre.

5 Design of Tests

- 5.1 The DUS Test Centre is responsible for selecting a suitable site which should be on ground that has normally not had a Cruciferous crop in the previous five years but may be less where it has been determined the risk is negligible.
- 5.2 Crop husbandry should follow best practice for all operations and particularly as regards cultivation, drilling, fertiliser and spray application, use of irrigation and control of weeds, pests and diseases.
- 5.3 The minimum duration of tests should normally be two independent growing cycles. The National List and Seeds Committee (NLSC) must be informed of any proposed changes to the number of cycles.
- 5.4 From information given in the Technical Questionnaire the candidate variety may be grown in plots and compared with varieties which are in the same classification for the following characters:

UPOV characteristics that could be used for grouping:

- (a) Ploidy (characteristic 1)
- (b) Petiole: intensity of anthocyanin coloration (characteristic 2)
- (c) Leaf: number of lobes (characteristic 6)
- (d) Root: degree of swelling (characteristic 15)
- (e) Only varieties with Root: degree of swelling: medium or strong: Root: colour of skin above soil (characteristic 16)
- (f) Only varieties with Root: degree of swelling: medium or strong: Root: colour of skin below soil (characteristic 18)
- (g) Only varieties with Root: degree of swelling: medium or strong: Root: colour of flesh (characteristic 19)
- (h) Only varieties with Root: degree of swelling: medium or strong: Root: shape in longitudinal section (characteristic 22)
- 5.5 Varieties known to be clearly distinct from the candidate on any other discontinuous or continuous characteristic may be excluded from the trial. If this exclusion is based on a characteristic which is not listed in section D 5.2 approval by the NLSC must be sought. See section F for further information on additional characters.
- 5.6 The tests are carried out using a grouped design, with a plot of each candidate and close control variety present in each replicate as follows:

| | Variety type | |
|-------------------------------------|-----------------|-----------------|
| | Horticultural | Fodder |
| Number of replications | 2 | 2 |
| Number of rows per plot | 3 | 2 |
| Spacing between plot rows | 0.4m (approx.) | 0.8m (approx.) |
| Plot length | 10 m | 10 m |
| Number of plants per replicate | 200 | 130 |
| Hence, number of plants per variety | 400 | 260 |
| Plant spacing (thinned to) | 0.15m (approx.) | 0.15m (approx.) |

Groups are randomised and varieties are randomised within groups.

- 5.7 Seed is sown in the field between April and early June according to a plan produced by the Test Centre and thinned to a stand to achieve the plant number per plot as indicated in C 5.6. Varieties are coded by the Test Centre.
- 5.8 Any candidate with distinctness problems in the first test cycle may be grown side by side with their close controls in the second or third test cycles.
- 5.9 Recordings are taken on each trial approximately 6 8 weeks after sowing until harvest maturity stage. Characters recorded are listed in Section D.

6 Records and Recording

- 6.1 All records and plot data should be in a form determined and validated by the Test Centre.
- 6.2 Characters, recording details and instructions are given in Section D. Any variant and abnormal plant or plants resulting from an adverse reaction to husbandry practice are recorded but excluded from analysis.
- 6.3 In the first recording year, characters, as indicated in Section D, are recorded on all candidates and their controls. The data for measured characters are analysed and used to assess the distinctness and uniformity of the candidate varieties. (For details see Section G).
- 6.4 In the second test cycle, characters, as indicated in Section D, are assessed on all varieties in test and the data analysed and, together with those from the first test cycle, used to assess the distinctness and uniformity of the candidate varieties. (For details see Section G).
- 6.5 If a third test cycle is necessary, characters, as indicated in Section D, are recorded on all candidates and their controls. The data for measured characters are analysed and, together with those from the first and second test cycles, are used to assess the distinctness and uniformity of the candidate varieties. (For details see Section G).
- 6.6 If the Test Centre notices unusual or novel characters in a candidate, a note may be made of these at any time and a photographic record made.

7 Communications with the Applicant

- 7.1 The Test Centre will notify the applicant or the agent of any DUS problems at the earliest practical opportunity as they arise during the growing season. All such notifications must be copied to APHA.
- 7.2 In the case of distinctness problems, if confidentiality considerations allow, the applicant should be informed which variety their candidate is not yet distinct from and be invited to submit any information which may help to distinguish them.
- 7.3 If DUS problems arise, applicants will be invited to visit the DUS tests by arrangement so that the material can be examined (if appropriate), and discussions held with the Test Centre.
- 7.4 After each test cycle the results are summarised and reported to the applicant and APHA by the Test Centre.

Section D – Summary of DUS Characteristics to be Assessed, Method of Assessment and Standards Applied

1 Purpose

1.1 The purpose of this section is to summarise the characteristics to be assessed.

2 Scope

2.1 This section summarises characteristics, states of expression, method of observation and standards required for DUS assessment.

3 Responsibilities

3.1 The Test Centre is responsible for co-ordinating the procedures in this summary of characteristics.

4 Organisation

4.1 The minimum duration of tests to assess characteristics is normally two independent growing cycles. Shorter durations may be applied for assessment of additional characteristics. Proposed changes to the number of growing cycles must be approved by the NLSC.

5 DUS Characteristics to be Assessed

5.1 Routine Characteristics

The following table summarises the DUS characteristics to be routinely examined.

Note:

- * a characteristic which must be examined according to the UPOV Guidelines.
- G a grouping characteristic.

Type of observation of characteristics:

- MG Single measurement of a group of plants of parts of plants
- MS Measurement of a number of individual plants or parts of plants
- VG Visual assessment by a single observation of a group of plants or parts of plants

- VS Visual assessment by observation of individual plants or parts of plants
- C Special test

Number of plants or sample size for assessment: Sample size of 60

5.2 Turnip Characteristics Routinely Recorded in DUS Tests

| UK | UPOV TG/37/11 2022 | Character | Material examined | Number of plants or sample size for assessment | Method of assessment and recording | States of expression | D Method and Minimum Distance required | U Method: standard applied |
|----|--------------------------|--|----------------------|--|------------------------------------|-------------------------------------|---|----------------------------------|
| 05 | 1 | Ploidy | Young plants | 2 replicates of 5 plants | By flowcytometer | 2 = diploid 4 = tetraploid | Clear visual difference | Off-type standard |
| | | | | | | | or 1 state | |
| 25 | 2 | Petiole: intensity of anthocyanin coloration | DUS plot | At least 40 plants in total | | 1 = absent or very weak 2 = weak | Clear visual difference | Off-type standard |
| | from 2 replicates | | 3 = medium | or | and | | | |
| | | | | | | 4 = strong | 2 states | Uniformity |
| | | | | | | 5 = very strong | | score >5 |
| 29 | 3 | Leaf: attitude | DUS plot | At least 40 | Visual observation | | Clear visual | Off-type |
| | | | | plants in total | | 2 = erect to semi-erect | difference | standard |
| | | | | from 2 replicates | | 3 = semi-erect | or | and |
| | | | | | | 4 = semi-erect to | 2 states | Uniformity score >5 |
| | | | | | | horizontal 5 = horizontal | | Score >3 |
| 23 | 4 | Leaf: degree of recurving of the apex | DUS plot | At least 40 | Visual observation | 1 = absent or very weak | Clear visual | Off-type |
| | - | | 200 p.u. | plants in total | | 2 = very weak to weak | difference | standard |
| | | | | from 2 replicates | | 3 = weak | or | and |
| | | | | | | 4 = weak to medium | 2 states | Uniformity |
| | | | | | | 5 = medium | | score >5 |
| | | | | | | 6 = medium to strong | | |
| | | | | | | 7 = strong | | |
| | | | | | | 8 = strong to very strong | | |
| 20 | 5 | Leaf: intensity of green colour | DUS plot | At least 40 | Visual observation | 9 = very strong 1 = very light | Clear visual | Off-type |
| | | 20011 Interiority or groom concur | Boo plot | plants in total | | 2 = very light to light | difference | standard |
| | | | | from 2 replicates | | 3 = light | or | and |
| | | | | | | 4 = light to medium | 2 states | Uniformity score >5 |
| | | | | | | 5 = medium 6 = medium to dark | | |
| | | | | | | 7 = dark | | |
| | | | | | | 8 = dark to very dark | | |
| | | | | | | 9 = very dark | | |

| 16 | 6 | Leaf: number of lobes | DUS plot | At least 40 plants | Visual observation | 1 = absent or very few | Clear visual | Off-type |
|----|---|-----------------------|------------------|--------------------|--------------------|------------------------|----------------|------------------|
| | | | or | in total from 2 | or | 2 = very few to few | difference | standard |
| | | | 40 single plants | replicates | visual score | 3 = few | or | and |
| | | | | | or | 4 = few to medium | 2 states | Uniformity score |
| | | | | | counts on 20 | 5 = medium | difference | >5 |
| | | | | | single plants per | 6 = medium to many | or | or |
| | | | | | plot | 7 = many | COYD @ 5% | COYU at 0.1% |
| | | | | | | 8 = many to very many | for both 2 and | for both 2 and |
| | | | | | | 9 = very many | 3 year tests | 3 year tests |

| UK | UPOV TG/37/11 2022 | Character | Material examined | Number of plants or sample size for assessment | Method of assessment and recording | States of expression | D Method and Minimum Distance required | U Method: standard applied |
|----|--------------------------|---|----------------------|---|--|---|---|---|
| 61 | 7 | Only varieties with Leaf: number of lobes: absent or very few: Leaf: depth of the incisions of margin at basal part | DUS plot | At least 40 plants in total from 2 replicates | Visual observation or visual score | 1 = absent or very shallow 2 = very shallow to shallow 3 = shallow 4 = shallow to medium 5 = medium 6 = medium to deep 7 = deep 8 = deep to very deep 9 = very deep | Clear visual difference or 2 states | Off-type standard and Uniformity score >5 |
| 52 | 8 | Leaf: depth of incisions of margin of upper part of leaf | DUS plot | At least 40 plants in total from 2 replicates | Visual observation or visual score | 1 = absent or very shallow 2 = very shallow to shallow 3 = shallow 4 = shallow to medium 5 = medium 6 = medium to deep 7 = deep 8 = deep to very deep 9 = very deep | Clear visual difference or 2 states | Off-type standard and Uniformity score >5 |

| 21 | 9 | Leaf: undulation of margin | DUS plot | At least 40 | Visual observation | 1 = absent or very weak | Clear visual | Off-type |
|------|------------------|----------------------------|------------------|--------------------------------------|-----------------------------|-----------------------------------|---------------------------------|--------------------------------|
| | | | | plants in total | or | 2 = very weak to weak | difference | standard |
| | | | | from 2 replicates | visual score | 3 = weak | or | and |
| | | | | | | 4 = weak to medium | 2 states | Uniformity |
| | | | | | | 5 = medium | | score >5 |
| | | | | | | 6 = medium to strong | | |
| | | | | | | 7 = strong | | |
| | | | | | | 8 = strong to very strong | | |
| | | | | | | 9 = very strong | | |
| 11 | 10 | Leaf: length | DUS plot | At least 40 plants | Visual observation | · · | Clear visual | Off-type |
| | | | or | in total from 2 | | 2 = very short to short | difference | standard |
| | | | 40 single plants | replicates | visual score | 3 = short | or | and |
| | | | | | or | 4 = short to medium | 2 states | Uniformity score |
| | | | | | | 5 = medium | difference | >5 |
| | | | | | per plot | 6 = medium to long | or | or |
| | | | | | per piot | 7 = long 8 = long to very long | COYD @ 5% for both 2 and | COYU at 0.1% |
| | | | | | | 9 = very long | 3 year tests | for both 2 and 3 year tests |
| 17 | 11 | Leaf: width | DUS plot | At least 40 plants | Visual observation | | Clear visual | Off-type |
| 1 | '' | Louis Width | or | in total from 2 | or | 2 = very narrow to narrow | difference | standard |
| | | | 40 single plants | replicates | visual score | 3 = narrow | or | and |
| | | | | | | 4 = narrow to medium | 2 states | Uniformity score |
| | | | | | | 5 = medium | difference | >5 |
| | | | | | measurements | 6 = medium to broad | or | or |
| | | | | | per plot | 7 = broad | COYD @ 5% | COYU at 0.1% |
| | | | | | | 8 = broad to very broad | for both 2 and | for both 2 and |
| | | | | | | 9 = very broad | 3 year tests | 3 year tests |
| 1117 | LIDOV | | M. (| N | Made | 01.1 | | 1111 |
| UK | UPOV TG/37/11 | Character | Material | Number of | Method of | States of expression | D Method and | U Method: |
| | 2022 | | examined | plants or sample size for assessment | assessment and recording | | Minimum Distance required | standard applied |

| 12 | 12 | Leaf: length of terminal lobe | DUS plot | | Visual observation | - | Clear visual | Off-type |
|----|----|--|------------------|--------------------|--------------------|---------------------------|----------------|------------------|
| | | | or | in total from 2 | | 2 = very short to short | difference | standard |
| | | | 40 single plants | replicates | | 3 = short | or | and |
| | | | | | | 4 = short to medium | 2 states | Uniformity score |
| | | | | | J - J | 5 = medium | difference | >5 |
| | | | | | | 6 = medium to long | or | or |
| | | | | | | 7 = long | COYD @ 5% | COYU at 0.1% |
| | | | | | | 8 = long to very long | for both 2 and | for both 2 and |
| | | | | | | 9 = very long | 3 year tests | 3 year tests |
| 13 | 13 | Leaf: width of terminal lobe | DUS plot | At least 40 plants | Visual observation | 1 = very narrow | Clear visual | Off-type |
| | | | or | in total from 2 | | 2 = very narrow to narrow | difference | standard |
| | | | 40 single plants | replicates | visual score | 3 = narrow | or | and |
| | | | | | | 4 = narrow to medium | 2 states | Uniformity score |
| | | | | | | 5 = medium | difference | >5 |
| | | | | | | 6 = medium to broad | or | or |
| | | | | | | 7 = broad | COYD @ 5% | COYU at 0.1% |
| | | | | | | 8 = broad to very broad | for both 2 and | for both 2 and |
| | | | | | | 9 = very broad | 3 year tests | 3 year tests |
| 24 | 14 | Leaf: hairiness of upper side | DUS plot | At least 40 | | 1 = absent or very weak | Clear visual | Off-type |
| | | | | plants in total | | 2 = very weak to weak | difference | standard |
| | | | | from 2 replicates | visual score | 3 = weak | or | and |
| | | | | | | 4 – weak to medium | 2 states | Uniformity |
| | | | | | | 5 = medium | | score >5 |
| | | | | | | 6 = medium to strong | | |
| | | | | | | 7 = strong | | |
| | | | | | | 8 = strong to very strong | | |
| | | | | | | 9 = very strong | | |
| | 15 | Root: degree of swelling | DUS plot | At least 40 | Visual observation | 1 = absent or weak | Clear visual | Off-type |
| | | | | plants in total | or | 2 = medium | difference | standard |
| | | | | from 2 replicates | visual score | 3 = strong | or | and |
| | | | | | | | 2 states | Uniformity |
| | | | | | | | | score >5 |
| 58 | 16 | Only varieties with Root: degree of | DUS plot | At least 40 | Visual observation | 1 = white | Clear visual | Off-type |
| | | swelling: medium or strong: Root: color of | | plants in total | or | 2 = green | difference | standard |
| | | skin <u>above</u> soil | | from 2 replicates | visual score | 3 = yellow-orange | or | and |
| | | | | | | 4 = red | 1 state | Uniformity |
| | | | | | | 5 = reddish purple | | score >5 |
| | | | | | | 6 = bluish purple | | |
| | | | | | | 7 = black | | |

| 95 | 17 | Only varieties with Root: degree of | DUS plot | At least 40 | Visual observation | 1 = light | Clear visual | Off-type |
|----|----------|--|----------|-----------------------------|---------------------------------------|--|--------------------------|------------------|
| | " | swelling: medium or strong: Root: | 200 piot | plants in total | | 2 = medium | difference | standard |
| | | intensity of color of skin <u>above</u> soil | | from 2 replicates | visual score | 3 = dark | or | |
| | | intensity of color of skill <u>above</u> son | | · | Viodal 55515 | 3 – uark | 2 state | |
| | | <u> </u> | | <u> </u> | | | 2 otato | |
| UK | UPOV | Character | Material | Number of | Method of | States of expression | D Method and | U Method: |
| | TG/37/11 | | examined | plants or | assessment | | Minimum | standard |
| | 2022 | | | sample size for | and recording | | Distance | applied |
| 36 | 18 | Only variation with Donty donner of | DUS plot | assessment At least 40 | Visual observation | 4 —bita | required Clear visual | Off to up a |
| 36 | 10 | Only varieties with Root: degree of | DOS PIOL | plants in total | | | _ | Off-type |
| | | swelling: medium or strong: Root: color of | | from 2 replicates | | 2 = yellow | difference | standard |
| | | skin <u>below</u> soil | | nom 2 rophodioc | | 3 = red | or | and |
| | | | | | | 4 = purple | 1 state | Uniformity score |
| | | | | | | 5 = black | | >5 |
| | | | | | | | | |
| | | | | | | | | |
| | 10 | | DUO 1 / | | \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ | 4 19 | 0, , | 0.55 |
| 37 | 19 | Only varieties with Root: degree of | DUS plot | At least 40 plants in total | Visual observation | | Clear visual | Off-type |
| | | swelling: medium or strong: Root: color of flesh | | from 2 replicates | | 2 = yellow | difference | standard |
| | | of flesh | | nom z replicates | visual score | | or | |
| | | | | | | | 1 state | |
| 53 | 20 | Only varieties with Root: degree of | DUS plot | At least 40 | Visual observation | 1 - abcent | Clear visual | Off-type |
| | | swelling: medium or strong: Root: | DOO PIOT | plants in total | | 9 = present | difference | standard |
| | | anthocyanin coloration of flesh | | from 2 replicates | visual score | 9 - present | or | |
| | | anthocyanin coloration of nesh | | · | Viodai 66616 | | 1 state | |
| 99 | 21 | Only varieties with Root: degree of | DUS plot | At least 40 | Visual observation | 1 = very shallow | Clear visual | Off-type |
| | | swelling: medium or strong: Root | 200 plot | plants in total | | 2 = very shallow to | difference | standard |
| | | position in soil | | from 2 replicates | | shallow | or | and |
| | | | | · | | 3 = shallow | 2 states | Uniformity |
| | | | | | | 4 = shallow to medium | 2 otatoo | score >5 |
| | | | | | | 5 = medium | | |
| | | | | | | 6 = medium to deep | | |
| | | | | | | 7 = deep | | |
| | | | | | | 8 = deep to very deep | | |
| | | | | | | 9 = very deep | | |
| 54 | 22 | Only varieties with Root: degree of | DUS plot | At least 40 | Visual observation | | Clear visual | Off-type |
| | | swelling: medium or strong: Root: shape | | plants in total | | 2 = narrow oblate | difference | standard |
| | | in longitudinal section | | from 2 replicates | | 3 = broad oblate | or | and |
| | | | | | | 4 = circular | 2 states | Uniformity |
| | | | | | | 5 = broad oblong | | score >5 |
| | 1 | | | | | 6 = narrow oblong 7 = broad obovate | | |
| | | | | | | r – broad obovate | | |

| | | | | | | 8 = narrow obovate 9 = triangular | | |
|----|----|--|----------|---|--------------|---|--|--|
| 40 | 23 | Only varieties with Root: degree of swelling: medium or strong: Root: length | DUS plot | At least 40 plants in total from 2 replicates | visual score | 2 = very short to short 3 = short 4 = short to medium 5 = medium 6 = medium to long | Clear visual difference or 2 states or COYD @ 5% for both 2 and 3 year tests | |

| UK | UPOV TG/37/11 2022 | Character | Material examined | Number of plants or sample size for assessment | Method of assessment and recording | States of expression | D Method and Minimum Distance required | U Method: standard applied |
|----|--------------------------|--|------------------------------------|---|---|---|---|---|
| 56 | 24 | Only varieties with Root: degree of swelling: medium or strong: Root: curvature of vertical axis | DUS plot | At least 40 plants in total from 2 replicates | Visual observation or visual score | 1 = absent 9 = present | Clear visual difference or 1 state | Off-type standard |
| 55 | 25 | Only varieties with Root: degree of swelling medium or strong: Root: position of broadest part | DUS plot or 40 single plants | At least 40 plants in total from 2 replicates | | 1 = above middle 2 = at middle 3 = below middle | Clear visual difference or 2 states difference | Off-type standard and Uniformity score >5 |
| 41 | 26 | Only varieties with Root: degree of swelling: medium or strong: Root: diameter at broadest part | DUS plot or 40 single plants | At least 40 plants in total from 2 replicates | visual score or 20 single measurements per plot | 1 = very small 2 = very small to small 3 = small 4 = small to medium 5 = medium 6 = medium to large 7 = large 8 = large to very large 9 = very large | Clear visual difference or 2 states difference or COYD @ 5% for both 2 and 3 year tests | Off-type standard and Uniformity score >5 or COYU at 0.1% for both 2 and 3 year tests |
| 57 | 27 | Only Varieties with Root: degree of swelling: medium or strong: Root: shape of collar | DUS plot | At least 40 plants in total from 2 replicates | or visual score | 1 = strongly depressed 2 = strongly depressed to moderately depressed 3 = moderately depressed 4 = moderately depressed to flat 5 = flat 6 = flat to moderately raised 7 = moderately raised 8 = moderately raised to | Clear visual | Off-type standard and Uniformity score >5 |

| | | | | | | strongly raised 9 = strongly raised | | |
|----|----|---|----------|---|--------------|---|--|---|
| 96 | 28 | Only varieties with Root: degree of swelling: medium or strong: Root: shape of apex | DUS plot | At least 40 plants in total from 2 replicates | visual score | 1 = narrow acute 2 = broad acute 3 = rounded 4 = truncate 5 = depressed | Clear visual difference or 2 states | Off-type standard and Uniformity score >5 |

| UK | UPOV TG/37/112 022 | Character | Material examined | Number of plants or sample size for assessment | Method of assessment and recording | States of expression | D Method and Minimum Distance required | U Method: standard applied |
|----|--------------------------|--|----------------------|--|---|---|--|---|
| 30 | 29 | Only varieties with Root: degree of swelling: medium or strong: Time of harvest maturity | DUS plot | At least 40 plants in total from 2 replicates | visual score or dated (one representative value per plot) | 1 = very early 2 = very early to early 3 = early 4 = early to medium 5 = medium 6 = medium to late 7 = late 8 = late to very late 9 = very late | Clear visual difference or 2 states or COYD @ 5% for both 2 and 3 year tests | Off-type standard and Uniformity score >5 or COYU at 0.1% for both 2 and 3 year tests |
| | 30 | Plant: number of sprouts | DUS plot | At least 40 plants in total from 2 replicates | visual score | 1 = very few 2 = very few to few 3 = few 4 = few to medium 5 = medium 6 = medium to many 7 = many 8 = many to very many 9 = very many | Clear visual difference or 2 states | Off-type standard and Uniformity score >5 |
| 01 | 31 | Time of flowering | DUS plot | At least 40 plants in total from 2 replicates | visual score or dated (one representative | , , | Clear visual difference or 2 states or COYD @ 5% for both 2 and 3 year tests | Off-type standard and Uniformity score >5 or COYU at 0.1% for |

| | | | | 8 = late to very late 9 = very late | | both 2 and 3 year tests |
|----|----------------------------------|------------|---|--|--|---|
| 32 | Petal: intensity of yellow colou | r DUS plot | At least 40 plants in total from 2 replicates | 1 = light 2 = medium 3 = dark | Clear visual difference or 2 states | Off-type standard and Uniformity score >5 |

5.3 Previously Approved Characteristics Not Routinely Recorded in DUS Tests

The following table summarises characteristics no longer used which have been approved by the NLSC and can be examined at the request of the applicant where necessary to establish Distinctness. A fee may be charged for examination of these characteristics as advised by APHA, Plant Varieties and Seeds.

| UPOV | UK | Character | Material examined | Number of plants or sample size for assessment | Method of assessment and recording | States of expression | D Method and Minimum distance required | U Method and Standard applied |
|------|-------------------|-------------------|--------------------------------------|--|------------------------------------|--------------------------------|--|--|
| | S | | | | | | | |
| | 02 QN VG/MS | Cotyledon: length | Seedlings raised in glasshouse | At least 40 plants in total from 2 replicates | | 3=short 5=medium 7=long | Clear visual difference or 2 states or COYD @ 5% for both 2 and 3 year tests | Off-type standard and Uniformity score >5 or COYU at 0.1% for both 2 and 3 year tests |
| | 03 QN VG/MS | Cotyledon: width | Seedlings raised in glasshouse | At least 40 plants in total from 2 replicates | or | 3=narrow 5=medium 7=wide | Clear visual difference or 2 states or COYD @ 5% for both 2 and 3 year tests | Off-type standard and Uniformity score >5 or COYU at 0.1% for both 2 and 3 year tests |

| UPOV | UK | Character | Material examined | Number of plants or sample size for assessment | Method of assessment and recording | States of expression | D Method and Minimum distance required | U Method and Standard applied |
|------|-------------------|---|---------------------------------------|---|--|---------------------------------|---|--|
| | 42 QN VG/MS | Root: diameter (1/3 from top) | DUS plot or 40 single plants | At least 40 plants in total from 2 replicates | Visual observation or visual score or 20 single measurements per plot | 3=narrow 5=medium 7=broad | Clear visual difference or 2 states difference or COYD @ 5% for both 2 and 3 year tests | Off-type standard and Uniformity score >5 or COYU at 0.1% for both 2 and 3 year tests |
| | | Root: diameter (1/3 from base) | DUS plot or 40 single plants | At least 40 plants in total from 2 replicates | Visual observation or visual score or 20 single measurements per plot | 3=narrow 5=medium 7=broad | Clear visual difference or 2 states difference or COYD @ 5% for both 2 and 3 year tests | Off-type standard and Uniformity score >5 or COYU at 0.1% for both 2 and 3 year tests |
| | 44 QN VG | Flower: width of petal | DUS plot | At least 40 plants in total from 2 replicates | Visual observation or visual score | 3=narrow 5=medium 7=wide | Clear visual difference or 2 states | Off-type standard and Uniformity score >5 |
| | 46 QN VG | Flower: length of petal | DUS plot | At least 40 plants in total from 2 replicates | Visual observation or visual score | 3=short 5=medium 7=long | Clear visual difference or 2 states | Off-type standard and Uniformity score >5 |
| | 70 QN VG/MG | Plant: stem length after flowering (at harvest) | DUS plot | At least 40 plants in total from 2 replicates | Visual observation or visual score | 3=short 5=medium 7=long | Clear visual difference or 2 states | Off-type standard and Uniformity score >5 |
| | 09 | Leaf: type | DUS plot | At least 40 plants in total from 2 replicates | | 1 = entire 2 = lobed | Clear visual difference or 1 state | Off-type standard |

| 31 | Root: thick cork layer around skin | DUS plot | At least 40 plants in total from 2 replicates | Visual observation or visual score | 1 = absent 9 = present | Clear visual difference or 1 states | Off-type standard |
|----|---|----------|---|--|-------------------------------------|---|---|
| 39 | Root: intensity of yellow colour of flesh | DUS plot | At least 40 plants in total from 2 replicates | Visual observation or visual score | 3 = light 5 = medium 7 = dark | Clear visual difference or 2 states | Off-type standard and Uniformity score >5 |

New Additional DUS Characteristics: Applicants can suggest new characters on the TQ for testing DUS or after notification by the DUS Test Centre of distinctness problems (for procedures see Section F).

Section E – Reference Seed Stock Maintenance and VCU Seed Stock Authentication Procedures, where applicable

1 Purpose

1.1 This section sets out the procedures for reference seed stock maintenance and VCU seed stock authentication (if applicable).

2 Scope

2.1 These procedures apply to all reference collection varieties.

3 Responsibilities

3.1 The Test Centre is responsible for conducting these procedures.

4 Procedures for Reference Seed Stock Maintenance

- 4.1 The seed sample submitted with the successful or pending application is considered to be the definitive stock of the variety. Subject to meeting the required certification standards a small portion of the seed is sown for observation and measurement. The remainder is stored under controlled and monitored storage conditions as part of the official reference collection.
- 4.2 If during the normal tests there is any evidence that a seed stock is deteriorating in storage, or that stocks have low quantity, a request will be made to the maintainer asking for a replacement stock of the variety. This replacement stock must be authenticated by comparing plots established from the replacement seed with that of the definitive seed, over a maximum of two recording cycles.
- 4.3 If the replacement seed sample cannot be visually distinguished from the definitive reference stock, it will be accepted as representing the variety. If there are visual differences, the new sample will be recorded, and will be accepted as representing the variety if there are no significant (P=0.05) differences in the first test cycle, or no significant (P=0.05) differences over two test cycles in a COYD analysis (see associated document UPOV TGP/8/1 for details). It may then be accepted as definitive and substituted for the existing definitive stock in the reference collection. These procedures may be modified, where, in the opinion of the technical officer, differences are the result of environmental or cultural factors.

Section E

- 4.4 A replacement sample or an additional replacement sample will be considered sufficiently uniform after one test cycle, if the level of off-types is the same or less than the number at 1% population standard and 95% acceptance probability, and the standard deviations of the measured characters are not significantly greater at the 0.1% (P=0.001) significance level than that of the mean standard deviations of the control varieties. Over 2 years the additional replacement sample will be considered sufficiently uniform if the Combined Over Years Uniformity (COYU) is not significantly greater at the 0.1% (P=0.001) significance level than that of the reference varieties. These procedures may be modified, where, in the opinion of the technical officer, differences are the result of environmental or cultural factors.
- 4.5 In the event of the replacement sample not meeting the required acceptance standards, an additional replacement sample will be requested. Plots will be established from any additional replacement seed sample and compared over a maximum of two test cycles. If the additional replacement sample does not meet standards, APHA will be informed, and the variety will be deleted from the reference collection and the Variety Lists will be reviewed.

5 Procedures for VCU Seed Stock AuthenticationNot applicable.

6 Procedures for the Inclusion of New Common Knowledge Varieties into the Reference Collection

- 6.1 When a new variety enters into common knowledge it must be included in the reference collection if seed is available. A request will be sent by the Test Centre to the maintainer of the variety and an official description will be requested from the Test Authority which registered the variety. If an official description is provided, seed received will be assumed to be definitive if the seed conforms to the official description. Small differences in the expression of quantitative characters are likely to be the result of recording in a different environment and will be considered as conforming to the description. If no official description is available, seed will be assumed to be definitive.
- 6.2 If the seed does not conform to the official description to an acceptable standard, the sample shall not be accepted as definitive. A replacement sample will be requested and the relevant Testing Authority informed. The standards for the validation will be as for authentication of replacement seed (seed E4).

7 Release of Reference Samples for Authorised Purposes

7.1 Seed of reference samples can be supplied by the Test Centre, on request, to UK and UPOV DUS Testing Authorities and UK and OECD Seed Certification Agencies, provided the recipient is notified in writing that this material, or any material derived from it, must not be supplied to a third party or used for any other purpose than as a reference for official DUS testing or seed certification.

7.2 Provision of reference samples, other than in 7.1, to any other parties must be authorised by APHA.

Section F – Procedures for Assessment of New Additional DUS Characters

1 Purpose

1.1 This section sets out the procedures for assessment of new additional DUS characters for varieties of Turnip entered for Variety Lists and/or PBR trials.

2 Scope

2.1 These procedures apply to applications where additional DUS characteristics which have not been previously approved by the NLSC, are requested for use in the examination of DUS.

3 Responsibilities

- 3.1 The Test Centre is responsible for liaising with the applicant to produce a proposed procedure for the conduct of new tests. This procedure must ensure that Distinctness, Uniformity and Stability requirements will be met.
- 3.2 All new additional characteristics must be authorised by the NLSC in consultation with the PVSC.

4 Reference Varieties

- 4.1 The reference varieties must include varieties from which the candidate variety is not distinct, as well as other varieties for control purposes.
- 4.2 Seed of reference varieties will be supplied by the Test Centre.

5 Procedures

- 5.1 Details of the proposed special test or assessments will be submitted to the NLSC
- 5.2 The NLSC may commission a test or trial to further investigate a proposal. The applicant will be advised by APHA of arrangements and costs.
- 5.3 Where the test for a character is approved by the NLSC it should be subsequently listed in Section D 5.2 or 5.3 as appropriate.

Section G - Procedures for DUS Decisions

1 Purpose

1.1 This section sets out the standards used to assess distinctness, uniformity and stability of varieties of Turnip.

2 Scope

2.1 These procedures apply to all varieties of Turnip entered for Variety List and/or Plant Breeders' Rights tests and those being tested on behalf of Foreign Authorities.

3 Responsibilities

- 3.1 The Test Centre is responsible for applying the criteria for DUS, set out in this procedure.
- 3.2 The Test Centre is responsible for producing the DUS report in accordance with these procedures and for ensuring that they are in accordance with UPOV guidelines.

4 Reference Varieties

4.1 Appendix 1 sets out which varieties are considered as reference varieties for these procedures.

5 Distinctness

- 5.1 In accordance with associated document UPOV TG/1/3, varieties can be considered distinct where they have a different expression in a grouping character.
- 5.2 Distinctness is normally assessed in two independent test cycles, but a candidate variety could be considered distinct after one test cycle if there are no other similar varieties. A third independent test cycle may be undertaken if distinctness is not established after two test cycles.
- 5.3 The distinctness standard applied for qualitative is a difference of one state, unless otherwise indicated in Section D. For pseudo-qualitative characters the distinctness standard is a difference of 1, 2 or 3 states depending on the characteristic.
- 5.4 If a candidate is clearly different in a visually observed quantitative character, it is considered to be distinct, without the need for a repeated observation.

- 5.5 Where varieties are grown in close proximity under the same conditions, and a direct comparison can be made, a candidate is considered to be distinct, without the need for a repeated observation.
- 5.6 Where varieties are not grown in close proximity, a candidate is considered to be distinct if a difference of at least two states (see table in section D 5.2) is recorded in a visually observed quantitative character.
- 5.7 The standard for measured or counted quantitative characters, is, at least, a 5% (P=0.05) significant difference in one character over two or three growing cycles in a Combined Over Years Distinctness (COYD) analysis. Please see associated documents UPOV TGP/8/1 for details.
- 5.8 Where COYD cannot be applied, alternative methods should be considered.
- 5.9 When the number of varieties grown does not provide sufficient degrees of freedom for use of the standard COYD analysis, alternative methods should be adopted. If there is sufficient historical data (at least 5 years and sufficient degrees of freedom) then the long-term LSD is applied. This LSD is calculated using up to 10 years of the most recent data. If there is insufficient historical data, the 2 x 1% method should be used.
- 5.10 Where the candidate has a full complement of data for two test cycles, but there is only data for control varieties for one test cycle, the use of FITC (Fitted Constant program in DUST) may be applied. This situation may arise due to the loss of plant material within plots in any one year or where suitable control varieties were not grown in both test cycles.

 The standard applied for Distinctness in such cases is P=0.01.

6 Uniformity

6.1 Uniformity is assessed for all characteristics used to establish distinctness.

Uniformity based on the assessment of 'Off-types'

- 6.2 The assessment of Off-types is undertaken in both test cycles and the total number of 'off-types' combined should not exceed the number allowed using the population standards.
- 6.3 Off-type plants in the glasshouse or field are identified and marked for exclusion from recording.
- 6.4 For cross-pollinated varieties and hybrid varieties (excluding single cross hybrids), relative uniformity standards are applied; the total number of off-type plants recorded in the test should not exceed that of similar varieties.

Section G

6.5 In vegetatively propagated varieties, single cross hybrids and inbred lines, the total number of off-type plants should not exceed that indicated in UPOV TGP/8/1 using a population standard of 1% and a 95% acceptance probability. In a population of 60 plants, 2 off-types are allowed.

Section G

6.6 Providing that the number of off-types in the first test cycle does not exceed 3, the applicant may submit a new seed sample (Resubmission) in the second test cycle with the aim of meeting the off-type standard. Distinctness will be assessed on data from the original seed submitted in the first test cycle and on data from the resubmitted seed in the second test cycle. The resubmitted seed will be authenticated against the original seed in side-by-side plots.

- 6.7 In addition, the number of aneuploid or inbred plants allowed in F1 hybrids should not exceed the numbers indicated in UPOV TGP/8/1 for a 3% population standard and a 95% acceptance probability.
- 6.8 After the variants have been excluded, the characteristics listed in Section D5 are used to assess the uniformity of the remaining plants, according to the methods described.

<u>Uniformity based on the assessment of general variation where no measurements are</u> recorded.

6.9 Uniformity of continuous variation (quantitative characters) is assessed visually according to the following scale:

Score Description1-5 unacceptable (1 is worst)6-9 acceptable (9 is best)

A candidate with a visual uniformity score of 6 or more is satisfactory.

<u>Uniformity based on the assessment of general variation where measurements are recorded:</u>

6.10 Provided a variety meets the off-type standard, it can be considered sufficiently uniform after two or three, test cycles if, for all measured characters necessary for distinctness, the Combined Over Years Uniformity (COYU) analysis is not significantly greater than that of the reference varieties at the 0.1% (P=0.001) significance level (see document TGP/8/1). In all cases an examination of data from individual test cycles is carried out to investigate the uniformity problem indicated by the COYU result. Decisions on whether any outlier plants (off-types) identified by data analysis should be excluded from the calculation of variety means and standard deviations, should be taken by the Test Centre.

7 Stability

7.1 A variety is considered sufficiently stable when there is no evidence to indicate that it lacks uniformity or fails to conform to the essential characteristics of its description in different submissions or in different tests.

8 DUS Report and Variety Description

- 8.1 Upon completion of the DUS examination the DUS Summary report will be submitted to APHA and will be discussed at the relevant DUS Test Centre Meeting. This report will specify all non-routine characteristics for establishing distinctness.
- 8.2 The final DUS report, including the full variety description for positive results, will be submitted to APHA by the specified date. The characteristics to be used in the description are identified in Section D.

Appendix 1 – Reference Collection Varieties1 Variety Listing and Plant Breeders Rights

- 1.1 The DUS reference collection, for any given category of plant variety comprises the following at the time when the application for the candidate is made:
- 1.1.1 All other candidate varieties already in DUS test in the UK or entering testing at the same time as the candidate.
- 1.1.2 All varieties with UK PBR.
- 1.1.3 All varieties on the OECD variety list that are listed by countries with comparable climatic conditions to the UK.
- 1.1.4 All varieties protected under National PBR (UPOV contracting parties) with comparable climatic conditions to the UK
- 1.1.5 Any varieties nominated by the applicant as being comparable i.e., known to be similar.
- 1.1.6 Any other varieties considered to be comparable i.e., known to be similar by the appropriate Test Centre or DUS Centre Group.
- 1.1.7 Other available comparable varieties in common knowledge.



© Crown copyright 2024

You may re-use this information (excluding logos) free of charge in any format or medium, under the terms of the Open Government Licence v.3. To view this licence visit www.nationalarchives.gov.uk/doc/open-government-licence/version/3/ or email PSI@nationalarchives.gsi.gov.uk

Data Protection:

For information on how we handle personal data visit www.gov.uk and search Animal and Plant Health Agency Personal Information Charter.

This publication is available at www.gov.uk/government/publications

Any enquiries regarding this publication should be sent to us at

www.gov.uk/apha

APHA is an Executive Agency of the Department for Environment, Food and Rural Affairs and also works on behalf of the Scottish Government, Welsh Government and Food Standards Agency to safeguard animal and plant health for the benefit of people, the environment and the economy.