



Animal &
Plant Health
Agency

Great Britain and Northern Ireland Variety List Trials: Trial Procedures for Official Examination of Value for Cultivation and Use (VCU) Harvest 2025

**Cereals – Wheat, Barley, Oats, Triticale, Rye, Spelt
Wheat, Durum Wheat**

Appendices

November 2024

Changes

- Updated title from United Kingdom to Great Britain and Northern Ireland
- Updated year to 2025
- Appendix 4 – Table A
 - KWS UK Ltd – updated location of trial from Orford to Framlingham
 - Agrii – updated location of trial from Great Dunmow to Thaxted
 - Envirofield – updated location of trial from Kidlington to Woodstock
 - Frontier Agriculture Ltd. – removed Saaten Union as Seed Handling Operator
 - SRUC – removed Scottish Agronomy as Seed Handling Operator. Removed West of Ormiston from location of trial
- Appendix 4 – Table C
 - KWS UK Ltd – updated location of trial from Fulbourn to Newton
 - Frontier Agriculture – removed Saaten Union as Seed Handling Operator
 - Removed SRUC Edinburgh Growing Trial Operator
 - Changed SRUC in Ellon, Aberdeenshire to Scottish Agronomy and removed NIAB as Seed Handling Operator
 - Added Scottish Agronomy in St Boswells, Scottish Borders as Growing Trial Operator
- Appendix 4 – Table E
 - Agrii – changed Trial description from “T only” to “T and UnT”
 - NIAB TAG – updated location of trial to Sutton Scotney, Hampshire from Hampshire
- Appendix 4 – Table F
 - NIAB TAG – changed Trial description from “T only” to “T and UnT”
 - Changed SRUC to Scottish Agronomy and removed NIABSHU as Seed Handling Operator
- Appendix 4 – Table H
 - Saaten Union UK Ltd – changed Trial description from “T and UnT” to “T only”
- Added Appendix 4 – Table K
 - Durum Wheat including Saaten Union UK Ltd as Growing Trial Operator
- Appendix 5
 - Removed “(Spring Controls to be reviewed in September 2024)” and updated controls
 - Winter Wheat - Replaced Gleam with Champion. Added Bamford as Additional NL1 Control
 - Winter Wheat – Removed Cleam (VL2) (Feed). Removed (VL2) from LG Astronomer (Biscuit) (VL2). Removed (VL1) from Bnmford (Biscuit) (VL1) and added (Add. VL1)
 - Winter Barley – Removed Buccaneer (Malting) Added Electrum (Malting). Added Add. To LG Caravelle (2 row Feed) (VL1)

- Winter Barley – Replaced Electrum with Buccaneer. Added LG Caravelle as Additional NL1 Control
 - Winter Oats – Replaced Grafton with Peloton
 - Winter Oats – Removed (VL2 Comparator) from Grafton (VL2 Comparator) - naked oat
 - Winter Triticale – Replaced KWS Fido with Lumaco
 - Winter Triticale – Removed KWS Fido. Added Lumaco
 - Spring Wheat - Replaced Hexham with KWS Fixum
 - Spelt Wheat – Added Zllernfit
 - Spring Barley – Removed SY Tungsten and Propino
 - Spring Oats – Removed Aspen. Added Merlin
- Appendix 7
 - Changed year
 - Updated Product Table: Added Ascra Xpro. Removed Fandango, Imtrex and Siltra Xpro. Changed Tebuconazole to Tebucur.
 - Updated Table “WINTER WHEAT”:
 - Removed Contact Trials Manager line.
 - For T0: Replaced Sunorg Pro Metconazole + with Tebucur 250 + and updated rate. Added Arizona.
 - For T1: Removed “For eye spot situation” from Revystar XE+. Removed Antargo + line. Removed Revystar XE+ line. Removed Arizona + line. Added “Optional for a rust situation” for Elatus Era line. Replaced Talius/Justice with Entargo+ and updated rate.
 - For T1.5: Replaced Tebuconazole with Sunorg Pro (Metconazole 250) and updated rate. Updated Arizona to note compulsory timings.
 - For T2: Added note for Univoq + and updated rate. Updated Tebuconazole to Tebuconazole 250 and added note for compulsory regions.
 - For T3: Removed “+” from Comet 200. Added noted for compulsory timings for Arizona.
 - Removed table for “SPRING WHEAT – AUTUMN SOWN”
 - Updated table “SPRING WHEAT – SPRING SOWN”
 - For T1: Replaced Revystar XE with Ascra Xpro and updated rate.
 - For T2: Replaced Revystar XE with Univoq and updated rate.
 - Added Note on yellow rust situation.
 - Updated table “SPRING BARLEY”
 - For T1: Replaced Siltra with Ascra and updated rate.
 - Updated table “WINTER OATS”:
 - For T1: Replaced Siltra with Ascra, updated rate and removed note on Siltra Xpro. Replaced Comet 200 with Tebuconazole 250 and removed note on Comet 200.

- For T2: Replaced Siltra Xpro + with Elatus Era +, updated rate and removed note on Siltra Xpro. Removed Comet 200
 - For T3: removed note regarding Comet 200 and removed + on product. Removed Toledo and updated Tebuconazole to Tebuconazole 250 +.
- Updated table “SPRING OATS”
 - For T0: Removed “+” from Cyflamid.
 - For T1: Replaced Siltra with Ascra, updated rate and removed note for Siltra Xpro. Replaced Comet 200 with Tebuconazole 250.
 - For T2: Replaced Siltra Xpro with Elatus Era and updated rate. Updated Tebuconazole to Tebuconazole 250 +.
- Updated table “WINTER RYE & TRITICALE”
 - For T0: Updated Tebuconazole to Tebuconazole 250.
- Updated Table “SPRING TRITICALE” to include “RYE”
 - Added treatment timings and updated products to reflect both crops.

Contents

| | |
|--|----|
| Scope..... | 6 |
| Appendix 1 – Approved Trial Organisers/ Operators for wheat, barley, oats, triticale, rye and spelt wheat..... | 7 |
| Appendix 2 – Seed treatment products for use on NL trials..... | 8 |
| Appendix 3 – Seed dispatch deadline dates | 9 |
| Appendix 4 – Growing Trial Operators and trial locations | 10 |
| Appendix 5 – Control varieties for VCU assessments | 14 |
| Appendix 6 – Plant growth regulator protocol for cereal variety trials - 2024/2025 | 15 |
| Appendix 7 – Fungicide protocol for cereal variety trials – 2024/25 | 22 |
| Appendix 8 – Moisture content determination for yield | 3 |
| Appendix 9 – Dates by which records should be submitted..... | 6 |
| Appendix 10 – Growth stages of cereals | 7 |
| Appendix 11 – Assessment keys for cereal diseases | 9 |

Scope

This document contains the appendices for the main guidance document:

Trial Procedures for Official Examination of Value for Cultivation and Use (VCU) Harvest 2025 –
Cereals – Wheat, Barley, Oats, Triticale, Rye, Spelt Wheat, Durum Wheat

Appendix 1 – Approved Trial Organisers or Operators for wheat, barley, oats, triticale, rye and spelt wheat.

| Activity | Organisers or operators responsible |
|--|--|
| Data Handling Operators | BioSS* AHDB Cereals and Oilseeds** NIAB*** |
| Trials Organiser | BSPB |
| Pathology Trials Operator | NIAB |
| Trial Inspection and Technical Validation Operator | AHDB Cereals and Oilseeds and SASA |
| Quality Testing Operators | NIAB and Campden BRI |
| Data Review and Standards Setting Operator | NIAB |

*Wheat, Barley, Oats, Spring Triticale and Spring Rye

**Triticale, Rye

*** Spelt wheat
Durum wheat

Appendix 2 – Seed treatment products for use on NL trials

Where there are alternative treatments, all varieties in the trial must be treated with the same product

| Crop | Treatment |
|------------------|---|
| Winter wheat | Prothioconazole + Tebuconazole (Redigo Pro) Kinto Plus (Fluxapyroxad, Triticonazole and Fludioxonil) may be used |
| Winter barley | Prothioconazole + Tebuconazole (Redigo Pro). Raxil Star. Kinto Plus (Fluxapyroxad, Triticonazole and Fludioxonil) may be used |
| Winter oats | Prothioconazole + Tebuconazole (Redigo Pro). Kinto Plus (Fluxapyroxad, Triticonazole and Fludioxonil) may be used |
| Spring wheat | Prothioconazole + Tebuconazole (eg Redigo Pro) |
| Spring barley | Prothioconazole + Tebuconazole (Redigo Pro) |
| Spring oats | Prothioconazole + Tebuconazole (Redigo Pro) |
| Winter Triticale | Prothioconazole + Tebuconazole (Redigo Pro) Kinto Plus (Fluxapyroxad, Triticonazole and Fludioxonil) may be used |
| Rye | Prothioconazole + Tebuconazole (Redigo Pro) Kinto Plus (Fluxapyroxad, Triticonazole and Fludioxonil) may be used |
| Spelt wheat | No Treatment |
| Durum wheat | Prothioconazole + Tebuconazole (eg Redigo Pro) |

Appendix 3 – Seed dispatch deadline dates

VCU seed must be delivered to each Growing Trials Operator or Seed Handling Operator by:

| | |
|---------------|--------------|
| Winter wheat | 15 September |
| Triticale | 1 September |
| Rye | 1 September |
| Spelt wheat | 1 September |
| Durum wheat | 1 September |
| Winter barley | 8 September |
| Winter oats | 15 September |
| Spring wheat | 23 October |
| Spring barley | 15 January |
| Spring oats | 15 January |

Appendix 4 – Growing Trial Operators and trial locations

1 Growing Trial Operators or Seed Handling Operators

A. WINTER WHEAT

| Growing Trial Operator | Seed Handling Operator (if not Trial Operator) | Location of trial | Trial description |
|--------------------------|--|-------------------------------------|-------------------|
| Saaten Union UK Ltd | not applicable | Wold Newton, Lincolnshire | T and UnT |
| NIAB | not applicable | Terrington, Norfolk | T only |
| NIAB | not applicable | Callow, Herefordshire | T and UnT |
| Elsoms Wheat Ltd | not applicable | Spalding, Lincolnshire | T, UnT and L |
| KWS UK Ltd | not applicable | Framlingham, Suffolk | T and UnT |
| KWS UK Ltd | not applicable | Frisby on the Wreak, Leicestershire | T and L |
| RAGT Seeds Ltd | not applicable | Ickleton, Cambridgeshire | T only |
| Agrii | not applicable | Thaxted, Essex | T only |
| Limagrain UK Ltd | not applicable | Elmswell, Bury St Edmunds, Suffolk | T only |
| Syngenta Ltd | not applicable | Horncastle, Lincolnshire | T only |
| Envirofield | not applicable | Woodstock, Oxfordshire | T only |
| NIAB | not applicable | Broughton, Hants | T only |
| NIAB | not applicable | Petham, Kent | T only |
| Frontier Agriculture Ltd | not applicable | Driffield, Yorkshire | T only |
| SRUC | not applicable | East Lothian | T, UnT and L |
| DSV UK Ltd | not applicable | Wardington, Banbury, Oxon | T only |
| Scottish Agronomy | not applicable | Tayside, Angus | T and UnT |

B. SPRING WHEAT

| Growing Trial Operator | Seed Handling Operator (if not Trial Operator) | Location of trial | Trial description |
|-------------------------------|--|---------------------------|-------------------|
| Saaten Union UK Ltd | not applicable | Cowlinge, Suffolk | T and DOP |
| John Innes Enterprise | not applicable | Norwich, Norfolk | T and DOP |
| KWS UK Ltd | not applicable | Fowlmere, Cambridgeshire | T and DOP |
| NIAB | not applicable | Sutton Scotney, Hampshire | T only |
| Stockbridge Technology Centre | Saaten Union | Cawood, North Yorkshire | T only |
| Frontier Agriculture | not applicable | Bleasby, Lincolnshire | T only |

C WINTER BARLEY

| Growing Trial Operator | Seed Handling Operator (if not Trial Operator) | Location of trial | Trial description |
|------------------------|--|-------------------------------|-------------------|
| Agrii | not applicable | Great Dunmow, Essex | T only |
| Saaten Union UK Ltd | not applicable | Cowlinge, Suffolk | T only |
| KWS UK Ltd | not applicable | Newton, Cambridgeshire | T only |
| Syngenta Ltd | not applicable | Great Sturton, Lincolnshire | T and UnT |
| Scottish Agronomy | not applicable | Balgonie, Fife | T and UnT |
| NIAB | not applicable | Wimborne, Dorset | T and UnT |
| NIAB | not applicable | Callow, Herefordshire | T and UnT |
| Frontier Agriculture | not applicable | Drifffield, Yorkshire | T only |
| Scottish Agronomy | not applicable | Ellon, Aberdeenshire | T and UnT |
| Scottish Agronomy | not applicable | St Boswells, Scottish Borders | T and UnT |

D SPRING BARLEY

| Growing Trial Operator | Seed Handling Operator (if not Trial Operator) | Location of trial | Trial description |
|-------------------------------|--|--------------------------------|-------------------|
| KWS UK LTD | not applicable | Fulbourn, Cambridgeshire | T only |
| NIAB | not applicable | Morley, Norfolk | T only |
| Scottish Agronomy | not applicable | Gateside, Nr Kinross | T and UnT |
| Syngenta Ltd | not applicable | Great Sturton, Lincolnshire | T and UnT |
| Stockbridge Technology Centre | Saaten Union | Cawood, North Yorkshire | T only |
| Scottish Agronomy | not applicable | Hill of Fearn, Tain | T only |
| Trials Force | Agrii | Steelstrath, Aberdeenshire | T and UnT |
| NIAB | not applicable | Andover, Hampshire | T and UnT |
| Trials Force | Agrii | Cullen, Banffshire | T only |
| SRUC | not applicable | West of Ormiston, East Lothian | T only |
| Scottish Agronomy | not applicable | Tayside, Angus | UnT only |
| AFBI | Agrii | Crossnacreevy | T and UnT |

E WINTER OATS

| Growing Trial Operator | Seed Handling Operator (if not Trial Operator) | Location of trial | Trial description |
|------------------------|--|---------------------------|-------------------|
| Agrii | not applicable | Great Dunmow, Essex | T and UnT |
| Scottish Agronomy | not applicable | Balgonie, Fife | T and UnT |
| NIAB | not applicable | Callow, Herefordshire | T and UnT |
| NIAB | not applicable | Sutton Scotney, Hampshire | T only |

F SPRING OATS

| Growing Trial Operator | Seed Handling Operator (if not Trial Operator) | Location of trial | Trial description |
|------------------------|--|-----------------------|-------------------|
| Scottish Agronomy | not applicable | Fife | T and UnT |
| Saaten Union | not applicable | Cowlinge, Suffolk | T and UnT |
| NIAB | not applicable | Callow, Herefordshire | T and UnT |
| Scottish Agronomy | not applicable | Aberdeenshire | T and UnT |
| AFBI | Agrii | Crossnacreevy | T and UnT |

G WINTER RYE

| Growing Trial Operator | Seed Handling Operator (if not Trial Operator) | Location of trial | Trial description |
|------------------------|--|-------------------|-------------------|
| Saaten Union UK Ltd | not applicable | Cowlinge, Suffolk | T and UnT |

H SPRING RYE

| Growing Trial Operator | Seed Handling Operator (if not Trial Operator) | Location of trial | Trial description |
|------------------------|--|-------------------|-------------------|
| Saaten Union UK Ltd | not applicable | Cowlinge, Suffolk | T only |

I WINTER TRITICALE

| Growing Trial Operator | Seed Handling Operator (if not Trial Operator) | Location of trial | Trial description |
|------------------------|--|-------------------|-------------------|
| Saaten Union UK Ltd | not applicable | Cowlinge, Suffolk | T only |

J SPRING TRITICALE

| Growing Trial Operator | Seed Handling Operator (if not Trial Operator) | Location of trial | Trial description |
|------------------------|--|-------------------|-------------------|
| Saaten Union UK Ltd | not applicable | Cowlinge, Suffolk | T only |

K DURUM WHEAT

| Growing Trial Operator | Seed Handling Operator (if not Trial Operator) | Location of trial | Trial description |
|------------------------|--|-------------------|-------------------|
| Saaten Union UK Ltd | not applicable | Cowlinge, Suffolk | T only |

2. Pathology Trials Operator

| Pathology Trials Operator | Location of trial |
|---------------------------|-------------------|
| NIAB | UK |

Appendix 5 – Control varieties for VCU assessments

The Control varieties are:

| | | | |
|-------------------------|---|-------------------------|---|
| Winter wheat | LG Skyscraper (Feed) Champion (VL1) (Feed) Skyfall (Bread making) KWS Extase (Bread making) LG Astronomer (Biscuit) Bamford (Biscuit) (Add. VL1) | Spelt wheat | Zollernspelz |
| Winter barley | Craft (Malting) KWS Feeris (6 row Feed) SY Kingsbarn (6 row Hybrid Feed) Electrum (Malting) KWS Tardis (Feed) LG Caravelle (2 row Feed) (Add. VL1) | Spring barley | RGT Planet LG Diablo Laureate Skyway |
| Winter oats | Cromwell Mascani Dalguise Peloton (VL2 Comparator) - naked oat Grafton– naked oat | Spring triticale | Doublet Trimour |
| Winter triticale | Kasyno Lumaco | Spring wheat | KWS Ladum KWS Cochise Hexham KWS Fixum |
| Spring | Miradoux | Spring oats | Canyon WPB Isabel MerlinLennon |
| | | Winter rye | SU Performer |
| | | Spring rye | SU Vergil Ovid |

Appendix 6 – Plant growth regulator protocol for cereal variety trials - 2024/2025

Last updated: December 2023

Recommendations by Paul Gosling, BASIS registration number R/E/8107/IFM.

RL Trials Co-ordinator: Mark Bollebakker

01480 482989

NL Co-ordinator: Jeremy Widdowson

01353 653846

The following notes are given for guidance in the use of plant growth regulators (PGRs) on Recommended List trials. The full manufacturer instructions should be consulted prior to the storage, handling or use of any agrochemical product. The instructions and advice given on product labels should be followed at all times. There should be no conflicting advice between that given in this protocol and on the product label; if there appears to be any conflict, inform the Trials Co-ordinators before the application is made.

Table 1: Summary of applications for NL trials

| Crop | Fungicide treated plots | Plots or DOPs without fungicide | Lodging trials |
|---|---|---|----------------|
| Winter wheat | Yes | No | No |
| Spring wheat (late autumn or spring sown) | Yes | No | not applicable |
| Winter barley | Yes, unless the crop is stressed and the risk of lodging is negligible. | Yes, unless the crop is stressed and the risk of lodging is negligible. | No |
| Spring barley | Only if a high risk of lodging | No | No |
| Winter oats | Yes, +F/+PGR plots only. | Yes | No |
| Spring oats | Yes, +F/+PGR plots only. | No | not applicable |
| Winter rye | Yes | Yes | not applicable |
| Winter triticale | Yes | Yes | not applicable |

Plant growth regulators should not be applied to trials grown specifically for the assessment of lodging. There are important restrictions relating to the use of plant growth regulators.

Crop damage can occur if manufacturers' guidelines are not followed.

Common restrictions are those relating to crops that:

- are sited on soils of low fertility
- are suffering from herbicide damage
- are under stress from drought, waterlogging or any other cause
- were sown in the very late spring

Trial managers should consult the manufacturer's instructions to see if any of these restrictions apply.

IN ALL CASES, PLANT GROWTH REGULATOR SHOULD BE APPLIED TO DESIGNATED PLOTS ONLY AS SPECIFIED IN THE AHDB RECOMMENDED LIST CEREAL TRIALS PROTOCOL

General guidance and notes specific to the use of PGRs on variety trials

Products should be used according to current manufacturers' instructions. It is the responsibility of the Trial Manager to ensure that the growth stages of all of the varieties in the trial are within the manufacturer's guidelines for use. The following notes are intended to highlight matters of particular relevance to the use of PGR products on variety trials and timings are relevant to earliest varieties.

The **Terpal** doses in this protocol are below full rate (**2.0 l/ha**).

Consult the Trials Co-ordinators if you feel that a higher dose is needed e.g. if there is a risk of severe lodging. However, be aware that late applications of **Terpal** (e.g. at around the flag leaf emergence stage) at higher doses can lead to crop damage or stimulate the production of secondary tillers.

Winter wheat and late autumn sown Spring wheat

| Product | Crop | Rate and timing |
|--|-------------------|--|
| Either split dose 3C Chlormequat 750+ Moddus | Winter wheat only | 3C Chlormequat 750 at 1.0 l/ha at G.S. 25 – 30 PLUS †Moddus at 0.1 l/ha if applied at the G.S 30 timing. Followed at G.S. 31 – 32 by 3C Chlormequat 750 at 1.0 l/ha PLUS †Moddus at 0.1 – 0.2 l/ha. Do not apply if any variety is beyond the G.S. 32 timing. |
| <u>OR SINGLE DOSE</u> (Winter wheat and late sown autumn sown spring wheat) 3C Chlormequat 750 + Moddus | Winter wheat | 3C Chlormequat 750 at 1.5 – 2.0 l/ha at GS30 – 31 (in the North and North-west regions this can be delayed to GS32) PLUS †Moddus at 0.1 – 0.2 l/ha. Do not apply if any variety is beyond G.S. 32. |
| | Spring wheat | 3C Chlormequat 750 at 1.25 l/ha at GS30 – 31 (in the North and North-west regions this can be delayed to G.S. 32) PLUS †Moddus at 0.1 – 0.2 l/ha. Do not apply if any variety is beyond G.S. 32. |
| Optional: *Terpal | not applicable | Either single-dose of 0.75 – 1.5 l/ha (depending on lodging risk and *condition of the crop) at GS32 – 37 OR (particularly for the early sown crops) a split dose of 1.0 l/ha at GS33 plus 0.75 l/ha at GS37 |

***Terpal** should not be applied to any variety in the trial if the leaf sheaths have split and the ears are visible.

- **DO NOT** apply **Terpal** if the crop is suffering from herbicide damage or physical stress caused by e.g., waterlogging, drought, take-all.
- **DO NOT** apply in temperatures above 21°C. If, in Winter wheat/Spring wheat trials there are large differences in growth stages; contact the RL Trials Co-ordinator/ NL Coordinator.

† **Moddus** - Apply unless the crop is stressed or the lodging risk is negligible.

Spring sown Spring wheat trials

| Product | Rate and timing |
|---------------------------------------|--|
| Optional: 3C Chlormequat 750 + Moddus | Single half dose of 0.6 l/ha at GS 30 – 31. Consult the Trials Co-ordinator if the crop is late sown and/or under stress. Do not apply if any variety is beyond G.S. 31. |

Terpal and Moddus should not be applied to spring sown Spring Wheat trials.

Winter barley trials

| Product | Rate and timing |
|---------------------------------------|--|
| Optional: 3C Chlormequat 750 + Moddus | Single half dose of 0.6 l/ha at GS 30 – 31. Consult the Trials Co-ordinator if the crop is late sown and/or under stress. Do not apply if any variety is beyond G.S. 31. |
| Moddus | Optional: G.S. 31 – 32 in high fertility situations (0.1 – 0.2 l/ha) |
| *Terpal | 0.75 – 1.0 l/ha at *G.S. 32 – 39 |

*The preferred option for Terpal is for the product to be applied separately from the T2 application. Terpal should not be applied on any variety in the trial if the leaf sheaths have split and the ears are visible.

- Do not apply Terpal if the crop is suffering from herbicide damage or physical stress caused by e.g. waterlogging, drought, take-all.
- Do not apply in temperatures above 21°C.

Spring barley trials

| Product | Rate and timing |
|---------|--|
| Moddus | Optional: Moddus may be applied at 0.1-0.2 l/ha at GS30 for increased rooting and tiller survival and where lodging may be expected. |
| Terpal | Optional: Terpal 0.5 l/ha at GS32-37. Apply only if the risk of lodging is high and the crop shows no signs of stress. |

Winter and Spring oat trials

| Product | Rate and timing |
|--|--|
| Moddus | Optional: Moddus at 0.1 – 0.2 l/ha, in high lodging risk situations and if applied at G.S. 30. |
| 3C Chlormequat 750 Option for an additional Canopy in high-risk situations | Single dose: 3C Chlormequat 750 1.5 – 2.0 l/ha at G.S.31 – 32. A non-ionic wetting agent should be used - see product label. <u>In high-risk situations Canopy 0.75 – 1.5 l/ha up to G.S. 41.</u> |

Apply routinely as a single application to all appropriate trials unless it is felt that the application is inadvisable, in which case the Trials Co-ordinators should be consulted

Winter rye trials

| Product | Rate and timing |
|--------------------------------------|--|
| 3C Chlormequat 750 | Either single dose: 3C Chlormequat 750 at 1.0 l/ha PLUS †Moddus at 0.1 – 0.2 l/ha at G.S. 30 |
| | Or split dose: 3C Chlormequat 750 1.0 l/ha at G.S. 31 – 32 PLUS †Moddus at 0.1 – 0.2 l/ha |
| Medax Max Check timing with label | Medax Max 0.3 - 0.4 kg/ha should be used if Terpal timing is missed at GS37. This is safe up to GS49 and is safer for use on the ear in comparison to Terpal |
| Terpal | Terpal 1.5 – 2.0 l/ha at G.S. 37 |

† Moddus. Apply unless the crop is stressed or the lodging risk is negligible

Winter triticale trials

| Product | Rate and timing |
|---------------------|--|
| 3C Chlormequat 750 | Single dose: 3C Chlormequat 750 at 1.0 l/ha PLUS †Moddus at 0.1 – 0.2 l/ha at GS30 |
| Optional: Medax Max | Medax Max 0.3 – 0.4 kg/ha |

Spring Rye (VL trial)

| Product | Rate and timing |
|--------------------|---|
| 3C Chlormequat 750 | Single dose: 3C Chlormequat 750 at 2.0 l/ha |

Add recommendation for Spring Rye, Meddax Max full rate at GS 25 – 30 (at point of breaking dormancy)

Spring triticale (VL trial)

| Product | Rate and timing |
|--------------------|---|
| 3C Chlormequat 750 | Single dose: 3C Chlormequat 750 at 2.0 l/ha |

Appendix 7 – Fungicide protocol for cereal variety trials – 2024/25

Last updated: December 2023

Recommendations by Paul Gosling, BASIS registration number R\8107\IFM.

| | | |
|-------------------------|------------------|--------------|
| RL Trials Co-ordinator: | Mark Bollebakker | 01480 482989 |
| NL Co-Ordinator: | Jeremy Widdowson | 01353 653846 |

This programme is for use on AHDB Recommended List and National List cereal trials in 2021/22:

- For spring and summer applications to trials for harvest 2022.
- For autumn applications to trials to be harvested in 2023

It is an experimental protocol and is designed to meet the protocol aim of keeping disease levels in treated plots below 5% infection in all varieties and in all trials. It is not intended to follow commercial practice.

Please note that most treatments are compulsory, and the rates and timings specified should be adhered to as closely as possible. The protocol is robust and, if applied correctly, should be effective. **If, however, disease levels rise above 5% (e.g., if weather conditions do not allow optimal application), please contact Mark Bollebakker (RL) or Jeremy Widdowson (NL) to discuss an appropriate course of action.**

Fungicides should be applied at the stated dose rates unless agreed otherwise with the RL Trials Co-ordinator or NL Co-ordinator. Changes to dose rates will only be sanctioned in exceptional circumstances, such as drought-stressed trials under low disease pressure.

Please contact the RL Trials Co-ordinator or NL Co-ordinator if you have any difficulty in sourcing a particular product.

In some cases, two or more products may be available from a company with the same active substances and formulation; if you wish to use such a product and it is not listed in this protocol, contact the RL Trials Co-ordinator or NL Co-ordinator. Generic products should be avoided as they may contain the same active substances but in a different formulation.

Important: Every care has been taken to ensure that all mixtures, rates, and timings are approved, meeting COSSH regulations and manufacturers and statutory guidelines. However, it is the responsibility of the Trial Manager to ensure that they meet all current regulations at the time of application. It is recommended to seek advice from a qualified BASIS advisor for

suitability to local conditions and regulatory compliance. The RL Trials Co-ordinator or NL Co-ordinator should be notified of any conflict between the protocol and current regulations.

In accordance with FRAC guidelines, only two applications of strobilurin fungicides and two SDHI fungicides are to be applied to any crop.

| Product | Active substances | Amount of active substance | Manufacturer |
|----------------|---------------------|----------------------------|--------------------|
| Ascra Xpro | Bixafen | 65g/l | Bayer Crop Science |
| | Fluopyram | 65g/l | |
| | Prothioconazole | 130g/l | |
| Comet 200 | Pyraclostrobin | 200 g/l | BASF |
| Cyflamid | Cyflufenamid | 50 g/l | Certis |
| Elatius ERA | Benzovindiflupyr | 75 g/l | Syngenta |
| | Prothioconazole | 150 g/l | |
| Entargo | Boscalid | 500 g/l | BASF |
| Arizona | Folpet | 500 g/l | Adama |
| Proline 275 | Prothioconazole | 275 g/l | Bayer |
| Prosaro | Prothioconazole | 125 g/l | Bayer |
| | Tebuconazole | 125 g/l | |
| Revystar XE | Mefentrifluconazole | 47.5/100 g/l | BASF |
| | Fluxapyroxad | | |
| Sunorg-Pro | Metconazole | 90g/l | BASF |
| Talius/Justice | Proquinazid | 200 g/l | DuPont |
| Tebucur 250 | Tebuconazole | 250 g/l | Belchim/Rotam |
| Univoq | Fenpicoxamid | 50 g/l | Corteva |
| | Prothioconazole | 100 g/l | |

WINTER WHEAT

| Treatment Timing | Growth Stage (G.S.) - target timing or disease | Product or active ingredient | Rate |
|------------------|--|------------------------------|------------------|
| T0 | G.S. 30 (with no later than when 50% of varieties at G.S.30) | | |
| | | Cyflamid + | 0.25 – 0.35 l/ha |
| | | Tebucur 250 + | 0.6-0.75 l/ha |
| | | Comet 200 | 0.4 – 0.6 l/ha |
| | <i>Note: Arizona is compulsory at T1 and T2 but can only then be used at one timing either at T0, T1.5 or T3</i> | Arizona | 1.0 l/ha |
| T1 | G.S. 32 (with most varieties at G.S. 32) | | |
| | | Revystar XE + | 1.0 – 1.25 l/ha |
| | | Arizona + | 1.0 l/ha |
| | | Talius/Justice | 0.15 l/ha |
| | Optional for a rust situation | Elatus Era | 0.6 l/ha |

| Treatment Timing | Growth Stage (G.S.) - target timing or disease | Product or active ingredient | Rate |
|--|--|------------------------------|------------------|
| Optional for an eyespot situation | | Entargo + | 0.5 l/ha |
| T 1.5 | G.S. 33 (targeting leaf 2 emerging) | | |
| <i>Optional in a rust situation</i> Note: 14-day interval between T1.5 Tebuconazole application and T2 application | | Sunorg Pro (Metconazole 250) | 0.5 l/ha |
| <i>Optional in a Septoria situation</i> | | Prothioconazole | 0.6 l/ha |
| Note: Arizona is compulsory at T1 and T2 but can only then be used at one timing either at T0, T1.5 or T3 | | Arizona | 1.0 l/ha |
| T2 | G.S 39 – 45 and no later than 4 weeks after T1 application | | |
| Please refer to the Corteva website on guidance for water volumes | | Univoq + | 1.0 – 1.5 l/ha |
| <i>Optional nationally but compulsory for the East and Southern regions</i> | | Tebuconazole 250 | 0.75 – 1.0 l/ha |
| | | Arizona | 1.0 l/ha |
| <i>Optional - If mildew established</i> | | Cyflamid | 0.25 – 0.35 l/ha |
| T3 | G.S. 55 – 61 Timing for Fusarium control (very early anthesis preferred). | | |
| | | Prosaro + | 0.8 – 1.0 l/ha |

| Treatment Timing | Growth Stage (G.S.) - target timing or disease | Product or active ingredient | Rate |
|---|--|------------------------------|----------------|
| | | Comet 200 | 0.4 – 0.6 l/ha |
| <p><i>Note: Arizona is compulsory at T1 and T2 but can only then be used at one timing either at T0, T1.5 or T3. If used at T3 it can only be used up to GS59.</i></p> | | Arizona | 1.0 l/ha |
| Post T3 | | | |
| <p>For extreme septoria or brown rust situations please contact the relevant trials co-ordinator.</p> <p><i>Tebucur 250 has a maximum total dose limit of 2l/ha and can be applied up to GS71. In extreme yellow rust situations, there may be flexibility for a third application of Tebucur 250 at 0.5 l/ha, but please contact the relevant trials co-ordinator before doing so.</i></p> | | | |
| NOTE | | | |
| <p>No more than TWO applications of a SDHI product should be applied to any cereal crop</p> | | | |
| <p><i>Depending whether "Knock down" or protectant activity is required, applications of Cyflamid (eradicator) and Talius/Justice (protectant) can be swapped at T0 or T1. DO NOT APPLY CONSECUTIVE APPLICATIONS OF PRODUCTS CONTAINING CYFLAMID</i></p> | | | |
| <p><i>For Cyflamid, the maximum number of treatments is TWO PER CROP on ALL recommended cereals, to be applied ONLY IN SPRING</i></p> | | | |

SPRING WHEAT - SPRING SOWN

| Treatment Timing | Growth Stage (G.S.) - target timing or disease | Product or active ingredient | Rate |
|--|---|------------------------------|------------------------|
| Pre T1 | | | |
| <i>For disease infections before G.S. 29 consult the relevant trials co-ordinator.</i> | | | |
| T1 | G.S. 29 – 31 | | |
| | | Ascra Xpro + | 0.8 - 1.0 l/ha |
| | | Comet 200 + | 0.4 – 0.6 l/ha |
| | | Arizona + | 1.0 l/ha |
| | | Talius/Justice | 0.15 l/ha |
| T2 | G.S. 37 and no later than 3 weeks after T1 application | | |
| | | Univoq + | 1.0-1.5 l/ha |
| | | Arizona | 1.0 l/ha |
| | <i>Optional if mildew established</i> | <i>Cyflamid</i> | <i>0.25 – 0.5 l/ha</i> |
| T3 | G.S. 51 – 61 | | |
| | | Prosaro + | 0.8 l/ha |
| | | Comet 200 + | 0.4 – 0.6 l/ha |
| | <i>If including Arizona, must not exceed G.S. 59</i> | <i>Arizona</i> | <i>1.0 l/ha</i> |

Note

In a yellow rust situation, an application of tebuconazole 250 (0.75- 1.0 l/ha) can be made at an appropriate timing.

WINTER BARLEY

| Treatment Timing | Growth Stage (G.S.) - target timing or disease | Product or active ingredient | Rate |
|--|---|------------------------------------|---|
| Before T0 | | | |
| <i>Optional - If net blotch or mildew present in Autumn or early Spring please contact the trials co-ordinator</i> | | | |
| T0 | G.S. 26 – 30 at start of Spring growth | | |
| | | Proline 275 + | 0.3 – 0.5 l/ha |
| | | Comet 200 | 0.35 – 0.5 l/ha |
| T1 | G.S. 30 – 31 No later than 4 weeks after T0 application | | |
| | | Ascra Xpro + | 0.7 – 1.25 l/ha |
| | | Arizona + | 1.5 l/ha |
| | | Cyflamid | 0.25 – 0.35 l/ha |
| T2 | G.S 39 – 45 (varieties should not exceed G.S. 45) | | |
| | | Revystar XE + | 1.0 – 1.25 l/ha |
| | | Arizona | 1.5 l/ha |
| | <i>Optional: If net blotch or rhynchosporium is developing.</i> Not to be applied after the start of flowering. | <i>Proline 275</i> | <i>0.3 – 0.5 l/ha</i> |
| T3 | G.S. 59 – 61 | | |
| | <i>Optional (TO BE CONSIDERED COMPULSORY IF BROWN RUST IS A RISK). Must not be applied after the start of flowering</i> | <i>Proline 275 + Comet 200</i> | <i>0.3 – 0.5 l/ha 0.35 – 0.5 l/ha</i> |

| | | |
|---|--|--|
| NOTE: Comet 200 must be applied before G.S. 59 | | |
|---|--|--|

SPRING BARLEY

| Treatment Timing | Growth Stage (G.S.) - target timing or disease | Product or active ingredient | Rate |
|---|---|------------------------------|-------------------------|
| T0 | G.S. 13 – 15 | | |
| <i>Optional: If disease is present</i> | | <i>Proline 275</i> | <i>0.2 – 0.4 l/ha</i> |
| T1 | G.S. 25 – 31 Applications at the early end of this range may be necessary if rhynchosporium or mildew are developing. | | |
| | | Ascra Xpro + | 0.6 – 1.0 l/ha |
| | | Arizona | 1.0 l/ha |
| <i>Optional: if mildew is present</i> | | <i>Cyflamid</i> | <i>0.25 – 0.35 l/ha</i> |
| T2 | G.S 45 – 59 (earliest varieties should not exceed G.S.59) no later than 3 weeks after T1 application. If any varieties have passed G.S. 59 contact relevant trials co-ordinator. | | |
| <i>Note: If trial is grown for malting quality Revystar XE must not be applied after G.S. 45</i> | | Revystar XE + | 0.75 – 1.0 l/ha |
| | | Arizona | 1.5 l/ha |
| T3 | G.S. 59 – 69 | | |
| <i>Optional: If net blotch or rhynchosporium or fusarium developing</i> | | <i>Proline 275</i> | <i>0.3 – 0.5 l/ha</i> |

WINTER OATS

| Treatment Timing | Growth Stage (G.S.) - target timing or disease | Product or active ingredient | Rate |
|------------------|---|------------------------------|------------------|
| T0 | G.S. Mid to late tillering | | |
| | | Cyflamid + | 0.25 – 0.35 l/ha |
| | | Prothioconazole | 0.35 l/ha |
| T1 | G.S. 31 | | |
| | | Ascra Xpro + | 0.7 - 1.2 l/ha |
| | | Talius/Justice | 0.15 l/ha |
| | <i>Optional - if crown rust is a problem.</i> | <i>Tebuconazole 250</i> | <i>0.5 l/ha</i> |
| T2 | G.S. 39 – 45 | | |
| | | Elatus Era + | 0.6 – 0.8 l/ha |
| | | Cyflamid | 0.25 – 0.35 l/ha |
| T3 | G.S. 45 – 59 | | |
| | <i>Optional: If crown rust pressure has remained high before G.S.59 – 61.</i> | <i>Comet 200</i> | <i>0.5 l/ha</i> |
| | | Tebuconazole 250 + | 0.5 l/ha |

SPRING OATS

| Treatment Timing | Growth Stage (G.S.) - target timing or disease | Product or active ingredient | Rate |
|------------------|---|------------------------------|-------------------------|
| T0 | G.S. 13 – 15 | | |
| | <i>Optional: If mildew present.</i> | <i>Cyflamid</i> | <i>0.25 – 0.35 l/ha</i> |
| | | Prothioconazole | 0.35 l/ha |
| T1 | G.S. Mid to late tillering | | |
| | | Ascra Xpro + | 0.7 – 1.2 l/ha |
| | | Talius/Justice | 0.2 l/ha |
| | <i>Optional: If crown rust is a problem.</i> | <i>Tebuconazole 250</i> | <i>0.5 l/ha</i> |
| T2 | G.S. 39 – 45 | | |
| | | Elatus Era + | 0.6 – 0.8 l/ha |
| | | Cyflamid | 0.25 – 0.35 l/ha |
| | <i>Optional: If crown rust pressure is a problem.</i> | <i>Comet 200+</i> | <i>0.5 l/ha</i> |
| | | <i>Tebuconazole 250 +</i> | <i>0.5 l/ha</i> |

WINTER RYE & TRITICALE

| Treatment Timing | Growth Stage (G.S.) - target timing or disease | Product or active ingredient | Rate |
|--|--|------------------------------|------------------|
| T0 | G.S. 30 | | |
| | | Tebuconazole 250 + | 0.75 – 1.0 l/ha |
| | | <i>Cyflamid</i> | 0.25 – 0.35 l/ha |
| <i>Optional: If mildew present, TRIALS OPERATORS' DISCRETION WETHER TO USE EITHER AT T0 OR T1.</i> | | | |
| T1 | G.S. 31 – 32 | | |
| | Increase rate for high rust. | Elatus Era + | 1.0 l/ha |
| | | <i>Cyflamid</i> | 0.25 – 0.35 l/ha |
| <i>Optional: If mildew present, TRIALS OPERATORS' DISCRETION WETHER TO USE EITHER AT T0 OR T1.</i> | | | |
| T2 | G.S. 39 – 45 | | |
| | | Revystar XE | 0.5 – 1.0 l/ha |
| T3 | G.S. 59 – 61 | | |
| | | <i>Prosaro</i> | 0.8 l/ha |
| <i>Optional: Rye only if rust remain a problem before G.S.61.</i> | | | |

SPRING RYE AND TRITICALE

| Treatment Timing | Growth Stage (G.S.) - target timing or disease | Product or active ingredient | Rate |
|------------------|--|------------------------------|--------------------|
| T1 | G.S. 31 – 32 | | |
| | | Tebuconazole 250 + | 0.75 – 1.0 l/ha |
| | | Talius/Justice | 0.15 l/ha |
| T2 | G.S. 32 – 45 | | |
| | | Elatus Era | 1.0 l/ha |

Recommended List Fungicide and PGR Ag Chem product labels

Introduction

This list can be used to help find the relevant product labels for PGRs, fungicides, herbicides, and insecticides listed in the main protocols. The links below can be used to navigate to the company page and search for the product labels.

Adama UK

[Listen > Learn > Deliver | ADAMA](#)

BASF

[Product Labels, MSDSs & EISs \(basf.co.uk\)](#)

Bayer Crop Science

[Crop Protection Products from Bayer Crop Science UK](#)

Belchim UK

[Products UK – Belchim Crop Protection](#)

Certis Europe

[Products A-Z and Key Documents | Certis Europe United Kingdom](#)

Corteva/DU Pont

[Find a Product | Corteva Agriscience](#)

Syngenta

[Labels and SDS downloads | Syngenta](#)

Appendix 8 – Moisture content determination for yield

Yield data must be corrected to 15% moisture content. In order to do this, the moisture content of the harvested plot grain is required. One of three methods of determining dry matter must be used – the oven method, electronic moisture analysers method or NIR (Near Infrared) determination.

1 Oven method

Samples are dried until constant mass is achieved. For expediency it is permissible to dry samples for a fixed time provided it can be demonstrated that this is sufficient to reliably achieve constant mass for samples even when the chosen apparatus is fully loaded with samples.

Apparatus and Equipment.

Oven. Electrically heated and controlled in such a way that, during normal working, the mean temperature of the air and of the shelves carrying the test samples is 100° C and operates within the range 96 - 104° C. (Temperature to be reviewed by the Procedures Development Group).

The oven should be regularly maintained and regularly checked for correct operation.

Sample drying trays. Durable under test conditions and being of a size which enables the test sample to be distributed evenly within the tray and at depth which does not protract the drying time.

Balance. Accuracy 0.1 g \pm 0.05 g. The balance should be regularly serviced and calibrated. Frequent checks on its correct operation should be made during the period when the balance is in use.

Method

The test samples are received direct from the combine in hermetically sealed bags or containers. Weigh a fully representative 100 g sub-sample or an accurately recorded catch-weight between 100-200 g and place into the drying tray with an identifying label.

Place the drying trays containing the test samples into the pre-heated oven. Dry the test samples for the pre-determined period or until constant mass is achieved (see below).

Remove the test samples from the oven and allow to cool to ambient temperature.

Record the dry weight of the test sample to 0.1 g.

If achievement of constant mass is to be directly measured, five check samples should be removed from a range of positions within the oven after a period of about 16hrs. The dry weight of these samples should be recorded as above. The check samples should be returned to the oven and dried for a further 2 hours and the dry weight again recorded. A dry matter content of less than 0.3% between the two determinations will be accepted as representing constant mass. If constant mass has not been achieved, the check samples should be returned to the oven for further periods of two hours until constant mass is observed.

Results

The dry matter content of the test sample is calculated as follows;

$$\text{Dry matter (\%)} = \frac{\text{Dry test sample weight}}{\text{Original test sample weight}} \times 100$$

When all samples from a given trial have been recorded, the fresh and dry weights are immediately reported to the Data Handling Operator electronically. When the dry weights are reported as a percentage, the fresh weight should be reported as 100.

2 Electronic moisture assessment (moisture analysers)

Principles

Moisture analysers, either separate instruments or probes on combines, may be used for determining the dry matter of harvested grain. There are no restrictions on the make or model of moisture analyser that may be used, provided the conditions described below are met.

The manufacturer's recommendations for use must be followed. On-combine analysis must only be carried out on equipment specifically manufactured for this purpose. 'Desktop' analysers should not be used on the combine because it has been shown that heat and vibration can cause inaccuracy.

Equipment

The analysing equipment must:

- be calibrated at least once annually for each crop according to the manufacturer's instructions using check samples (see reference below) and have a moisture content accuracy of plus/minus 0.5%. The calibration data should be retained for a minimum of 1 year.

- be serviced regularly, especially just prior to harvest, according to manufacturer recommendations. The action taken should be documented and the information held for a minimum of 1 year.
- be fit for use in accordance with manufacturer instructions. It should have an adequate power supply throughout operation. Instructions should be held with the machine and all operators adequately trained in its operation.

In the field:

- the determination of dry matter must be the same for all plots in a trial replicate. For this reason, there should be minimal risk of rainfall during the harvest of a replicate. If there is a significant risk then backup samples should be taken from all plots to allow comparison through the oven method.
- the grain samples to be analysed must be between 83 and 88% dry matter (12 to 17% moisture content). If it is possible that samples in a replicate may fall outside this range, samples must be taken from all plots so that the oven method may be used should it be necessary. Polythene bags and plot identity labels must be carried at all times to allow this to be carried out.
- The grain to be analysed must be fully ripe with no green ears/grains in any sample. In these cases the samples for the oven method should be used.
- The data sent to the Data Handling Operator must be in the form DRY MATTER%.

References:

BS 4317-24:1990, ISO 7700/1-1984 Methods of test for cereals and pulses. Method of checking the calibration of moisture meters for cereals.

3 NIR determination

The NIR method is permitted for the measurement of moisture content provided that the instrument uses current UK NIR Network calibrations for the appropriate crops. The operator must also participate in the monthly ring checks for the various calibrations being used to demonstrate that the instrument and operating practices are performing within specification. Records of the results of the monthly ring checks should be available for inspection if required.

Appendix 9 – Dates by which records should be submitted

1 To Trials Organiser

| Record | Latest date of receipt by Trials Organiser |
|--|---|
| Site data part 1 (including site sketch) | Within 2 months of drilling trial (autumn sown trials) Within 1 month of drilling trial (spring sown trials) |
| Site data part 2 | By the time trials harvested |
| Plot records (in approved electronic format) | Growing Trial Operator should notify Trials Organiser that trial has been harvested within 2 days of harvest |

2 Plot records to Data Handling Operator

| Record | Date |
|---|---|
| Plot records SHOULD be sent to Data Handling Operator | Yield and moisture data within 3 days of harvest Other data within 10 days of record being taken |

3 Plot samples to Quality Testing Operator

| Samples | Date |
|---|--------------------------|
| Plot samples for quality testing SHOULD be sent to Quality Testing Operator | Within 2 days of harvest |

Appendix 10 – Growth stages of cereals

SEEDLING GROWTH

- 10 first leaf through coleoptile
- 11 first leaf unfolded
- 12 2 leaves unfolded
- 13 3 leaves unfolded
- 14 4 leaves unfolded
- 15 5 leaves unfolded
- 16 6 leaves unfolded
- 17 6 leaves unfolded
- 18 8 leaves unfolded
- 19 9 or more leaves unfolded

TILLERING

- 20 main shoot only
- 21 main shoot and 1 tiller
- 22 main shoot and 2 tillers
- 23 main shoot and 3 tillers
- 24 main shoot and 4 tillers
- 25 main shoot and 5 tillers
- 26 main shoot and 6 tillers
- 27 main shoot and 7 tillers
- 28 main shoot and 8 tillers
- 29 main shoot and 9 or more tillers

STEM ELONGATION

- 30 Ear at 1 cm
- 31 1st node detectable
- 32 2nd node detectable
- 33 3rd node detectable
- 34 4th node detectable
- 35 5th node detectable
- 36 6th node detectable
- 37 flag leaf just visible
- 39 flag leaf ligule/collar just visible

BOOTING

- 41 flag leaf sheath extending
- 43 boots just visibly swollen
- 45 boots swollen
- 47 flag leaf sheath opening
- 49 first awns visible

INFLORESCENCE (EAR EMERGENCE)

- 51 First spikelet of inflorescence just visible
- 52 $\frac{1}{4}$ of inflorescence emerged
- 55 $\frac{1}{2}$ of inflorescence emerged
- 57 $\frac{3}{4}$ of inflorescence emerged
- 59 inflorescence completed

ANTHESIS

- 60 beginning of anthesis
- 61
- 64 anthesis half-way
- 65
- 68 anthesis completed
- 69

MILK DEVELOPMENT

- 71 caryopsis watery ripe
- 73 early milk
- 75 medium milk
- 77 late milk

DOUGH DEVELOPMENT

- 83 early dough
- 85 soft dough
- 87 hard dough

RIPENING

- 91 caryopsis hard (difficult to divide by thumb-nail)
- 92 caryopsis hard (can no longer be dented by thumb-nail)
- 93 caryopsis loosening in daytime

Reference: Tottman D R, Broad H (1987) Decimal Code for the Growth Stages of Cereals
Annals of Applied Biology 100, 683-687.

Appendix 11 – Assessment keys for cereal diseases

- 1) Examine top 4 leaves. If top leaf has been fully expanded for less than 14 days, refer to 2nd leaf as 'top leaf'.
- 2) Ignore all naturally senescent leaf tissue.
- 3) Include all chlorosis and necrosis attributable to disease.
- 4) Record % infection; use interpolated values (e.g. 3%) if necessary.

If foci present, record average over the plot as a whole.

| % Infection | Crown rust | Yellow rust | Brown rust |
|-------------|--|--|--|
| 0 | No infection observed | | |
| 0.1 | 1 small cluster of pustules per tiller | 1 stripes per tiller | 25 pustules per tiller |
| 1 | 1 cluster per leaf | 2 stripes per leaf | 100 pustules per leaf |
| 5 | Most tillers infected but some top leaves uninfected | Most tillers infected but some top leaves uninfected | Top leaf – numerous pustules but leaves appear green overall |
| 10 | All leaves infected but leaves appear green overall | All leaves infected but leaves appear green overall | Top leaf – pustules sufficiently dense to give brown appearance in patches |
| 25 | Leaves appear ½ infected ½ green | | |
| 50 | Leaves appear more infected than green | | |
| 75 | Very little green leaf tissue left | | |
| 100 | Leaves dead - no green tissue left | | |

| % Infection | Septoria | Rhynchosporium | Net blotch |
|--------------------|---|--|---|
| 0 | No infection observed | | |
| 0.1 | 1 lesion per 10 tiller | 1 lesion per 10 tillers | 1 small lesion per 10 tillers |
| 1 | 2 small lesions per tiller | 1 lesion per tiller | 1 small lesion per tiller |
| 5 | Small lesions beginning to form areas of dead tissue across width of leaf | Discrete lesions on most tillers, about 2 per leaf | 2 lower leaves appear ¼ infected. Other leaves - few lesions |
| 10 | 2 lower leaves – large areas of diseased tissue some covering 1/3 of leaf | Lesions coalescing but leaves appear green overall | 2 lower leaves appear ½ infected. Other leaves - numerous lesions |
| 25 | Leaves appear ½ infected ½ green | | |
| 50 | Leaves appear more infected than green | | |
| 75 | Very little green leaf tissue left | | |
| 100 | Leaves dead - no green tissue left | | |

| % Infection | Mildew |
|-------------|--|
| 0 | |
| 0.1 | 3 pustules per tiller |
| 1 | 5 pustules per leaf |
| 5 | 2 lower leaves appear $\frac{1}{4}$ infected |
| 10 | 2 lower leaves appear $\frac{1}{2}$ infected |
| 25 | Leaves appear $\frac{1}{2}$ infected $\frac{1}{2}$ green |
| 50 | Leaves appear more infected than green |
| 75 | Very little green leaf tissue left |
| 100 | Leaves dead - no green tissue left |

| % Infection | Ramularia |
|-------------|--|
| 0 | |
| 1 – 5 | Sparse lesions on upper leaves |
| 6 – 10 | More lesions on upper leaves |
| 11 – 20 | Numerous lesions on middle and upper leaves with some necrosis |
| 21 – 30 | Many lesions and severe necrosis on upper leaves and lesions on middle leaves |
| 31 – 40 | Extensive lesions on upper leaves many lesions on middle leaves and necrosis |
| 41 – 50 | Severe damage to upper leaves more lesions and necrosis on middle and lower leaves |
| 51 – 75 | 100% lesions on upper leaves severe necrosis on middle leaves |
| 75 – 100 | Almost all leaves necrotic with lesions on all leaves |

Ear Blight of wheat (*Fusarium* Spp.)

Percentage area infected
(0 = Healthy)

Notes on assessment



- 1) Carry out the assessment between GS 80-90.
- 2) Conduct a 'whole-plot' assessment using the diagram above as a guide to infection levels.
- 3) Estimate the infection level at several points in the plot, giving a single score per plot that is representative of the whole plot.

For further details see Parry D W, Bayles R A & Priestley R H (1984).
Resistance of winter wheat varieties to Ear Blight (*Fusarium culmorum*).
Journal the National Institute of Agricultural Botany 16, 465-468.

Wheat glume blotch (*Septoria nodorum* Berk.)



Percentage of ear affected

Notes on assessment

- 1) Carry out the assessment between GS 80-90.
- 2) Conduct a 'whole-plot' assessment using the diagram above as a guide to infection levels.
- 3) Estimate the infection level at several points in the plot, giving a single score per plot that is representative of the whole plot.

For further details see Parry D W, Bayles R A & Priestley R H (1984).
Resistance of winter wheat varieties to Ear Blight (*Fusarium culmorum*).
Journal the National Institute of Agricultural Botany 16, 465-4

For further information and photos see the HGCA cereals encyclopaedia at <http://cereals.ahdb.org.uk/>.



© Crown copyright 2024

You may re-use this information (excluding logos) free of charge in any format or medium, under the terms of the Open Government Licence v.3. To view this licence visit www.nationalarchives.gov.uk/doc/open-government-licence/version/3/ or email PSI@nationalarchives.gsi.gov.uk

This publication is available at www.gov.uk/government/publications

Any enquiries regarding this publication should be sent to us at

webmaster@apha.gov.uk

www.gov.uk/apha

The Animal and Plant Health Agency (APHA) is an executive agency of the Department for Environment, Food & Rural Affairs, and also works on behalf of the Scottish Government and Welsh Government.