

Monthly water situation report: North East

1 Summary – November 2024

November was a very dry month with only one day of significant rainfall recorded. Monthly rainfall totals were classed as below average for November. Monthly mean river flows have decreased this month at all indicator sites and fall within the notably low, below normal and normal ranges. Groundwater levels vary across the area and generally remain above normal due to the high rainfall recorded in previous months. Soil moisture deficit (SMD) data still shows wet soils across the area. All reservoirs in the area have seen a decrease in stocks this month but remain healthy for the time of year.

1.1 Rainfall

Monthly rainfall totals were classed as below the long term average (LTA) for all catchments across the area. Monthly rainfall totals ranged from 42% of the LTA in the Seaham catchment to 47% of the LTA in the Northumbria North Sea Tribs catchment.

Analysis of the daily rainfall shows a very dry first 3 weeks of November with almost no rainfall recorded across the area. Significant rainfall was recorded on 23 November with two thirds of the months total recorded on that one day at some rain gauges. The final week of November was also dry with little rainfall recorded.

Cumulative 3-month rainfall totals are all in the normal range.

1.2 Soil moisture deficit and recharge

Soils are classed as wet with less than 10mm soil moisture deficit across the area.

1.3 River flows

Monthly mean river flows have decreased this month at all indicator sites and fall within the notably low, below normal or normal ranges. Monthly mean flows ranged from 35% of the LTA at Hartford Bridge on the River Blyth to 57% of the LTA at Rutherford Bridge on the River Greta.

Analysis of the daily mean flows shows that flows were in the notably high or above normal ranges at the start of the month. Daily mean flows decreased at all indicator sites over the first 3 weeks of November following a period of very little rainfall. Daily mean flows increased on 23 November following a 24-hour period of intense rainfall. In the final week of November flows decreased across all indicator sites following a dry period and fell within the normal, below normal or notably low ranges by the end of the month.

1.4 Groundwater levels

Groundwater levels vary across the area. Levels in West Hall Farm borehole in the Wear Magnesian Limestone, remain in the exceptionally high range. Aycliffe NRA2 in the Skerne Magnesian Limestone has decreased slightly and now falls within the notably high range. Royalty Observation is classed as above normal in the Fell Sandstone. Red Lion in the Skerne Magnesian Limestone and Town Law on the Fell Sandstone fall within the normal ranges for November.

1.5 Reservoir stocks

All reservoirs in the area have seen a decrease in stocks this month, with the exception of Kielder which recorded a small increase. Stocks in the Durham group, Lune and Balder Group and Cow Green reservoir ended the month at above 90%. Overall, reservoir stocks remain healthy for the time of year.

Reservoir or reservoir group	Percentage of current stocks	Percentage of previous month stocks
Kielder	81.9	81.5
North Tynedale group	76.6	80.1
Derwent	85.9	89.2
Durham group	90.1	92.1
Lune and Balder group	97.4	98.5
Cow Green	97.6	100


Author: GWHCL North East, hydrology.northeast@environment-agency.gov.uk

All data are provisional and may be subject to revision. The views expressed in this document are not necessarily those of the Environment Agency. Its officers, servants or agents accept no liability for any loss or damage arising from the interpretation or use of the information, or reliance upon views contained in this report.

2 Rainfall

2.1 Rainfall map


Figure 2.1: Total rainfall for hydrological areas for the current month (up to 30 November), the last 3 months, the last 6 months, and the last 12 months, classed relative to an analysis of respective historic totals. November rainfall totals were classed as notably low across the area. Table available in the appendices with detailed information.



Rainfall data for 2024, extracted from Environment Agency 1km gridded rainfall dataset derived from Environment Agency intensity rain gauges. (Source: Environment Agency. Crown Copyright, 100024198, 2024). Rainfall data prior to 2023, extracted from Met Office HadUK 1km gridded rainfall dataset derived from registered rain gauges (Source: Met Office. Crown copyright, 2024).

2.2 Rainfall charts

Figure 2.2: Monthly rainfall totals for the past 12 months as a percentage of the 1961 to 1990 long term average for each hydrometric area in the North East. Monthly rainfall totals are classed as below average for all catchments for November.




Rainfall data for 2024, extracted from Environment Agency 1km gridded rainfall dataset derived from Environment Agency intensity rain gauges. (Source: Environment Agency. Crown

Copyright, 100024198, 2024). Rainfall data prior to 2023, extracted from Met Office HadUK 1km gridded rainfall dataset derived from registered rain gauges (Source: Met Office. Crown copyright, 2024).

3 Soil moisture deficit

3.1 Soil moisture deficit map

Figure 3.1: Map to the left shows soil moisture deficits for week ending 30 November. Map on the right shows the difference (mm) between the actual soil moisture deficit and the 1961 to 1990 long term average soil moisture deficits MORECS data for real land use. Soils are classed as wet with less than 10mm soil moisture deficit across the area.




(Source: Met Office. Crown copyright, 2024). All rights reserved. Environment Agency, 100024198, 2024.

4 River flows

4.1 River flows map






Figure 4.1: Monthly mean river flow for indicator sites for November 2024, expressed as a percentage of the respective long term average and classed relative to an analysis of historic November monthly means. Monthly means are classed as normal at Hartford Bridge and below normal at Rothbury, Rutherford Bridge and Witton Park. Flows are classed as notably low at Haydon Bridge, Heaton Mill and Middleton in Teesdale. There are current ongoing data quality issues at Mitford on the River Wansbeck and therefore the site is marked as no data for the month. Table available in the appendices with detailed information.






Source: Environment Agency). Crown copyright. All rights reserved. Environment Agency, 100024198, 2024.

4.2 River flow charts

Figure 4.2: Daily mean river flow for index sites over the past year, compared to an analysis of historic daily mean flows, and long term maximum and minimum flows. There are current ongoing data quality issues at Mitford on the River Wansbeck and therefore there is no data from October 2023 onwards and hence no graph below.






Source: Environment Agency, 2024.

5 Groundwater levels

5.1 Groundwater levels map







Figure 5.1: Groundwater levels for indicator sites at the end of November 2024, classed relative to an analysis of respective historic November levels. Groundwater levels are classed as exceptionally high at West Hall Farm on the Magnesian Limestone, as notably high at Aycliffe and normal at Red lion. In the north of the area Town Law is classed as normal and Royalty Observation is classed as above normal on the Fell Sandstone. Table available in the appendices with detailed information.



(Source: Environment Agency). Geological map reproduced with kind permission from UK Groundwater Forum, BGS copyright NERC. Crown copyright. All rights reserved. Environment Agency, 100024198, 2024.

5.2 Groundwater level charts


Figure 5.2: End of month groundwater levels at index groundwater level sites for major aquifers in the North East. 12 months compared to an analysis of historic end of month levels and long term maximum and minimum levels.



Source: Environment Agency, 2024.

6 Reservoir stocks

Figure 6.1: End of month reservoir stocks compared to long term maximum, minimum and average stocks. Note: Historic records of individual reservoirs and reservoir groups vary in length.



(Source: Water Company).

7 Glossary

7.1 Terminology

Aquifer

A geological formation able to store and transmit water.

Areal average rainfall

The estimated average depth of rainfall over a defined area. Expressed in depth of water (mm).

Artesian

The condition where the groundwater level is above ground surface but is prevented from rising to this level by an overlying continuous low permeability layer, such as clay.

Artesian borehole

Borehole where the level of groundwater is above the top of the borehole and groundwater flows out of the borehole when unsealed.

Cumecs

Cubic metres per second (m^3s^{-1}).

Effective rainfall

The rainfall available to percolate into the soil or produce river flow. Expressed in depth of water (mm).

Flood alert and flood warning

Three levels of warnings may be issued by the Environment Agency. Flood alerts indicate flooding is possible. Flood warnings indicate flooding is expected. Severe flood warnings indicate severe flooding.

Groundwater

The water found in an aquifer.

Long term average (LTA)

The arithmetic mean calculated from the historic record, usually based on the period 1961 to 1990. However, the period used may vary by parameter being reported on (see figure captions for details).

mAOD

Metres above ordnance datum (mean sea level at Newlyn Cornwall).

MORECS

Met Office Rainfall and Evaporation Calculation System. Met Office service providing real time calculation of evapotranspiration, soil moisture deficit and effective rainfall on a 40 by 40 km grid.

Naturalised flow

River flow with the impacts of artificial influences removed. Artificial influences may include abstractions, discharges, transfers, augmentation and impoundments.

NCIC

National Climate Information Centre. NCIC area monthly rainfall totals are derived using the Met Office 5 km gridded dataset, which uses rain gauge observations.

Recharge

The process of increasing the water stored in the saturated zone of an aquifer. Expressed in depth of water (mm).

Reservoir gross capacity

The total capacity of a reservoir.

Reservoir live capacity

The capacity of the reservoir that is normally usable for storage to meet established reservoir operating requirements. This excludes any capacity not available for use (for example, storage held back for emergency services, operating agreements or physical restrictions). May also be referred to as 'net' or 'deployable' capacity.

Soil moisture deficit (SMD)

The difference between the amount of water actually in the soil and the amount of water the soil can hold. Expressed in depth of water (mm).

7.2 Categories

Exceptionally high

Value likely to fall within this band 5% of the time.

Notably high

Value likely to fall within this band 8% of the time.

Above normal

Value likely to fall within this band 15% of the time.

Normal

Value likely to fall within this band 44% of the time.

Below normal

Value likely to fall within this band 15% of the time.

Notably low

Value likely to fall within this band 8% of the time.

Exceptionally low

Value likely to fall within this band 5% of the time.

8 Appendices

8.1 Rainfall table

Hydrological area	November 2024 rainfall % of long term average 1961 to 1990	November 2024 band	September 2024 to November 2024 cumulative band	June 2024 to November 2024 cumulative band	December 2023 to November 2024 cumulative band
Northumbria North Sea Tribs	47	Notably Low	Normal	Below normal	Notably high
Seaham Area	42	Notably Low	Normal	Normal	Above normal
Tees	45	Notably Low	Normal	Normal	Notably high
Tweed	44	Notably Low	Normal	Below normal	Notably high
Tyne	45	Notably Low	Normal	Normal	Notably high
Wear	42	Notably Low	Normal	Normal	Notably high

8.2 River flows table

Site name	River	Catchment	November 2024 band	October 2024 band
Hartford Bridge	Blyth	Blyth	Normal	Notably high
Haydon Bridge	South Tyne	Tyne	Notably low	Above normal
Heaton Mill	Till	Till	Notably low	Notably high
Middleton In Teesdale	Tees	Tees	Notably low	Notably high
Mitford	Wansbeck	Wansbeck	No data	No data
Rothbury	Coquet	Coquet	Below normal	Notably high
Rutherford Bridge	Greta	Tees	Below normal	Above normal
Witton Park	Wear	Wear	Below normal	Above normal

8.3 Groundwater table

Site name	Aquifer	End of November 2024 band	End of October 2024 band
Aycliffe Nra2	Skerne Magnesian Limestone	Notably high	Exceptionally high
Red Lion	Skerne Magnesian Limestone	Normal	Normal
Royalty Observ.	Till Fell Sandstone	Above normal	Exceptionally high
Townlaw	Till Fell Sandstone	Normal	Below normal
West Hall Farm	Wear Magnesian Limestone	Exceptionally high	Exceptionally high