

Risk of airborne introduction of BTV and EHDV to Great Britain from the near Continent.

Time period: 11 to 17 December 2024.

This report describes the retrospective risk of entry of bluetongue virus (BTV) or epizootic haemorrhagic disease virus (EHDV)- infected midges into Great Britain (GB) from the near Continent over the previous week. It does not attempt to predict the future risk of virus entry or consider the historical risk earlier than the time period stated above.

We estimate the overall risk of airborne introduction of either infectious BTV-infected or infectious EHDV-infected midges to GB from the near Continent over the last week to have been “Negligible” (see Appendix A for definitions), meaning that the risk was low enough to not merit consideration. We did not consider meteorological conditions to have been suitable for vector incursions from any country over the last week.

We also consider vector activity and temperatures in coastal and near-coastal areas in the south and east of England (the areas of GB most at risk of airborne virus incursions) over the previous two weeks to estimate the potential risk of onward spread of BTV within these areas should an incursion have occurred. This risk only considers the spread risk from infectious vectors entering the country over the previous two weeks and does not consider the risk of spread from vectors infected before this time.

We estimate that the risk of spread of BTV over the last two weeks if incursions did occur was “Negligible” in all four regions (the North East, East Anglia, the South East and the South West). This means that temperatures in all four regions were considered to be consistently unsuitable for sustained and onward BTV transmission by local midge populations as a result of infectious vectors entering the country over the previous two weeks.

Preliminary outbreak assessments of the BTV and EHDV situation in Europe are available, which also consider other potential routes of virus entry.

Read about [Bluetongue virus in Europe \(GOV.UK\)](#)

Read about [Epizootic haemorrhagic disease in Europe \(GOV.UK\)](#)

More details on our risk estimates and the evidence underlying these are provided in the seven tables in the report below. For each of BTV and EHDV, we provide three tables which describe the risk of airborne incursions. These represent:

- Our estimate of the risk of incursion of midge vectors (regardless of infection status) from each high-risk source country (with this risk shown in the top row of the table).
- Our estimate of the risk of incursion of infectious midge vectors from each high-risk source country (with this risk shown in the top row of the table).
- Our estimate of the uncertainty surrounding the country-specific risk of incursion of infectious vectors, accounting for our knowledge of relevant processes and the available data (with this uncertainty shown as the top row of the table).

We also provide a table showing our estimates of the risk of virus spread within high-risk areas of GB (which considers both vector feeding behaviour and temperature suitability for virus replication and spread).

Airborne Orbivirus Assessment
No. 39
18 December 2024



1. Risk of airborne BTV incursion into GB from high-risk countries on the Continent.

	France	Belgium	Netherlands	Germany	Denmark
Risk of vector incursion into GB from source country	Negligible: No midges are expected to have been blown into GB over the previous week.	Negligible: No midges are expected to have been blown into GB over the previous week.	Negligible: No midges are expected to have been blown into GB over the previous week.	Negligible: No midges are expected to have been blown into GB over the previous week.	Negligible: No midges are expected to have been blown into GB over the previous week.
Incursion risk distribution	Not Applicable	Not Applicable	Not Applicable	Not Applicable	Not Applicable

	France	Belgium	Netherlands	Germany	Denmark
Risk of incursion of infectious BTV-infected vectors into GB from source country	Negligible: No infectious BTV-infected midges are expected to have been blown into GB over the previous week.	Negligible: No infectious BTV-infected midges are expected to have been blown into GB over the previous week.	Negligible: No infectious BTV-infected midges are expected to have been blown into GB over the previous week.	Negligible: No infectious BTV-infected midges are expected to have been blown into GB over the previous week.	Negligible: No infectious BTV-infected midges are expected to have been blown into GB over the previous week.
Vector activity in source country	Thought to be active and feeding in source country.	Thought to be active and feeding in source country.	Thought to be active and feeding in source country.	Thought to be active and feeding in source country.	Thought to be active and feeding in source country.
Temperatures in source country	Unsuitable for development of infectiousness in newly infected vectors in the high-risk area over the previous week.	Unsuitable for development of infectiousness in newly infected vectors in the high-risk area over the previous week.	Unsuitable for development of infectiousness in newly infected vectors in the high-risk area over the previous week.	Unsuitable for development of infectiousness in newly infected vectors in the high-risk area over the previous week.	Unsuitable for development of infectiousness in newly infected vectors in the high-risk area over the previous week.
Current BTV status of source country	8,710 BTV-3 outbreaks reported in country between 01 June and 12 December. In the high-risk area, most reports have been in the east but reports have been	3,672 BTV-3 cases reported in country between 01 June and 03 November, including in the high-risk area.	10,621 BTV-3 outbreaks (2,116 clinical and 8,505 PCR-confirmed) reported in country between 01 June and 12 December, including in the high-risk area.	15,372 BTV-3 outbreaks reported in country between 01 June and 17 December, including in the high-risk area.	1,024 reports of BTV-3 in country between 01 June and 17 December, including in the high-risk area.

Airborne Orbivirus Assessment
No. 39
18 December 2024



	made as far west as the Cherbourg peninsula. Country considered endemic for BTV-4 and BTV-8 (but BTV-4 not currently thought to be present in the mainland).		Total of 12 reports of BTV-12 (10 in high-risk area) between 10 October and 12 December.		
Livestock immunity in the source country	Some immunity to BTV-8 expected due to natural infection and vaccination. No immunity to BTV-3 expected before the current outbreak, and no expected immunity to BTV-12. Voluntary BTV-3 vaccination recently implemented for all cattle and sheep in the country.	Some immunity to BTV-3 expected due to natural infection and vaccination. No expected immunity to BTV-12.	Some immunity to BTV-3 expected due to natural infection and vaccination. No expected immunity to BTV-12.	Some immunity to BTV-3 expected due to natural infection and vaccination. No expected immunity to BTV-12.	No immunity to any strain of BTV expected before the current outbreak as no natural spread or vaccination in recent years. Voluntary BTV-3 vaccination recently implemented in the country.

	France	Belgium	Netherlands	Germany	Denmark
Uncertainty in BTV incursion risk estimate	Low: Relevant knowledge good and most/ all data are complete. No subjective judgement is introduced.	Low: Relevant knowledge good and most/ all data are complete. No subjective judgement is introduced.	Low: Relevant knowledge good and most/ all data are complete. No subjective judgement is introduced.	Low: Relevant knowledge good and most/ all data are complete. No subjective judgement is introduced.	Low: Relevant knowledge good and most/ all data are complete. No subjective judgement is introduced.
Knowledge of current livestock infection in high-risk area of source country	Moderate uncertainty. Recent reports of infection with new strain of BTV-8, but this strain has not yet been detected in the high-risk area. The exact number of BTV-8 cases is unknown. A total of 274 new reports of BTV-3 infection in the	Moderate uncertainty. no new reports of BTV infection in the country reported since 03 November, with no regular updates on numbers of infected premises since 07 October. Level of surveillance activities is unknown.	Low uncertainty. A total of 143 new outbreaks of BTV-3 in country reported between 05 and 12 December. BTV-3 vaccine efficacy is unclear. Current level of immunity to BTV-3 is unknown. No	Low uncertainty. A total of 206 new outbreaks of BTV-3 in the country have been reported 10 and 17 December. Vaccine efficacy is unclear. Current level of immunity is also unknown.	Low uncertainty. A total of 7 new outbreaks of BTV-3 in the country have been reported between 09 and 17December.

Airborne Orbivirus Assessment
No. 39
18 December 2024



	country between 06 and 12 December. Most outbreaks of BTV-3 remain near to the Belgian and German borders, although cases have now been reported in the French high-risk area as far west as the western coast of Cherbourg peninsula. Level of surveillance activities is unknown. Current level of immunity also unknown. Vaccine coverage or efficacy remains unclear.	Vaccine coverage has been reported as being low. Vaccine efficacy is unclear. Current level of immunity is also unknown.	expected immunity to BTV-12.		
Knowledge of current vector infection in high-risk area of source country	Moderate uncertainty. Uncertainty in livestock BTV-8 infection status in the high-risk area (although cases of livestock infection with BTV-3 and the new strain of BTV-8 are being reported in the country).	Moderate uncertainty. No update on numbers of infected livestock premises since 03 November.	Low uncertainty. Knowledge of recent livestock infection in high-risk area.	Low uncertainty. Knowledge of recent livestock infection in high-risk area.	Low uncertainty. Knowledge of recent livestock infection in high-risk area.
Other comments on uncertainty	Uncertainty estimate constrained by lack of potential vector incursions.	Uncertainty estimate constrained by lack of potential vector incursions.	None	None	None

Airborne Orbivirus Assessment
No. 39
18 December 2024



2. Risk of airborne EHDV incursion into GB from high-risk countries on the Continent.

	France	Belgium	Netherlands	Germany	Denmark
Risk of vector incursion into GB from source country	Negligible: No midges are expected to have been blown into GB over the previous week.	Negligible: No midges are expected to have been blown into GB over the previous week.	Negligible: No midges are expected to have been blown into GB over the previous week.	Negligible: No midges are expected to have been blown into GB over the previous week.	Negligible: No midges are expected to have been blown into GB over the previous week.
Incursion risk distribution	Not Applicable	Not Applicable	Not Applicable	Not Applicable	Not Applicable

	France	Belgium	Netherlands	Germany	Denmark
Risk of incursion of infectious EHDV-infected vectors into GB from source country	Negligible: No infectious EHDV-infected midges are expected to have been blown into GB over the previous week.	Negligible: No infectious EHDV-infected midges are expected to have been blown into GB over the previous week.	Negligible: No infectious EHDV-infected midges are expected to have been blown into GB over the previous week.	Negligible: No infectious EHDV-infected midges are expected to have been blown into GB over the previous week.	Negligible: No infectious EHDV-infected midges are expected to have been blown into GB over the previous week.
Vector activity in source country	Thought to be active and feeding in source country.	Thought to be active and feeding in source country.	Thought to be active and feeding in source country.	Thought to be active and feeding in source country.	Thought to be active and feeding in source country.
Temperatures in source country	Unsuitable for development of infectiousness in newly infected vectors over the previous week in the high-risk area.	Unsuitable for development of infectiousness in newly infected vectors over the previous week in the high-risk area.	Unsuitable for development of infectiousness in newly infected vectors over the previous week in the high-risk area.	Unsuitable for development of infectiousness in newly infected vectors over the previous week in the high-risk area.	Unsuitable for development of infectiousness in newly infected vectors over the previous week in the high-risk area.
Current EHDV status of source country	3,438 EHDV outbreaks have been reported in the country between 01 June and 12 December. A focus of infection has been present around the Maine-et-Loire	Country considered free of EHDV.	Country considered free of EHDV.	Country considered free of EHDV.	Country considered free of EHDV.

Airborne Orbivirus Assessment
No. 39
18 December 2024



	and Loire-Atlantique departments in the northwest of the country, with cases reported in the adjacent Morbihan and Ile-et-Vilaine departments (with one outbreak reported around 10km from the high-risk area). However, no cases have been reported inside the high-risk area to date.				
Livestock immunity in the source country	Voluntary vaccination against EHDV commenced in a zone near to the border of the 150km EHDV control zone on 23 September. Part of the high-risk area is included in this zone.	No immunity to EHDV expected as no natural spread or vaccination.	No immunity to EHDV expected as no natural spread or vaccination.	No immunity to EHDV expected as no natural spread or vaccination.	No immunity to EHDV expected as no natural spread or vaccination.

	France	Belgium	Netherlands	Germany	Denmark
Uncertainty in EHDV incursion risk estimate	Low: Relevant knowledge good and most/ all data are complete. No subjective judgement is introduced.	Low: Relevant knowledge good and most/ all data are complete. No subjective judgement is introduced.	Low: Relevant knowledge good and most/ all data are complete. No subjective judgement is introduced.	Low: Relevant knowledge good and most/ all data are complete. No subjective judgement is introduced.	Low: Relevant knowledge good and most/ all data are complete. No subjective judgement is introduced.
Knowledge of current livestock infection in high-risk area of source country	Moderate uncertainty. Infection was reported last season. All outside high-risk area but some close to this area. 64 new EHDV outbreaks between 05 and 12 December.	Low uncertainty. No reports and no suspicion of virus presence in country.	Low uncertainty. No reports and no suspicion of virus presence in country.	Low uncertainty. No reports and no suspicion of virus presence in country.	Low uncertainty. No reports and no suspicion of virus presence in country.

Airborne Orbivirus Assessment
No. 39
18 December 2024



	Level of surveillance activities unknown. Level of immunity unknown. Vaccine coverage or efficacy remains unclear.				
Knowledge of current vector infection in high-risk area of source country	Moderate uncertainty. Uncertainty in livestock infection status.	Low uncertainty. No reports and no suspicion of virus presence in country.	Low uncertainty. No reports and no suspicion of virus presence in country.	Low uncertainty. No reports and no suspicion of virus presence in country.	Low uncertainty. No reports and no suspicion of virus presence in country.
Other comments on uncertainty	Uncertainty estimate constrained by lack of potential vector incursions.	None	None	None	None

**Airborne Orbivirus Assessment
No. 39
18 December 2024**



3. BTV spread risk within GB.

High risk region	Counties in region	Estimated number of cattle in region	Estimated number of sheep in region	Vector feeding levels	Temperature suitability for virus spread
South West	Cornwall Devon Dorset Somerset Wiltshire	1,533,622	1,540,170	Vectors are known to have been feeding on livestock over the last two weeks.	Negligible: Temperatures in the high risk region of GB were consistently unsuitable for sustained and onward BTV transmission by local midge populations as a result of infectious vectors entering the country over the previous two weeks.
South East	Hampshire Isle of Wight East Sussex West Sussex Kent	204,573	434,680	Vectors are known to have been feeding on livestock over the last two weeks.	Negligible: Temperatures in the high risk region of GB were consistently unsuitable for sustained and onward BTV transmission by local midge populations as a result of infectious vectors entering the country over the previous two weeks.
East Anglia	Essex Suffolk Norfolk	133,490	145,925	Vectors are known to have been feeding on livestock over the last two weeks.	Negligible: Temperatures in the high risk region of GB were consistently unsuitable for sustained and onward BTV transmission by local midge populations as a result of infectious vectors entering the country over the previous two weeks.
North East	Lincolnshire East Riding of Yorkshire North Yorkshire Durham Tyne & Wear Northumberland	764,208	2,044,607	Vectors are known to have been feeding on livestock over the last two weeks.	Negligible: Temperatures in the high risk region of GB were consistently unsuitable for sustained and onward BTV transmission by local midge populations as a result of infectious vectors entering the country over the previous two weeks.

Estimates of numbers of cattle and sheep in the different regions as shown in the table above are taken from recent LDDG reports:

- [Livestock Demographic Data Group: Cattle population report 2023](#)
- [Livestock Demographic Data Group: Sheep population report 2023](#)

**Airborne Orbivirus Assessment
No. 39
18 December 2024**



Appendix A: Risk and uncertainty levels used in the assessment.

Risk level	Probability range	Likelihood statement
Negligible	Less than 1 in 1 million	So rare that it does not merit to be considered
Very Low	Between 1 in one million and 1 in one thousand	Very unlikely but cannot be excluded
Low	Between 1 in one thousand and 0.05	Unlikely but could occur
Medium	Between 0.05 and 0.50	Likely
High	Between 0.50 and 0.90	Very likely but not certain
Very High	Over 0.90	Almost certain

Uncertainty level	Explanation
Low	Knowledge of the relevant processes is considered good and most/ all information or data are complete. No subjective judgement is introduced.
Moderate	There is a lack of knowledge of the relevant processes and/ or some information or data are lacking or incomplete. Subjective/ expert judgement is introduced with supporting evidence.
High	There is a lack of knowledge of the relevant processes and/ or most information or data are lacking or incomplete. Subjective/ expert judgement may be introduced without supporting evidence.

Appendix B: Modelling overview.

The Met Office Numerical Atmospheric-dispersion Modelling Environment (NAME) is run twice a day to estimate the likely transport of any infected midges.


NAME is run using meteorological data from the Met Office's numerical weather prediction (NWP) model output with 1.5 km horizontal spatial resolution and hourly time resolution. Modelled particles are released over a 2-hour period at sunrise and over a 3-hour period at sunset to represent the diel periodicity of midge activity. The model particles are released from 10 m above ground level. This height is assumed to be above the normal flight boundary layer of midges, where wind speed is greater than midge flight speed, and therefore excludes midges undertaking active local-scale flight. Particles are then dispersed for 12 hours, reflecting wind-tunnel experiments on the flight duration of midges.

Particles are released from nine hypothetical source locations on the near Continent, located from north west France to western Denmark. These nine sites do not represent current observed locations of high midge activity or disease presence, but are used to give broad coverage of the coastline of the near Continent. The particles released are the NAME "midge" species, for which the particle release rate is a function of the day of the year and the local temperature, wind speed and precipitation at the source location. These midge species particles are also removed from the atmosphere either if they encounter rain rates in excess of 1 mm per hour, to represent the washout of midges, or if they pass over land any time after their first 2 hours of flight, to represent the small distances midges travel over land. The NAME midge species represents the most likely scenario for midges active on the Continent, as it takes into account the effects of seasonality and the meteorology on both take-off and survivability along the flight trajectory.

The resulting midge plumes do not represent the spread of disease, rather the spread of midges (which may or may not be infected) had they been present at the source location. The risk of airborne incursion from a particular source into mainland GB coastal counties and the Channel Islands is based upon the total number of incursions by midge plumes from that source over the previous week.

Appendix C: High risk areas in Continental Europe and GB.

We consider the 50km wide area of continental Europe shown in red below to represent the area of main interest with regards to livestock infection and temperature suitability for onwards virus spread. The coloured areas in GB represent ceremonial counties at risk of airborne virus entry and/or potential subsequent onwards spread within GB, grouped into regions as described in the report. Note that although Somerset and Wiltshire are not coastal (and are therefore not considered counties of potential incursion), they are included as counties of interest for onwards spread within GB.



This image shows a map of GB and the adjacent portion of continental Europe. Within GB, four coastal and near-coastal regions in the south and east of England are indicated. These are named as follows (and composed of the listed ceremonial counties): the South West (Cornwall, Devon, Somerset, Dorset, Wiltshire), South East (Hampshire, Isle of Wight, West Sussex, East Sussex, Kent), East Anglia (Essex, Suffolk, Norfolk), and the North East (Lincolnshire, East Riding of Yorkshire, North Yorkshire, Durham, Tyne & Wear, Northumberland). Within continental Europe, a 50km band along the coast adjacent to GB is indicated in red. This runs from western France (Brittany), through north and northwestern Belgium, Netherlands, and Germany, through western Denmark to the north of the Jutland peninsula.