

The complete routine immunisation schedule

From January 2025

Age due	Diseases protected against	Vaccine given and trade name		Usual site ¹
Eight weeks old	Diphtheria, tetanus, pertussis (whooping cough), polio, <i>Haemophilus influenzae</i> type b (Hib) and hepatitis B	DTaP/IPV/Hib/HepB	Infanrix hexa or Vaxelis	Thigh
	Meningococcal group B (MenB)	MenB	Bexsero	Thigh
	Rotavirus gastroenteritis	Rotavirus ²	Rotarix ²	By mouth
Twelve weeks old	Diphtheria, tetanus, pertussis, polio, Hib and hepatitis B	DTaP/IPV/Hib/HepB	Infanrix hexa or Vaxelis	Thigh
	Pneumococcal (13 serotypes)	Pneumococcal conjugate vaccine (PCV)	Prevenar 13	Thigh
	Rotavirus	Rotavirus ²	Rotarix ²	By mouth
Sixteen weeks old	Diphtheria, tetanus, pertussis, polio, Hib and hepatitis B	DTaP/IPV/Hib/HepB	Infanrix hexa or Vaxelis	Thigh
	MenB	MenB	Bexsero	Thigh
One year old (on or after the child's first birthday)	Hib and MenC	Hib/MenC	Menitorix	Upper arm/thigh
	Pneumococcal	PCV booster	Prevenar 13	Upper arm/thigh
	Measles, mumps and rubella (German measles)	MMR	MMRvaxPro ³ or Priorix	Upper arm/thigh
	MenB	MenB booster	Bexsero	Thigh
Eligible paediatric age groups ⁴	Influenza (each year from September)	Live attenuated influenza vaccine LAIV ^{3,6}	Fluenz ^{3,6}	Both nostrils
Three years four months old or soon after	Diphtheria, tetanus, pertussis and polio	dTaP/IPV	REPEVAX	Upper arm
	Measles, mumps and rubella	MMR (check first dose given)	MMRvaxPro ³ or Priorix	Upper arm
Boys and girls aged twelve to thirteen years	Cancers and genital warts caused by specific human papillomavirus (HPV) types	HPV ⁵	Gardasil 9	Upper arm
Fourteen years old (school Year 9)	Tetanus, diphtheria and polio	Td/IPV (check MMR status)	REVAXIS	Upper arm
	Meningococcal groups A, C, W and Y	MenACWY	MenQuadfi	Upper arm
65 years old	Pneumococcal (23 serotypes)	Pneumococcal Polysaccharide Vaccine (PPV23)	Pneumovax 23	Upper arm
65 years of age and older	Influenza (each year from September)	Inactivated influenza vaccine	Multiple	Upper arm
65 from September 2023 ⁷	Shingles	Shingles vaccine	Shingrix	Upper arm
70 to 79 years of age (plus eligible age groups and severely immunosuppressed) ⁷	Shingles	Shingles vaccine	Zostavax ^{3,7} (or Shingrix if Zostavax contraindicated)	Upper arm
	75 years of age	Respiratory syncytial virus (RSV)	RSV vaccine	Abrysvo

1. Intramuscular injection into deltoid muscle in upper arm or anterolateral aspect of the thigh.
 2. Rotavirus vaccine should only be given after checking for SCID screening result.
 3. Contains porcine gelatine.
 4. See annual flu letter at: www.gov.uk/government/collections/annual-flu-programme
 5. See Green Book HPV Chapter 18a for details on immunising immunocompromised young people who will need 3 doses.

6. If LAIV (live attenuated influenza vaccine) is contraindicated or otherwise unsuitable use inactivated flu vaccine (check Green Book Chapter 19 for details).
 7. See Green Book Shingles Chapter 28a for details on eligible age groups including severely immunosuppressed individuals from age 50.



For vaccine supply information for the routine immunisation schedule please visit portal.immform.ukhsa.gov.uk and check Vaccine Update for all other vaccine supply information: www.gov.uk/government/collections/vaccine-update

Selective immunisation programmes

Target group	Age and schedule	Disease	Vaccines required
Babies born to hepatitis B infected mothers	At birth, four weeks and 12 months old ^{1,2}	Hepatitis B	Hepatitis B (Engerix B/HBvaxPRO)
Infants in areas of the country with TB incidence $\geq 40/100,000$	Around 28 days old ⁴	Tuberculosis	BCG
Infants with a parent or grandparent born in a high incidence country ³	Around 28 days old ⁴	Tuberculosis	BCG
Children in a clinical risk group	From 6 months to 17 years of age	Influenza	LAIV or inactivated flu vaccine if contraindicated to LAIV or under 2 years of age
Pregnant women	At any stage of pregnancy during flu season	Influenza	Inactivated flu vaccine
	From 16 weeks gestation ⁵	Pertussis	Tdap (ADACEL)
	From 28 weeks gestation	RSV	RSV vaccine (Abrysvo)

1. Take blood for HBsAg at 12 months to exclude infection.
2. In addition hexavalent vaccine (Infanrix hexa or Vaxelis) is given at 8, 12 and 16 weeks.
3. Where the annual incidence of TB is $\geq 40/100,000$ – see www.gov.uk/government/publications/tuberculosis-tb-by-country-rates-per-100000-people

4. Check SCID screening outcome before giving BCG.
5. Ideally before 32 weeks gestation but may still be given after 32 weeks.

Additional vaccines for individuals with underlying medical conditions

Medical condition	Diseases protected against	Vaccines required ¹
Asplenia or splenic dysfunction (including due to sickle cell and coeliac disease)	Meningococcal groups A, B, C, W and Y Pneumococcal Influenza	MenACWY MenB PCV13 (up to 10 years of age) ² PPV23 (from 2 years of age) Annual flu vaccine
Cochlear implants	Pneumococcal	PCV13 (up to 10 years of age) ² PPV23 (from 2 years of age)
Chronic respiratory and heart conditions (such as severe asthma, chronic pulmonary disease, and heart failure)	Pneumococcal Influenza	PCV13 (up to 10 years of age) ² PPV23 (from 2 years of age) Annual flu vaccine
Chronic neurological conditions (such as Parkinson's or motor neurone disease, or learning disability)	Pneumococcal (only if the individual is also at increased risk of aspiration) Influenza	PCV13 (up to 10 years of age) ² PPV23 (from 2 years of age) Annual flu vaccine
Diabetes	Pneumococcal Influenza	PCV13 (up to 10 years of age) ² PPV23 (from 2 years of age) Annual flu vaccine
Chronic kidney disease (CKD) (including haemodialysis)	Pneumococcal (stage 4 and 5 CKD) Influenza (stage 3, 4 and 5 CKD) Hepatitis B (stage 4 and 5 CKD)	PCV13 (up to 10 years of age) ² PPV23 (from 2 years of age) Annual flu vaccine Hepatitis B
Chronic liver conditions	Pneumococcal Influenza Hepatitis A Hepatitis B	PCV13 (up to 10 years of age) ² PPV23 (from 2 years of age) Annual flu vaccine Hepatitis A Hepatitis B
Haemophilia	Hepatitis A Hepatitis B	Hepatitis A Hepatitis B
Immunosuppression due to disease or treatment ⁴	Pneumococcal Shingles vaccine Influenza	PCV13 (up to 10 years of age) ^{2,3} PPV23 (from 2 years of age) Shingrix – over 50 years of age ⁵ Annual flu vaccine
Complement disorders (including those receiving complement inhibitor therapy)	Meningococcal groups A, B, C, W and Y Pneumococcal Influenza	MenACWY MenB PCV13 (up to 10 years of age) ² PPV23 (from 2 years of age) Annual flu vaccine

1. Check relevant chapter of the Green Book for specific schedule: www.gov.uk/government/collections/immunisation-against-infectious-disease-the-green-book
2. If aged 2 years to under 10 years of age and unimmunised or partially immunised against pneumococcal infection, give one PCV13 dose.
3. To any age in severely immunosuppressed.

4. Consider annual influenza vaccination for household members and those who care for people with these conditions.
5. Check Green Book Shingles Chapter 28a www.gov.uk/government/publications/shingles-herpes-zoster-the-green-book-chapter-28a