

# Monthly water situation report: Solent and South Downs Area

## Summary - August 2025

Solent and South Downs (SSD) had below average rainfall in August, receiving 61% (41mm) of the long term average (LTA) of 66mm. Monthly mean river flows across SSD ranged from notably low to normal. The end of month groundwater levels ranged from exceptionally low to above normal. Soils across SSD ended the month drier than the average for August. End of month reservoir stocks were below average at Ardingly Reservoir (Ouse) and at Arlington Reservoir (Cuckmere).

### 1.1 Rainfall

SSD had below average rainfall in August, receiving 61% (41mm) of LTA (66mm). The Sussex Coast areal unit received the highest rainfall during August with 85% (50mm) of LTA (59.3mm). The Pevensey Levels areal unit received the lowest rainfall with 32% (21mm) of LTA (64.6mm).

Most of August was dry until the 26 August when heavy rainfall occurred. The 28 August was the wettest day with some very high totals due to thunderstorms. The highest daily totals were recorded on 28 August at Westergate (Sussex Coast) and Calbourne (Isle of Wight) received with 59.8mm and 56.9mm respectively.

The last 6 months has been the sixth driest March–August period on record for SSD. It was the fourth driest March-August period on record for the Test Chalk areal unit. Only the March-August periods for 1976, 1921 and 1990 were drier. For the Hampshire Tertiaries areal unit was the fifth driest March-August period.

For the 24-month period (starting September 2023) it was the seventh highest total for SSD.

All these statistics are based on records going back to 1871.

### 1.2 Soil moisture deficit and recharge

Soils across SSD ended the month drier than the LTA for August.

### 1.3 River flows

Monthly mean river flows for August ranged from notably low to normal across SSD.

Flows were notably low on the:

- River Lymington at Brockenhurst
- River Meon at Mislingford

Flows were below normal on the:

- River Arun at Alfoldean
- River Test at Broadlands
- River Test at Chilbolton

Flows were normal on the:

- River Test at Chilbolton
- River Itchen at Allbrook and Highbridge
- River Wallington at North Fareham
- River Medina at Blackwater
- River Rother at Iping Mill
- River Cuckmere at Cowbeech
- River Adur at Sakeham
- River Ouse at Goldbridge

The notably low monthly mean flows for the River Lymington at Brockenhurst were the fifth lowest on record for August since 1960.

## 1.4 Groundwater levels

End of month groundwater levels for August ranged from exceptionally low to above normal across SSD.

Groundwater levels were exceptionally low at:

- Carisbrooke Castle (Isle of Wight)
- Beeding Hill (West Sussex Chalk)
- Chilgrove (West Sussex Chalk)

Groundwater levels were below normal at:

- Harting Common (Western Rother Greensand)
- Catherington (East Hampshire Chalk)

Groundwater levels were normal at:

- Clanville Gate (Test Chalk)
- Lopcombe Corner (Test Chalk)
- West Meon (East Hampshire Chalk)
- Cornish Farm (East Sussex Chalk)
- Houndean Bottom (East Sussex Chalk)

Groundwater levels were above normal at:

- Preston Candover (East Hampshire Chalk)
- Youngwoods Copse (Isle of Wight)

The groundwater level at Carisbrooke Castle (Isle of Wight) is the second lowest August level since 1977. Beeding Hill (West Sussex Chalk) is the third lowest August level in a record that dates back to 1979.

## **1.5 Reservoir stocks**

End of month reservoir stocks were below average for both Ardingly and Arlington Reservoirs. Ardingly Reservoir (Ouse) was at 34.3% of total capacity (LTA 73%) and Arlington Reservoir (Cuckmere) was at 56.1% of total capacity (LTA 62.75%).

## **1.6 Environmental impact**

### **1.6.1 Abstraction licence restriction**

At the start of August there were 22 licence restrictions in force, which increased to 27 licence restrictions by the middle of the month. By the end of the month 26 licence restrictions were in place. These included:

Cessation of abstraction for:

- three licences on the Arun and one licence on the Loxwood Stream (Arun),
- one licences on the Pagham Rife (Sussex Coast),
- three licences on the River Rother and one licence on the River Lod (Western Rother Greensand),
- one on the River Meon (East Hampshire Chalk),
- four on the River Medina and three licences on the Shepherds Chine (Isle of Wight),
- one on the Lymington River and three on Walkford Brook (Lymington),
- one on the River Blackwater (Hampshire Tertiaries).

Restrictions on abstraction rates for:

- one on the River Meon (East Hampshire Chalk),
- one on the Calbourne (Isle of Wight),
- one on the River Blackwater (Hampshire Tertiaries).

### **1.6.2 Flood Warnings in Force**

There were three flood alerts issued in August 2025 associated with the rainfall at the end of the month. The first flood alert was on 28 August in the Ouse catchment. On 29 August flood alerts were issued for the Western Yar on the Isle of Wight and the Aldingbourne and Barnham Rifes in West Sussex.

Author: [HydrologySSD@Environment-agency.gov.uk](mailto:HydrologySSD@Environment-agency.gov.uk)<Enter Authors email address>


All data are provisional and may be subject to revision. The views expressed in this document are not necessarily those of the Environment Agency. Its officers, servants or agents accept

no liability for any loss or damage arising from the interpretation or use of the information, or reliance upon views contained in this report.

## 2. Rainfall

### 2.1 Rainfall map

Figure 2.1: Total rainfall for hydrological areas for the current month (up to 31 August 2025), the last 3 months, the last 6 months, and the last 12 months, classed relative to an analysis of respective historic totals. Table available in the appendices with detailed information.




Rainfall data for October 2023 onwards, extracted from Environment Agency 1km gridded rainfall dataset derived from Environment Agency intensity rain gauges. (Source: Environment Agency. Crown Copyright, 100024198, 2025). Rainfall data prior to October 2023, extracted from Met Office HadUK 1km gridded rainfall dataset derived from registered rain gauges (Source: Met Office. Crown copyright, 2025).

## 3 River flows and Groundwater levels

### 3.1 River flows and Groundwater level map

Figure 3.1: Monthly mean river flow and groundwater levels at our indicator sites for August 2025, expressed as a percentage of the respective long term average and classed relative to an analysis of historic August monthly means. Table available in the appendices with detailed information.




(Source: Environment Agency). Crown copyright. All rights reserved. Environment Agency, 100024198, 2025. Geological map reproduced with kind permission from UK Groundwater Forum, BGS copyright NERC. Crown copyright. All rights reserved. Environment Agency, 100024198, 2025.

# 4 West Hampshire

## 4.1 West Hampshire Rainfall and effective rainfall charts


Figure 4.1: Monthly rainfall and effective rainfall totals for the past 12 months compared to the 1991 to 2020 long term average.



HadUK rainfall data. (Source: Met Office. Crown copyright, 2025).

## 4.2 West Hampshire River flow charts





Figure 4.2: Daily mean river flow for index sites over the past year, compared to an analysis of historic daily mean flows, and long term maximum and minimum flows.



Source: Environment Agency, 2025.

### 4.3 West Hampshire Groundwater level charts

Figure 4.3: End of month groundwater levels at index groundwater level sites for major aquifers. 34 months compared to an analysis of historic end of month levels and long term maximum and minimum levels.




Source: Environment Agency, 2025.

# 5 East Hampshire and Isle of Wight

## 5.1 East Hampshire and Isle of Wight Rainfall and Effective rainfall charts


Figure 5.1: Monthly rainfall and effective rainfall totals for the past 12 months compared to the 1991 to 2020 long term average.



HadUK rainfall data. (Source: Met Office. Crown copyright, 2025).

## 5.2 East Hampshire and Isle of Wight River flow charts



Figure 5.2 Daily mean river flow for index sites over the past year, compared to an analysis of historic daily mean flows, and long term maximum and minimum flows.



Source: Environment Agency, 2025.

### 5.3 East Hampshire and Isle of Wight Groundwater level charts

Figure 5.3: End of month groundwater levels at index groundwater level sites for major aquifers. 34 months compared to an analysis of historic end of month levels and long term maximum and minimum levels.




Source: Environment Agency, 2025.

# 6 West Sussex


## 6.1 West Sussex Rainfall and Effective Rainfall charts

Figure 6.1: Monthly rainfall and effective rainfall totals for the past 12 months as a percentage of the 1991 to 2020 long term average.



## 6.2 West Sussex River flow charts


Figure 6.2: Daily mean river flow for index sites over the past year, compared to an analysis of historic daily mean flows, and long term maximum and minimum flows.



Source: Environment Agency, 2025.

6.3 West Sussex Groundwater level charts

Figure 6.3: End of month groundwater levels at index groundwater level sites for major aquifers. 34 months compared to an analysis of historic end of month levels and long term maximum and minimum levels.




Source: Environment Agency, 2025.

# 7 East Sussex




## 7.1 East Sussex Rainfall and Effective Rainfall charts

Figure 7.1: Monthly rainfall and effective rainfall totals for the past 12 months compared to the 1991 to 2020 long term average.



7.2 East Sussex River flow charts


Figure 7.2: Daily mean river flow for index sites over the past year, compared to an analysis of historic daily mean flows, and long term maximum and minimum flows.



Source: Environment Agency, 2025.

7.3 East Sussex Groundwater level charts


Figure 7.3: End of month groundwater levels at index groundwater level sites for major aquifers. 34 months compared to an analysis of historic end of month levels and long term maximum and minimum levels.



Source: Environment Agency, 2025.

# 8 Reservoir stocks

Figure 8.1: End of month reservoir stocks compared to long term maximum, minimum and average stocks. Note: Historic records of individual reservoirs and reservoir groups making up the regional values vary in length.



(Source: water companies).

## 9 Glossary

### 9.1 Terminology

#### **Aquifer**

A geological formation able to store and transmit water.

#### **Areal average rainfall**

The estimated average depth of rainfall over a defined area. Expressed in depth of water (mm).

#### **Artesian**

The condition where the groundwater level is above ground surface but is prevented from rising to this level by an overlying continuous low permeability layer, such as clay.

#### **Artesian borehole**

Borehole where the level of groundwater is above the top of the borehole and groundwater flows out of the borehole when unsealed.

#### **Cumecs**

Cubic metres per second ( $\text{m}^3\text{s}^{-1}$ ).

#### **Effective rainfall**

The rainfall available to percolate into the soil or produce river flow. Expressed in depth of water (mm).

#### **Flood alert and flood warning**

Three levels of warnings may be issued by the Environment Agency. Flood alerts indicate flooding is possible. Flood warnings indicate flooding is expected. Severe flood warnings indicate severe flooding.

#### **Groundwater**

The water found in an aquifer.

## **Long term average (LTA)**

The arithmetic mean calculated from the historic record, usually based on the period 1991 to 2020. However, the period used may vary by parameter being reported on (see figure captions for details).

## **mAOD**

Metres above ordnance datum (mean sea level at Newlyn Cornwall).

## **MORECS**

Met Office Rainfall and Evaporation Calculation System. Met Office service providing real time calculation of evapotranspiration, soil moisture deficit and effective rainfall on a 40 by 40 km grid.

## **Naturalised flow**

River flow with the impacts of artificial influences removed. Artificial influences may include abstractions, discharges, transfers, augmentation and impoundments.

## **NCIC**

National Climate Information Centre. NCIC area monthly rainfall totals are derived using the Met Office 5 km gridded dataset, which uses rain gauge observations.

## **Recharge**

The process of increasing the water stored in the saturated zone of an aquifer. Expressed in depth of water (mm).

## **Reservoir gross capacity**

The total capacity of a reservoir.

## **Reservoir live capacity**

The capacity of the reservoir that is normally usable for storage to meet established reservoir operating requirements. This excludes any capacity not available for use (for example, storage held back for emergency services, operating agreements or physical restrictions). May also be referred to as 'net' or 'deployable' capacity.

## **Soil moisture deficit (SMD)**

The difference between the amount of water actually in the soil and the amount of water the soil can hold. Expressed in depth of water (mm).

## 9.2 Categories

### **Exceptionally high**

Value likely to fall within this band 5% of the time.

### **Notably high**

Value likely to fall within this band 8% of the time.

### **Above normal**

Value likely to fall within this band 15% of the time.

### **Normal**

Value likely to fall within this band 44% of the time.

### **Below normal**

Value likely to fall within this band 15% of the time.

### **Notably low**

Value likely to fall within this band 8% of the time.

### **Exceptionally low**

Value likely to fall within this band 5% of the time.

## 10 Appendices

### 10.1 Rainfall, effective rainfall and soil moisture deficit table

(Source: Met Office. Crown copyright, 2025). All rights reserved. Environment Agency, 100024198, 2025

Figure 10.1: This is areal rainfall, effective rainfall (percolation or runoff) and soil moisture deficit for the hydrological areas across the SSD. There may be significant variation within each area which must be considered when interpreting these data. When additional meteorological data is available estimates are revised which will affect the period totals in section 10.2

Hydrological Area	Rainfall (mm) 31 day Total	Rainfall August as %LTA	Effective Rainfall (mm) 31 day Total	Effective Rainfall August as %LTA	Soil Moisture Deficit (SMD) Day 31	SMD End of August LTA
Test Chalk	39	61%	4	60%	167	100
East Hampshire Chalk	40	56%	4	52%	154	95
West Sussex Chalk	53	72%	6	77%	137	92
East Sussex Chalk	27	40%	2	35%	153	98
Isle of Wight	50	84%	6	97%	145	109
Western Rother Greensand	50	69%	5	68%	143	95
Hampshire Tertiaries	42	64%	0	0	156	101
Lymington	48	76%	0	0	145	96
Sussex Coast	50	85%	0	0	148	109
Arun	47	73%	0	0	141	95
Adur	46	70%	0	0	135	92
Ouse	32	47%	0	0	139	88
Cuckmere	26	37%	0	0	145	87
Pevensey Levels	21	32%	0	0	153	95
SSD Average	41	61%	2	64%	147	97

## 10.2 Seasonal summary table of rainfall and effective rainfall

Summer season: 01/04/2025 to 31/08/2025

Hydrological Area	Seasonal Rainfall (mm)  Total	Seasonal Rainfall as % LTA	Seasonal Effective Rainfall (mm)  Total	Seasonal Effective Rainfall as % LTA
Test Chalk	156	53%	10	23%
East Hampshire Chalk	194	63%	16	29%
West Sussex Chalk	208	68%	19	35%
East Sussex Chalk	185	66%	16	38%
Isle of Wight	186	72%	16	44%
Western Rother Greensand	208	67%	18	33%
Hampshire Tertiaries	176	63%	0	0%
Lymington	183	66%	0	0%
Sussex Coast	172	68%	0	0%
Arun	193	69%	0	0%
Adur	196	71%	0	0%
Ouse	206	72%	0	0%
Cuckmere	189	68%	0	0%
Pevensey Levels	177	66%	0	0%
SSD Average	188	66%	7	21%

### 10.3 Rainfall banding table

Hydrological area	August 2025 band	June 2025 to August 2025 cumulative band	March 2025 to August 2025 cumulative band	September 2024 to August 2025 cumulative band
Test Chalk	Below normal	Normal	Exceptionally low	Normal
East Hampshire Chalk	Below normal	Normal	Exceptionally low	Normal
West Sussex Chalk	Normal	Normal	Notably low	Normal
East Sussex Chalk	Below normal	Normal	Exceptionally low	Normal
Isle of Wight	Normal	Below normal	Notably low	Normal
Western Rother Greensand	Normal	Below normal	Notably low	Normal
Hampshire Tertiaries	Below normal	Normal	Exceptionally low	Normal
Lymington	Normal	Normal	Exceptionally low	Normal
Sussex Coast	Normal	Below normal	Exceptionally low	Normal
Arun	Normal	Below normal	Notably low	Normal
Adur	Normal	Below normal	Notably low	Normal
Ouse	Below normal	Below normal	Notably low	Normal
Cuckmere	Notably low	Normal	Exceptionally low	Normal
Pevensey Levels	Notably low	Normal	Exceptionally low	Normal

## 10.4 River flows table

Site name	River	Catchment	August 2025 band	July 2025 band
Alfoldean Gs	Arun	Arun	Below normal	Below normal
Allbrook Gs+ Highbridge	Itchen (so)	Itchen	Normal	Normal
Blackwater	Medina	Isle of Wight	Normal	Normal
Broadlands	Test	Test Lower	Below normal	Normal
Brockenhurst GS	Lymington	New Forest	Notably low	Notably low
Chilbolton GS	Test	Test Upper	Below normal	Normal
Cowbeech Gs	Cuckmere	Cuckmere	Normal	Normal
Goldbridge Gs	Ouse [so]	Ouse Sussex	Normal	Above normal
Iping Mill Gs	Rother	West Rother	Normal	Normal
Mislingford GS	Meon	Meon	Notably low	Below normal
North Fareham GS	Wallington	Wallington	Normal	Normal
Sakeham GS	Adur	Adur	Normal	Above normal

## 10.5 Groundwater table

Site name	Aquifer	End of August 2025 band	End of July 2025 band
Carisbrooke Castle	Isle Of Wight Central Downs Chalk	Exceptionally low	Exceptionally low
Youngwoods Copse	Isle of Wight Lower Greensand	Above normal	Above normal
Clanville Gate Gwl	River Test Chalk	Normal	Normal
Lopcombe Corner Gwl	River Test Chalk	Normal	Normal
Preston Candover	River Itchen Chalk	Above normal	Above normal
West Meon Hut Gwl	River Itchen Chalk	Normal	Normal
Catherington	River Meon Chalk	Below normal	Below normal
Chilgrove House Gwl	Chichester-Worthing-Portsdown Chalk	Exceptionally low	Below normal
Beeding Hill Gwl	Brighton Chalk Block	Exceptionally low	Exceptionally low
Houndean Bottom Gwl	Brighton Chalk Block	Normal	Below normal
Harting Common Down	Western Rother Lower Greensand	Below normal	Below normal
Cornish Wigdens Gwtr	Eastbourne Chalk Block	Normal	Normal

## 10.6 Abstraction licence flow constraints

Number of flow constraints in force between 1 to 5 August 2025	Number of flow constraints in force between 6 to 12 August 2025	Number of flow constraints in force between 13 to 19 August 2025	Number of flow constraints in force between 20 to 31 August 2025
22	24	27	26

## 10.7 Solent and South Downs Areal Rainfall Units Map



## 10.8 SSD Areal Rainfall Monthly Long Term Averages

Hydrological Area	Jan LTA mm	Feb LTA mm	Mar LTA mm	Apr LTA mm	May LTA mm	Jun LTA mm	Jul LTA mm	Aug LTA mm	Sep LTA mm	Oct LTA mm	Nov LTA mm	Dec LTA mm
Test Chalk	92.6	65.7	59.4	60.5	53.7	57.8	57.5	63.8	62.8	95.3	100.9	93.1
East Hampshire Chalk	102.7	72.1	62.7	63.5	54.4	62.1	60.5	71.4	70.2	105.8	112.8	105.3
West Sussex Chalk	104.6	73.7	59.5	59.1	54.4	59.4	60.9	73.0	74.4	108.8	114.8	113.2
East Sussex Chalk	96.4	66.9	53.8	49.9	51.2	55.6	57.3	67.6	65.3	101.2	110.6	106.9
Isle of Wight	93.5	66.9	55.4	52.6	47.9	50.6	47.8	59.7	65.9	102.2	107.4	105.6
Western Rother Greensand	110.6	77.5	61.9	64.1	56.4	59.6	57.9	73.1	73.7	111.7	118.1	115.1
Hampshire Tertiaries	95.4	66.7	58.1	57.8	49.8	56.3	51.2	64.7	65.2	97.3	105.1	99.4
Lymington	96.7	67.4	59.6	58.2	50.1	54.5	51.2	62.9	66.6	100.1	109.5	105.3
Sussex Coast	86.8	59.1	48.5	49.6	45.3	50.6	48.8	59.3	59.9	89.3	95.3	93.3
Arun	96.8	67.1	54.7	55.3	53.2	53.3	54.7	64.6	65.8	97.7	103.4	98.9
Adur	94.7	65.6	52.7	52.8	53.5	52.5	53.8	65.1	63.5	99.4	102.9	100.2
Ouse	97.7	67.7	53.9	51.2	53.4	54.9	57.5	67.4	64.7	104.6	105.9	105.4
Cuckmere	94.4	65.8	51.0	49.0	50.5	55.8	56.8	68.7	65.4	101.9	106.8	105.5
Pevensey Levels	89.5	62.7	49.5	47.6	48.1	51.8	56.3	64.6	62.9	96.3	103.0	102.9
<b>SSD Average</b>	96.8	67.7	56.9	56.5	52.0	56.0	55.3	66.0	66.2	100.5	106.6	102.4