



Animal &
Plant Health
Agency

Great Britain and Northern Ireland Variety List Trials: Trial Procedures for Official Examination of Value for Cultivation and Use (VCU) Harvest 2026

Linseed – Spring and Winter

September 2025

Changes since last version

- Updated year of document and date of last update
- Updated references of NL to VL
- Updates to Appendices links

Contents

Section A – Summary of VCU trial assessments required	1
Section B – Seed handling procedures	3
B.1 Seed handling procedures.....	3
B.2 Authentication of VCU seed	3
Section C – Growing trial procedures.....	4
C.1 Responsibilities	4
C.2 Site suitability	4
C.3 Sowing the trial	4
C.4 Husbandry.....	7
C.5 Harvesting	9
C.6 Records.....	11
Section D – Disease testing procedures	15
D.1 Assessment of natural infection	15
D.2 Naturally occurring disease in VCU growing trials.....	15
D.3 Inoculated disease tests	16
Section E – Quality testing procedures	17
E.1 Responsibilities	17
E.2 Quality assessment methodology	17
Section F – Trial design and data handling procedures	18
F.1 Plan validation and storage	18
F.2 Data recording.....	18
F.3 Data processing	18
F.4 Other tests and trials	18
Supporting Document for Appendices.....	19

Section A – Summary of VCU trial assessments required

Bold = Obligatory

Italics = Additional. Assessed only if requested by the applicant

Winter linseed

Type of character	Reference	Description of assessment – Obligatory	Description of assessment – Additional (Assessed only if requested by applicant.)
Yield	Section C	Plot yield	
Yield	Section C	Moisture content	
Impact of environment	Section C	Standing ability	<i>Plant height</i> <i>Earliness of flowering</i> <i>Maturity</i>
Resistance to harmful organisms	Section D	None routinely recorded	
Quality characteristics (laboratory tests)	Section E		<i>Oil content</i>

Spring Linseed

Type of character	Reference	Description of assessment – Obligatory	Description of assessment – Additional (Assessed only if requested by applicant.)
Yield	Section C	Plot yield	
Yield	Section C	Moisture content	
Impact of environment	Section C	Standing ability	<i>Plant height</i> <i>Earliness of flowering</i> <i>Maturity</i>
Resistance to harmful organisms	Section D	None routinely recorded	
Quality characteristics (laboratory tests)	Section E	Oil content	

Further measurements

The following must be measured or recorded in all trials, following procedures in Section C:

Sowing date

Harvest date

Plot size

Bird damage

Seed or boll loss

Combine harvester losses

Section B – Seed handling procedures

B.1 Seed handling procedures

B.1.1 See GENERAL INFORMATION, SECTION 5 - Minor Crop VCU Procedures Introduction.

B.2 Authentication of VCU seed

B.2.1 APHA will notify the Seed Handling Operator of the DUS Test Centre to which a 200 g sample of each variety of linseed should be sent for authentication.

Section C – Growing trial procedures

C.1 Responsibilities

C.1.1 The Growing Trial Operators are responsible for conducting the trials according to these procedures.

C.2 Site suitability

C.2.1 The Growing Trial Operator will be responsible for providing a suitable site, which meets the following criteria:

C.2.2 The trial must be located within a commercial crop to aid management and for spring linseed reduce the risk of flax flea-beetle (FFB) damage. The Trials Organiser should be consulted if this proves impossible or impractical. Previous cropping must be appropriate for a linseed crop to be grown and should have no history of Fusarium wilt or likely herbicide residues that could damage the crop. There should be at least a 3-year (and preferably 5 year) gap between linseed and any other crop susceptible to sclerotinia.

C.2.3 Soil type should be typical of those on which linseed is grown locally. Soil fertility and texture should be uniform across the site. The soil should be as uniform as possible, with no substantial variations in previous cropping, ridges, furrows, etc.

C.2.4 The trial should be sited away from trees, hedges, headlands, and other features, which are likely to cause uneven growth or encourage grazing damage from wild fauna.

C.2.5 The trial area should be cultivated in the direction of ploughing and drilled across the direction of ploughing and cultivation such that each plot receives similar wheeling compaction. Cultivations should follow best local practice.

C.3 Sowing the trial

Time of sowing is critical for rapid emergence and even establishment and to reduce the risk of damage by FFB. As a guideline, winter linseed trials should be drilled between Mid-September and Mid-October, and spring linseed trials should be drilled between the last week of March and the 3rd week of April when soil temperatures reach 8°C and conditions are conducive to rapid and even establishment. To reduce FFB losses in spring linseed the drilling of the trial should coincide as closely as possible with that of the host crop of linseed. Seedbeds need to be well prepared but avoid excessive passes, over-consolidation and compacted soil. Prepare and compress the linseed seedbed so that moisture levels are preserved and even (especially on light soils). The trial can then be drilled when conditions are optimum.

Rolling after drilling is usually necessary and beneficial on lighter and stony soils. Heavier soils should be rolled if there is a risk of moisture loss, but it is essential to avoid capping.

Trial Managers must check the emerging crop regularly and, if, necessary, spray for FFB.

C.3.1 Plot size

C.3.1.1 The harvested plot area per variety should be not less than 20 m² per replicate for trials with a minimum of four replications. Plots should be drilled to a greater length than required and cut back to the required length prior to harvest. The plot width for calculating harvested area is measured centre gap to centre gap with an inter-plot gap in the range 0.5 m to 0.8 m.

C.3.2 Plant population

C.3.2.1 Winter and spring linseed is sown at a seed rate of 600 seeds per square metre (aiming for plant population of 400 seeds per metre).

The following formula will be used to calculate the seed rate for a given thousand seed weight:

$$\text{Seed rate (kg/ha)} = \frac{((\text{Target population} \times \text{Thousand seed weight}) \times 100)}{(\text{Establishment \%} \times \text{Germination \%})}$$

For operators using seed counters the following formula can be used to calculate required seed numbers per plot:

$$\text{Seed rate (kg/ha)} = \frac{((\text{Target population} \times \text{Drilled plot area}) \times 10,000)}{(\text{Establishment \%} \times \text{Germination \%})}$$

The likely establishment should be judged carefully depending on soil conditions and seedbeds. Growing Trial Operators are responsible for achieving the correct target populations.

C.3.3 Trial seed

Untreated seed must be sent as set out in accordance with the Seed and Fee Notice, directly to the Seed Handling Operator by the deadline set out in Appendix 3.

When drilling, every effort should be made to obtain even emergence.

C.3.4 Trial layout

C.3.4.1 The Trials Organiser following consultation with APHA produces provisional sowing lists. The Trials Organiser will make final sowing lists available to Growing Trial Operators, along with the trial plans produced by the Trial Design and Data Handling Operator.

C.3.4.2 The trial should be sown according to the plan produced by the Trial Design and Data Handling Operator and may be an incomplete block design. In an incomplete block design, each replicate is split into a number of sub-blocks. Any splitting of replicates must

be between sub-blocks and not through sub-blocks. Varieties can be moved within a sub-block but must not be moved from their sub-block. Varieties must not be moved around within the plan e.g. if drilling errors occur. If plots are moved out of their original sub-block they will have to be treated as missing plots. If there are any queries, please contact the Trials Organiser.

C.3.4.3 If there is a need to replace a planned variety e.g. if varieties are withdrawn, affected plots must be sown with any of the standard control varieties. Any such replacements must be agreed with the Trials Organiser. The control varieties are listed in Appendix 5.

C.3.5 Drilling

C.3.5.1 Care must be taken with drill settings and drilling speed to ensure satisfactory and uniform establishment and plant population from plot to plot.

Drill at 1.5 - 4cm into moisture in a firm and fine seedbed. In spring linseed, due to the very high risk of damage by flea beetles trial managers are advised to wait until conditions are conducive to good germination and rapid growth. It is also important to ensure that there is no carryover of seed between plots.

C.3.5.2 At least one discard plot must be drilled on either side of the trial with the same drill and at the same time that the trial is drilled.

C.3.5.3 Precautions must be taken to avoid any missing rows. Any missing rows or parts of rows must be noted in the trial diary and reported to the Trials Organiser within one month of emergence.

C.3.6 Confirmation of trial layout

C.3.6.1 After full establishment and within two months of sowing (autumn sown trials) or one month of sowing (spring sown trials), the Growing Trial Operator must confirm that the trial has been sown to plan or give details of any changes to plan. This should be done by clearly highlighting the changes in the electronic plan and returning it to the Trial Design and Data Handling Operator.

Return a completed site data 1 sheet including the following information:

- Site location details including how to get to the field.
- Sketch showing the layout of the trial in the field, in relation to other trials and showing access roads, gates, etc. The location of the access gates should utilise the navigation platform What3Words.com
- Trial sketch showing plot numbers and variety codes/and or names.
- A short post-establishment report of the condition of the trial.

C.4 Husbandry

C.4.1 Agronomy

Where not specified in these procedures' agronomy should follow best local practice, advisory and regulatory guidelines. Application of fertilisers and sprays should be uniform. It is normally best to apply these across the direction of the plots. Application wheelings should not run through the harvested plot area.

C.4.2 Fertiliser application

It should take into account inherent fertility, previous cropping, winter rainfall, the best local practice. All fertiliser applications should take account of AHDB Nutrient Management Guide (RB209), the corresponding advisory publications in England, Wales, Scotland and Northern Ireland and past trialling experience.

A typical rate of nitrogen is to use 80-100 kg/ha minimum (including SMN) as follows:

- Heavier soils: 100% in the prepared seedbed or when rows are first visible.
- Lighter soils: 60% when the rows are visible and 40% at early stem extension.

If the risk of the seedbed drying out is high, then it may be appropriate to apply all N to the seedbed even on lighter soils.

Late applications should be avoided as these can delay harvesting.

Trial managers should be aware of other nutrient requirements and should be prepared, if necessary, to apply appropriate treatments.

C.4.3 Herbicides

The herbicides to be used must be discussed with the Trials Organiser.

Chemicals should not be used to which any variety is known to be sensitive. Pre-emergence herbicides should be used, and it should be noted that under certain soil and weather conditions the linseed crop can be intolerant of some approved post-emergence herbicide products. Post-emergence sulfonyl urea products can be damaging and should be avoided.

The following factors should be considered:

Approved pre-emergence herbicides are effective with good (moist) seedbed conditions and with the appropriate application technique (e.g. water volume). Approved post-emergence herbicides (e.g. Bentazone – Basagran) can be effective against annual dicotyledons.

Post-emergence herbicide applications should be made when all varieties are taller than 15 - 20 cm (but always check the product label for any product variations on this). The risk is greater on light soils (e.g. chalks) and no variety should not be shorter than 10 cm.

Most damage is likely when soils are very dry and/or during extremes of temperature especially very hot conditions.

Experience has shown that the use of Jubilee (Metsulfuron-methyl) can be particularly damaging and should not be used without consulting AHDB/BSPB. Also note that the following products can lead to damage and should be **avoided**:

- Metazachlor e.g. Butisan
- Metazachlor + quimerac e.g. Katamaran
- Napropamide e.g. Devrinol
- Clopyralid + picloram e.g. Galera
- Bifenox e.g. Fox

If use is considered necessary for the success of the trial pay particular attention to restrictions and specific recommendations for use on linseed.

Use the minimum dose of herbicide that will kill the weeds.

C.4.4 Growth regulators

Plant growth regulators should not be used on spring linseed trials.

Up to four (two autumn and two spring) applications of plant growth regulator can be used on winter linseed to manage excessive or rapid growth. All applications should be discussed with the Trials Organiser.

If there is a high risk of severe lodging, applications of plant growth regulator can be used on winter linseed to manage excessive or rapid growth. All applications should be discussed with the Trials Organiser.

C.4.5 Pest and disease control

C.4.5.1 Pest control

Adequate measures should be taken to prevent or minimise damage by any pest. In spring linseed FFB, in particular, are likely to be a significant pest during establishment, in winter linseed pigeons can be a particular problem. Trial managers must ensure that adequate pre- and/or post-emergence control measures are taken.

Assessments should be made wherever pest damage occurs since decisions have to be made on the validity of each plot affected.

For seed dressings, see Appendix 2.

C.4.5.2 Disease control

Precautions should be taken to prevent disease levels in excess of about 10% leaf area cover, or about 10% of capsules infected, by applying appropriate fungicides according to the available approvals and label recommendations. Any disease which does develop should be recorded as described in Section E. In spring linseed, the diseases which are most likely to be encountered are *Botrytis* spp. and *Alternaria* spp. in wet seasons, and Powdery Mildew in dry seasons. In winter linseed *Mycosphaerella* (Pasma) is important.

C.4.6 Irrigation

If irrigation is required to establish the trial, seek the specific agreement of the Trials Organiser.

C.4.7 Pathways

Internal pathways should be made after the risk of pigeon damage has passed.

C.5 Harvesting

C.5.1 Timing of harvesting

It is the Trial Manager's responsibility to ensure that plots can be harvested without damaging neighbouring plots and without contamination: plots should be separated adequately as required.

C.5.2 Harvesting method:

C.5.2.1 Trials may be desiccated prior to combining unless there is a reason for not doing so, the control varieties must be at an overall suitable stage of development.

C.5.2.2 Plots should be trimmed to their final length prior to harvesting. The plot dimensions must be measured prior to harvesting. If it is necessary to reduce the size of any plot at harvest give clear details on the yield file. Individual harvested plot lengths should be recorded.

C.5.3 Samples

C.5.3.1 Samples are required from all plots for moisture determination using the oven method and oil content determination. If additional samples are required, they will be notified to the Growing Trial Operator by the Trials Organiser. All samples should be labelled with the labels provided, giving variety name/breeders reference, AFP number, replicate number and Growing Trial Operator identification number.

C.5.3.2 It is essential that all samples:

- Are representative of the variety/plot from which they are taken with minimal contamination. When sampling on-combine, it is essential to minimise the risk of contamination of grain from the previous plot.
- Are taken from the same source.
- Contain the weight of grain requested.

C.5.3.3 A single sample of 200 g sample should be taken in a polythene bag for moisture content and oil content determination. One label should be placed inside the bag, and this sealed by rolling over the top and securing the bags and the second labels with rubber bands.

C.5.3.4 All bagged samples must be kept in good condition at a moisture content and temperature appropriate for long term storage. They should be clearly marked both inside and outside the container/bag.

C.5.3.5 Sample drying should be undertaken using a cold/warm air drier to reduce moisture content to 9% or below.

C.5.3.6 All plot samples must be labelled with the trial identification number, variety name/breeders' reference, AFP number, plot number and Growing Trial Operator identification number.

C.5.4 Submission of data and samples

C.5.4.1 Appendix 6 lists the records, with deadlines, to be sent to the Trials Organiser. Diary sheets and any other field records should be returned to the Trials Organiser within 5 working days of harvest.

C.5.4.2 All plot records should be transmitted to the Trial Design and Data Handling Operator following the deadlines set out in Appendix 6. The Growing Trial Operator should ensure that data are free from errors before transmission. After scrutiny, copies of results will be returned to the Growing Trial Operator for action as agreed by the Trials Organiser.

C.5.4.3 All samples should be sent to the appropriate Quality Testing Operator following the deadlines set out in Appendix 6.

C.6 Records

6.1 There are four components:

1. **Diary** Field notes of trial status.
2. **Site data part 1*** Including full location details:
 - a) map of site location showing nearby settlements and roads,
 - b) a sketch showing the layout of trials in the field with access points, and
 - c) trial layout, showing plot numbers and variety codes/names.
3. **Site data part 2*** Details of agrochemical applications and irrigation.
4. **Plot records** Plot data.

* Template available from Trials Organiser

C.6.1.1 An entry in the Diary sheet should be made on every trials visit and any observations relevant to variety performance should be recorded. If the trial is in good condition, with no problems, this should be recorded.

C.6.2 Plot records

C.6.2.1 Plot data may be recorded direct onto a data logger using a system approved by the Trials Organiser or recorded on paper then entered and validated onto a computer using an approved system. A system of ensuring that data are recoverable, in the event of loss of original data, must be implemented, e.g. copy and safe storage. Whichever method is used, individual plot data will only be accepted by the appropriate Trial Design and Data Handling Operator in an approved format using the AFP number, variety name and units as listed in Sections C and D.

C.6.2.2 All observations should be checked at the time of recording to identify any unusual plot performance. These observations should be noted by the recorder and any possible causes identified, together with a recommendation for whether the data should remain in the analysis or should be excluded.

C.6.2.3 Plot numbers on record sheets must correspond with the numbering on the field plan.

C.6.2.4 If a character is not recorded or is missing the Growing Trial Operator should indicate in the diary or on the recording sheet the reason why it has been excluded.

C.6.2.5 Where a plot record is missing the Growing Trial Operator should record this in any data file or hard copy medium as a symbol thereby indicating there is no recorded value associated with this plot.

C.6.2.6 Specific plot records should be made as counts or on the scales shown for each character. Only the character names as listed may be used. All records should be returned

to the Trial Design and Data Handling Operator as soon as possible after they are completed to trials@ahdb.org.uk.

C.6.2.7 All records must be returned as soon as reasonably possible and when complete for the whole trial. Indicative deadlines are given in Appendix 6. All records must be returned by the final deadlines.

C.6.3 Procedures for recording characters

The following procedures must be followed for measuring all characters to be used in VL decision-making:

C.6.3.1 **PLOT YIELD AND MOISTURE CONTENT (OBLIGATORY) (kg)**

The following information must accompany the yield data:

- The moisture content % of the harvested grain, determined by oven method.
- Plot length: the plot length harvested in metres.
- Plot width: the width of the harvested plot in metres from outer row to outer row plus half of the inter-plot gap on either side. The adjustment for the inter-plot gap should be no greater than 0.8 m. If these are not the same for every plot a separate record must be submitted.
- Growth stage: usually 9.9 at harvest. The Growth Stage Chart for linseed is at Appendix 7.
- Yield (in kilograms): Note clearly any tare weight to be subtracted.

Yield, moisture content, plot length, plot width and harvest date should be sent to the appropriate Trial Design and Data Handling Operator within 5 days of harvesting the trial.

C.6.3.2 **STANDING ABILITY from all plots**

(Winter linseed - OBLIGATORY)

(1-9) *(Spring linseed - ADDITIONAL)*
(1-9)

1 very poor
9 very good

Growing Trials Operators should assess standing ability at a stage that provides good discrimination between varieties and be prepared to repeat the assessment if further lodging develops.

C.6.3.3 **PLANT HEIGHT from all plots (ADDITIONAL) (cm)**

Record average plot height at the end of flowering before leaning or lodging takes place (if practical take 3 measurements along the length of the plot). If lodging has occurred, choose a representative area of the plot, lift a number of plants against the measuring pole and record an average height.

C.6.3.4 *EARLINESS from all plots* (ADDITIONAL) (1-9)

- 1 very late
- 9 very early

Record when the earliest variety is in full flower and score all varieties relative to this. An assessment on one occasion is normally sufficient. Estimate the date of full flowering for the earliest control variety.

C.6.3.5 MATURITY from all plots

(Spring linseed - OBLIGATORY)
(1-9) (Winter linseed - ADDITIONAL)
(1-9)

Maturity should be judged by making a visual estimate of canopy senescence, where;

- 1 very late
- 9 very early

Unrepresentative areas of the plot should be avoided when making assessments, for example, localised diseased infections.

C.6.3.6 **SOWING DATE of each trial** (OBLIGATORY) (Day/month/year)

This is recorded in Part 1 of the Site Information Form.

C.6.3.7 **HARVEST DATE** (OBLIGATORY) (Day/month/year)

This is recorded in part 2 of the Site Information Form.

C.6.3.8 **BIRD DAMAGE from all plots** (OBLIGATORY) (1-9)

- 1 all plants severely damaged
- 9 no plants damaged

Indicate the cause of damage and, in the Diary section, what action has been taken to minimise further damage.

C.6.3.9 **SEED (or BOLL) LOSS from all plots** (OBLIGATORY) (1-9)

- 1 severe seed loss
- 9 no seed loss

and give an estimation of maximum % seed loss/boll loss.

Record before harvest if serious loss has already occurred. Base scores either on observation of boll loss or counts on the ground. Ensure that combines are set correctly to minimise losses at harvest. Assess any serious combining losses after harvest.

C.6.3.10 COMBINE LOSSES from all plots (OBLIGATORY) (1-9)

This must be recorded.

9 = no combine losses. Combine losses should be assessed if the losses are thought sufficient to exclude the yield data from results. Indicate the estimated number of grains lost per m² for the lowest score given on the 1 to 9 scale.

C.6.3.11 Site factors

Any factors which may have affected the yield of the trial or individual plots must be noted and accompany the yield data.

Where varietal differences are seen in pest or disease attack, records should be made either as an estimated % of plants affected, or as % leaf area attacked in accordance with the procedure in Section D for disease.

Records for other scores should be taken as % plants affected or on a 1 to 9 scale. Include definitions of 1 to 9 on the scale.

C.6.3.12 Trial inspection

All trials will be inspected by the Trial Inspection and Technical Validation Operator, and, in some cases, it may be necessary to visit on more than one occasion.

The requirements for Growing Trial Operators in respect of inspections are to:

1. Give inspectors reasonable access to trials and to provide full location and site details (if not already given with Site Data 1).
2. Provide the inspector with information (for example pesticide sprays applied etc) within seven days of a request.
3. Co-operate with the inspector in making any non-routine assessments required to establish the validity of the trial (for example population counts).
4. Carry out any action agreed in consultation with the inspector. In particular it is important that any requirement to shorten plots is undertaken. The data on plots that the trials operator and inspector agree to exclude should not be submitted.

Section D – Disease testing procedures

D.1 Assessment of natural infection

D.1.1 Disease observation tussocks

No disease observation tussocks are carried out routinely.

D.2 Naturally occurring disease in VCU growing trials

D.2.1 If disease levels increase to levels more than 5% of the leaf area (or 5% of infected plants as appropriate for the diseases) on the most affected variety a score should be made and sent to the Trials Organiser. Confirmation of the identity of a disease should be obtained from an appropriate plant pathologist if required. If disease infection persists, successive records should be made through the season.

D.2.1.1 The disease most likely to be encountered is Powdery mildew (*Oidium lini*), though both *Alternaria* and *Botrytis* may cause infections on the leaves. Capsules are most likely to be affected by *Alternaria* and *Botrytis*. Stem and whole plant symptoms are most likely to be caused by *Verticillium*, *Sclerotinia*, *Mycosphaerella* (Pasma disease), *Phoma*, *Fusarium* wilt and other *Fusarium* diseases. *Fusarium* wilt can cause the entire plant to wilt. If recorded, the trial location should not be used again due to long-term persistence in the soil.

D.2.2 Recording methods

D.2.2.1 Timing of assessments

A guide to probable assessment times in terms of growth stage is shown in the Table below:

Disease	Seedling/ Vegetative	Flower bud	Flowering	Capsule formation	Pre- maturity
Powdery mildew %	N/A	√	√	√	N/A
Botrytis %	N/A	√	√	√	N/A
Alternaria %	N/A	N/A	√	√	N/A
Fusarium %	√	√	√	√	N/A
Fusarium wilt %	√	N/A	√	N/A	N/A
Verticillium %	N/A	N/A	N/A	√	√
Sclerotinia %	N/A	N/A	√	√	N/A
Phoma %	√	√	√	√	N/A
Mycosphaerella %	√	N/A	N/A	√	√

D.2.2.2 Appropriate assessment keys are given in Appendix 8. All disease records to be sent to the Trial Design and Data Handling Operator as soon as they are made.

Disease data should be received by 13th August

D.3 Inoculated disease tests

No inoculated disease tests are carried out routinely.

Section E – Quality testing procedures

E.1 Responsibilities

E.1.1 The Quality Testing Operator appointed by the Trials Organiser is responsible for conducting approved quality tests according to these procedures.

E.2 Quality assessment methodology

E.2.1 Moisture content determination

The following procedure must be followed:

A 105 g sample (± 5 g) is placed in the drier which must be at a temperature of $100^{\circ}\text{C} \pm 4^{\circ}\text{C}$ with the air recirculator set in the range 80-100% recirculation in order to restore the temperature to $100^{\circ}\text{C} \pm 4^{\circ}\text{C}$ as rapidly as possible. When the temperature is restored to $100^{\circ}\text{C} \pm 4^{\circ}\text{C}$ the air regulator is set at 80% recirculation i.e. 20% fresh hot air. The air regulator is critical for even rapid drying. The samples are dried at $100^{\circ}\text{C} \pm 4^{\circ}\text{C}$ for such time as is necessary for complete drying.

The dried sample is carefully removed from the drier as soon as the sample is cool enough for accurate weighing. The dry weight is recorded to one decimal place.

When all samples from a given trial have been recorded, the fresh and dry weights are immediately reported to the Trials Organiser electronically using the character names given in Section D.10.3. When the dry weights are reported as a percentage, the fresh weight should be reported as 100.

Moisture content determination by conductance moisture meter is not acceptable to the Testing Authority.

E.2.2 Oil content determination

Analysis is performed using continuous emission NMR following ISO 5511:1992. Results are expressed as apparent oil as a percentage at 9% moisture.

The stability of the equipment is checked at two-hourly intervals through the working day by the use of weighed oil standards. A single determination is normally performed on each test sample.

Section F – Trial design and data handling procedures

F.1 Plan validation and storage

F.1.2 After the trial has been drilled, the Growing Trial Operator must:

- 1) Confirm that the trial has been drilled according to plan and provide the sowing date, by returning site data 1 and associated trial sketch to the appropriate Trial Design and Data Handling Operator.
- 2) If any amendments to the plan have been made, return a hard copy of the plan to the appropriate Trial Design and Data Handling Operator with any amendments clearly indicated. Alternatively, amendments may be notified electronically with the agreement of the Trial Design and Data Handling Operator.

F.1.3 The Trial Design and Data Handling Operator will check these for statistical validity.

F.2 Data recording

F.2.1 Data are recorded using the methods and characters given in Sections C, D and E.

F.2.2. Site information is recorded for each trial including, for example, data on previous cropping, seed rates, soil details and fertiliser applications.

F.2.3 Details of any agrochemical applications are also recorded and forwarded to the Trials Organiser.

F.3 Data processing

F.3.1 Processing of individual agronomic and disease variates.

F.3.2. A list of the agronomic, yield and disease variates, which may be recorded and processed, are specified in Sections C, D and E. After scrutiny, copies of the results will be returned to the Growing Trial Operator for action as agreed by the Trials Organiser.

F.4 Other tests and trials

F.4.1 Any additional or alternative designs required for the assessment of additional VCU characters not detailed in Annex A of the **MINOR CROPS VCU TRIAL PROTOCOL** will be added to these **Procedures** as and when approved by the NLSC.

Supporting Document for Appendices

Appendices for this main procedure are stored in a separate document, which is updated closer to the start of the growing trial to include the latest information on controls and trial organisers. This will be published on [VCU protocols and procedures for testing agricultural crops](#) - GOV.UK (www.gov.uk).



© Crown copyright 2025

You may re-use this information (excluding logos) free of charge in any format or medium, under the terms of the Open Government Licence v.3. To view this licence visit www.nationalarchives.gov.uk/doc/open-government-licence/version/3/ or email PSI@nationalarchives.gov.uk.

Data Protection:

For information on how we handle personal data visit www.gov.uk and search Animal and Plant Health Agency Personal Information Charter.

This publication is available at www.gov.uk/government/publications

Any enquiries regarding this publication should be sent to us at

webmaster@apha.gov.uk

www.gov.uk/apha

APHA is an Executive Agency of the Department for Environment, Food and Rural Affairs and also works on behalf of the Scottish Government, Welsh Government and Food Standards Agency to safeguard animal and plant health for the benefit of people, the environment and the economy.