

Monthly water situation report: Solent and South Downs Area

Summary - November 2025

Solent and South Downs (SSD) had close to average rainfall in November, receiving 87% (93mm) of the long term average (LTA) of 106.6mm. Monthly mean river flows for November ranged from below normal to normal across SSD. End of month groundwater levels for November ranged from notably low to above normal. Soils across SSD ended the month wetter than the LTA for November.

End of month reservoir stocks were below average at Ardingly Reservoir (Ouse) and at Arlington Reservoir (Cuckmere).

1.1 Rainfall

SSD had close to average rainfall in November, receiving 87% (93mm) of LTA (106.6mm). The Test Chalk areal unit received the highest monthly rainfall total with 112% (113mm) of LTA (100.9mm) during November. The Sussex Coast areal unit received the lowest rainfall total with 75% (71mm) of LTA (95.3mm).

The highest daily rainfall totals were recorded on 22 November. 30.4mm was recorded at Chilgrove (West Sussex Chalk), and 30mm at Folkington (Cuckmere). This was a widespread rainfall event with daily totals in excess of 20mm recorded across the whole area from the Test Chalk to the Pevensey Levels. There were no completely dry days recorded during November.

1.2 Soil moisture deficit and recharge

Soils across SSD ended the month wetter than the LTA for November.

1.3 River flows

Monthly mean river flows for November ranged from below normal to normal across SSD.

Flows were below normal on the:

River Meon at Mislingford

Flows were normal on the:

- River Test at Broadlands
- River Test at Chilbolton
- River Itchen at Allbrook and Highbridge
- River Lymington at Brockenhurst
- River Medina at Blackwater
- River Rother at Iping Mill
- River Ouse at Goldbridge

- River Cuckmere at Cowbeech
- River Arun at Alfoldean
- River Wallington at North Fareham
- River Adur at Sakeham

1.4 Groundwater levels

End of month groundwater levels for November ranged from notably low to above normal across SSD.

Groundwater levels were notably low at:

- Chilgrove (West Sussex Chalk)
- Beeding Hill (West Sussex Chalk)

Groundwater levels were below normal at:

• Catherington (East Hampshire Chalk)

Groundwater levels were normal at:

- Clanville Gate (Test Chalk)
- Lopcombe Corner (Test Chalk)
- West Meon (East Hampshire Chalk)
- Harting Common (Western Rother Greensand)
- Houndean Bottom (East Sussex Chalk)
- Carisbrooke Castle (Isle of Wight)
- Cornish Farm (East Sussex Chalk)
- Youngwoods Copse (Isle of Wight)

Groundwater levels were above normal at:

Preston Candover (East Hampshire Chalk)

The end of month groundwater level at Beeding Hill (East Sussex Chalk) was the fourth lowest for November in a record going back to 1979.

1.5 Reservoir stocks

End of month reservoir stocks were below average for both Ardingly and Arlington Reservoirs. Ardingly Reservoir (Ouse) was at 41.5% of total capacity (LTA 77%) and Arlington Reservoir (Cuckmere) was at 64.7% of total capacity (LTA 72.4%).

1.6 Environmental impact

1.6.1 Abstraction licence restriction

At the start of November there were 12 licence restrictions in force. By the end of the month 10 licence restrictions were in place. These included:

- one licence on the Ouse
- two licences on the River Meon
- one licence on the River Hamble
- one licence on the Sowley Stream
- one licence on the River Ems
- one licence on the Western Rother
- four licences on the Test
- one licence on the Itchen

1.6.2 Flood Alerts and Warnings in Force

There was 1 Fluvial Flood Alert issued in November for the Lower Itchen.

There were no Flood Warnings issued in November.

Author: <u>HydrologySSD@Environment-agency.gov.uk</u>


Contact Details: 03708 506506

All data are provisional and may be subject to revision. The views expressed in this document are not necessarily those of the Environment Agency. Its officers, servants or agents accept no liability for any loss or damage arising from the interpretation or use of the information, or reliance upon views contained in this report.

2. Rainfall

2.1 Rainfall map

Figure 2.1: Total rainfall for hydrological areas for the current month (up to 30 November 2025), the last 3 months, the last 6 months, and the last 12 months, classed relative to an analysis of respective historic totals. Table available in the appendices with detailed information.




Rainfall data for January 2025 onwards, extracted from Environment Agency 1km gridded rainfall dataset derived from Environment Agency intensity rain gauges. (Source: Environment Agency. Crown Copyright, 100024198, 2025). Rainfall data prior to January 2025, extracted from Met Office HadUK 1km gridded rainfall dataset derived from registered rain gauges (Source: Met Office. Crown copyright, 2025).

3 River flows and Groundwater levels

3.1 River flows and Groundwater level map

Figure 3.1: Monthly mean river flow and groundwater levels at our indicator sites for November 2025, expressed as a percentage of the respective long term average and classed relative to an analysis of historic November monthly means. Table available in the appendices with detailed information.




(Source: Environment Agency). Crown copyright. All rights reserved. Environment Agency, 100024198, 2025. Geological map reproduced with kind permission from UK Groundwater Forum, BGS copyright NERC. Crown copyright. All rights reserved. Environment Agency, 100024198, 2025.

4 West Hampshire

4.1 West Hampshire Rainfall and effective rainfall charts


Figure 4.1: Monthly rainfall and effective rainfall totals for the past 12 months compared to the 1991 to 2020 long term average.



HadUK rainfall data. (Source: Met Office. Crown copyright, 2025).


4.2 West Hampshire River flow charts

Figure 4.2: Daily mean river flow for index sites over the past year, compared to an analysis of historic daily mean flows, and long term maximum and minimum flows.



4.3 West Hampshire Groundwater level charts


Figure 4.3: End of month groundwater levels at index groundwater level sites for major aquifers. 34 months compared to an analysis of historic end of month levels and long term maximum and minimum levels.



5 East Hampshire and Isle of Wight

5.1 East Hampshire and Isle of Wight Rainfall and Effective rainfall charts


Figure 5.1: Monthly rainfall and effective rainfall totals for the past 12 months compared to the 1991 to 2020 long term average.



HadUK rainfall data. (Source: Met Office. Crown copyright, 2025).


5.2 East Hampshire and Isle of Wight River flow charts

Figure 5.2 Daily mean river flow for index sites over the past year, compared to an analysis of historic daily mean flows, and long term maximum and minimum flows.



5.3 East Hampshire and Isle of Wight Groundwater level charts


Figure 5.3: End of month groundwater levels at index groundwater level sites for major aquifers. 34 months compared to an analysis of historic end of month levels and long term maximum and minimum levels.



6 West Sussex

6.1 West Sussex Rainfall and Effective Rainfall charts


Figure 6.1: Monthly rainfall and effective rainfall totals for the past 12 months as a percentage of the 1991 to 2020 long term average.



HadUK rainfall data. (Source: Met Office. Crown copyright, 2025).


6.2 West Sussex River flow charts

Figure 6.2: Daily mean river flow for index sites over the past year, compared to an analysis of historic daily mean flows, and long term maximum and minimum flows.



6.3 West Sussex Groundwater level charts


Figure 6.3: End of month groundwater levels at index groundwater level sites for major aquifers. 34 months compared to an analysis of historic end of month levels and long term maximum and minimum levels.



7 East Sussex

7.1 East Sussex Rainfall and Effective Rainfall charts


Figure 7.1: Monthly rainfall and effective rainfall totals for the past 12 months compared to the 1991 to 2020 long term average.



HadUK rainfall data. (Source: Met Office. Crown copyright, 2025).


7.2 East Sussex River flow charts

Figure 7.2: Daily mean river flow for index sites over the past year, compared to an analysis of historic daily mean flows, and long term maximum and minimum flows.




7.3 East Sussex Groundwater level charts

Figure 7.3: End of month groundwater levels at index groundwater level sites for major aquifers. 34 months compared to an analysis of historic end of month levels and long term maximum and minimum levels.



8 Reservoir stocks

Figure 8.1: End of month reservoir stocks compared to long term maximum, minimum and average stocks. Note: Historic records of individual reservoirs and reservoir groups making up the regional values vary in length.



(Source: water companies).

9 Glossary

9.1 Terminology

Aquifer

A geological formation able to store and transmit water.

Areal average rainfall

The estimated average depth of rainfall over a defined area. Expressed in depth of water (mm).

Artesian

The condition where the groundwater level is above ground surface but is prevented from rising to this level by an overlying continuous low permeability layer, such as clay.

Artesian borehole

Borehole where the level of groundwater is above the top of the borehole and groundwater flows out of the borehole when unsealed.

Cumecs

Cubic metres per second (m^{3s-1}).

Effective rainfall

The rainfall available to percolate into the soil or produce river flow. Expressed in depth of water (mm).

Flood alert and flood warning

Three levels of warnings may be issued by the Environment Agency. Flood alerts indicate flooding is possible. Flood warnings indicate flooding is expected. Severe flood warnings indicate severe flooding.

Groundwater

The water found in an aquifer.

Long term average (LTA)

The arithmetic mean calculated from the historic record, usually based on the period 1991 to 2020. However, the period used may vary by parameter being reported on (see figure captions for details).

mAOD

Metres above ordnance datum (mean sea level at Newlyn Cornwall).

MORECS

Met Office Rainfall and Evaporation Calculation System. Met Office service providing real time calculation of evapotranspiration, soil moisture deficit and effective rainfall on a 40 by 40 km grid.

Naturalised flow

River flow with the impacts of artificial influences removed. Artificial influences may include abstractions, discharges, transfers, augmentation and impoundments.

NCIC

National Climate Information Centre. NCIC area monthly rainfall totals are derived using the Met Office 5 km gridded dataset, which uses rain gauge observations.

Recharge

The process of increasing the water stored in the saturated zone of an aquifer. Expressed in depth of water (mm).

Reservoir gross capacity

The total capacity of a reservoir.

Reservoir live capacity

The capacity of the reservoir that is normally usable for storage to meet established reservoir operating requirements. This excludes any capacity not available for use (for example, storage held back for emergency services, operating agreements or physical restrictions). May also be referred to as 'net' or 'deployable' capacity.

Soil moisture deficit (SMD)

The difference between the amount of water actually in the soil and the amount of water the soil can hold. Expressed in depth of water (mm).

9.2 Categories

Exceptionally high

Value likely to fall within this band 5% of the time.

Notably high

Value likely to fall within this band 8% of the time.

Above normal

Value likely to fall within this band 15% of the time.

Normal

Value likely to fall within this band 44% of the time.

Below normal

Value likely to fall within this band 15% of the time.

Notably low

Value likely to fall within this band 8% of the time.

Exceptionally low

Value likely to fall within this band 5% of the time.

10 Appendices

10.1 Rainfall, effective rainfall and soil moisture deficit table

(Source: Met Office. Crown copyright, 2025). All rights reserved. Environment Agency, 100024198, 2025

Figure 10.1: This is areal rainfall, effective rainfall (percolation or runoff) and soil moisture deficit for the hydrological areas across the SSD. There may be significant variation within each area which must be considered when interpreting these data. When additional meteorological data is available estimates are revised which will affect the period totals in section 10.2

Hydrological Area	Rainfall (mm) 30 day Total	Rainfall November as %LTA	Effective Rainfall (mm) 30 day Total	Effective Rainfall November as %LTA	Soil Moisture Deficit (SMD) Day 30	SMD End of November LTA
Test Chalk	113	112%	15	31%	23	26
East Hampshire Chalk	109	97%	38	56%	0	15
West Sussex Chalk	99	86%	63	82%	0	14
East Sussex Chalk	95	87%	47	76%	0	24
Isle of Wight	91	84%	27	57%	0	34
Western Rother Greensand	102	86%	53	67%	0	13
Hampshire Tertiaries	98	93%	33	59%	0	17
Lymington	99	91%	65	101%	0	15
Sussex Coast	71	75%	0	0%	4	28
Arun	83	81%	37	60%	0	14
Adur	86	83%	65	101%	0	14
Ouse	86	81%	71	103%	0	13
Cuckmere	84	79%	70	99%	0	14
Pevensey Levels	88	86%	58	98%	0	19
SSD Average	93	87%	46	74%	2	19

10.2 Seasonal summary table of rainfall and effective rainfall

Winter season: 01/10/2025 to 30/11/2025

Hydrological Area	Seasonal Rainfall (mm) Total	Seasonal Rainfall as % LTA	Seasonal Effective Rainfall (mm)Total	Seasonal Effective Rainfall as % LTA
Test Chalk	202	103%	27	38%
East Hampshire Chalk	221	101%	53	53%
West Sussex Chalk	226	101%	80	71%
East Sussex Chalk	220	104%	64	69%
Isle of Wight	212	101%	43	61%
Western Rother Greensand	229	99%	70	60%
Hampshire Tertiaries	205	101%	33	44%
Lymington	225	107%	65	75%
Sussex Coast	173	94%	0	0%
Arun	201	100%	37	43%
Adur	204	100%	65	70%
Ouse	210	100%	80	75%
Cuckmere	208	100%	79	73%
Pevensey Levels	210	105%	58	68%
SSD Average	210	101%	54	60%

10.3 Rainfall banding table

Hydrological area	November 2025 band	September 2025 to November 2025 cumulative band	June 2025 to November 2025 cumulative band	December 2024 to November 2025 cumulative band
Test Chalk	Above normal	Normal	Normal	Below normal
East Hampshire Chalk	Normal	Above normal	Normal	Normal
West Sussex Chalk	Normal	Above normal	Normal	Normal
East Sussex Chalk	Normal	Above normal	Normal	Normal
Isle of Wight	Normal	Above normal	Normal	Normal
Western Rother Greensand	Normal	Above normal	Normal	Normal
Hampshire Tertiaries	Normal	Above normal	Normal	Normal
Lymington	Normal	Above normal	Normal	Normal
Sussex Coast	Normal	Normal	Normal	Below normal
Arun	Normal	Normal	Normal	Normal
Adur	Normal	Above normal	Normal	Normal
Ouse	Normal	Above normal	Normal	Normal
Cuckmere	Normal	Above normal	Normal	Normal
Pevensey Levels Normal		Above normal	Normal	Normal

10.4 River flows table

Site name	River	River Catchment		October 2025 band	
Alfoldean Gs	Arun	Arun	Normal	Normal	
Allbrook Gs+ Highbridge	Itchen (so)	Itchen	Normal	Normal	
Blackwater	Medina	Isle of Wight	Normal	Normal	
Broadlands	Test	Test Lower	Normal	Below normal	
Brockenhurst GS	Lymington	New Forest	Normal	Normal	
Chilbolton GS	Test	Test Upper	Normal	Normal	
Cowbeech Gs	Cuckmere	Cuckmere	Normal	Normal	
Goldbridge Gs	Ouse [so]	Ouse Sussex	Normal	Normal	
Iping Mill Gs	Rother	West Rother	Normal	Normal	
Mislingford GS	Meon	Meon	Below normal	Below normal	
North Fareham GS	Wallington	Wallington	Normal	Normal	
Sakeham GS	Adur	Adur	Normal	Normal	

10.5 Groundwater table

Site name	Aquifer	End of November 2025 band	End of October 2025 band	
Carisbrooke Castle	Isle Of Wight Central Downs Chalk	Normal	Normal	
Youngwoods Copse	Isle of Wight Lower Greensand	Normal	Normal	
Clanville Gate Gwl	River Test Chalk	Normal	Normal	
Lopcombe Corner Gwl	River Test Chalk	Normal	Below normal	
Preston Candover	River Itchen Chalk	Above normal	Above normal	
West Meon Hut Gwl	River Itchen Chalk	Normal	Normal	
Catherington	River Meon Chalk	Below normal	Below normal	
Chilgrove House Gwl	Chichester- Worthing- Portsdown Chalk	Notably low	Exceptionally low	
Beeding Hill Gwl	Brighton Chalk Block	Notably low	Exceptionally low	
Houndean Bottom Gwl	Brighton Chalk Block	Normal	Below normal	
Harting Common Down	Western Rother Lower Greensand	Normal	Normal	
Cornish Wigdens Gwtr Eastbourne Chalk Block		Normal	Above normal	

10.6 Abstraction licence flow constraints

Number of	Number of	Number of	Number of
flow	flow	flow	flow
constraints in	constraints in	constraints in	constraints in
force between	force between	force between	force between
1 to 11	12 to 19	20 to 25	25 to 30
November	November	November	November
2025	2025	2025	2025
12	9	12	10

10.7 Solent and South Downs Areal Rainfall Units Map



10.8 SSD Areal Rainfall Monthly Long Term Averages

Hydrological Area	Jan LTA mm	Feb LTA mm	Mar LTA mm	Apr LTA mm	May LTA mm	Jun LTA mm	Jul LTA mm	Aug LTA mm	Sep LTA mm	Oct LTA mm	Nov LTA mm	Dec LTA mm
Test Chalk	92.6	65.7	59.4	60.5	53.7	57.8	57.5	63.8	62.8	95.3	100.9	93.1
East Hampshire Chalk	102.7	72.1	62.7	63.5	54.4	62.1	60.5	71.4	70.2	105.8	112.8	105.3
West Sussex Chalk	104.6	73.7	59.5	59.1	54.4	59.4	60.9	73.0	74.4	108.8	114.8	113.2
East Sussex Chalk	96.4	66.9	53.8	49.9	51.2	55.6	57.3	67.6	65.3	101.2	110.6	106.9
Isle of Wight	93.5	66.9	55.4	52.6	47.9	50.6	47.8	59.7	65.9	102.2	107.4	105.6
Western Rother Greensand	110.6	77.5	61.9	64.1	56.4	59.6	57.9	73.1	73.7	111.7	118.1	115.1
Hampshire Tertiaries	95.4	66.7	58.1	57.8	49.8	56.3	51.2	64.7	65.2	97.3	105.1	99.4
Lymington	96.7	67.4	59.6	58.2	50.1	54.5	51.2	62.9	66.6	100.1	109.5	105.3
Sussex Coast	86.8	59.1	48.5	49.6	45.3	50.6	48.8	59.3	59.9	89.3	95.3	93.3
Arun	96.8	67.1	54.7	55.3	53.2	53.3	54.7	64.6	65.8	97.7	103.4	98.9
Adur	94.7	65.6	52.7	52.8	53.5	52.5	53.8	65.1	63.5	99.4	102.9	100.2
Ouse	97.7	67.7	53.9	51.2	53.4	54.9	57.5	67.4	64.7	104.6	105.9	105.4
Cuckmere	94.4	65.8	51.0	49.0	50.5	55.8	56.8	68.7	65.4	101.9	106.8	105.5
Pevensey Levels	89.5	62.7	49.5	47.6	48.1	51.8	56.3	64.6	62.9	96.3	103.0	102.9
SSD Average	96.8	67.7	56.9	56.5	52.0	56.0	55.3	66.0	66.2	100.5	106.6	102.4