



UK Health
Security
Agency

Guidance for public health management of meningococcal disease in the UK

Updated December 2025

Contents

Document history	3
1. Background	5
1.1 Objectives of guidelines	6
2. Epidemiology of IMD in the UK	7
2.1 General epidemiology	7
2.2 Meningococcal group C	7
2.3 Meningococcal group W	7
2.4 Meningococcal group Y	7
2.5 Meningococcal group B	8
2.6 Other meningococcal groups	8
3. Vaccination programmes	9
3.1 MenC conjugate vaccines	9
3.2 Quadrivalent MenACWY vaccines	9
3.3 MenB vaccine	9
4. Additional guidance	11
5. Pre-admission management recommendations	12
6. Laboratory investigation of suspected cases	13
7. Role of public health	17
8. Public health action after a case	20
8.1 Risk to close contacts	20
8.2 Risk reduction through chemoprophylaxis	24
8.3 Choice of agent for chemoprophylaxis	27
8.4 Pregnancy and breastfeeding	31
8.5 Aim of vaccination	31
8.6 Disseminating information	34
9. Chemoprophylaxis in healthcare settings	37
10. Cluster and outbreak management	39
10.1 Clusters/outbreaks in a household setting	41
10.2 Management of clusters/outbreaks in an educational or residential setting	41
10.3 Vaccination in pregnancy	50
10.4 Management of clusters/outbreaks in the wider community	50
10.5 Disseminating information in cluster/outbreak management	51
11. Appendix 1	54
References	55
About the UK Health Security Agency	63

Document history

Date	Change details
August 2019	<p>The definition of a 'probable' IMD case has been clarified.</p> <p>Confirmation that the recent EU review of fluoroquinolones does not affect guidance on ciprofloxacin chemoprophylaxis for meningococcal disease.</p> <p>Definition of a cluster has been clarified.</p> <p>The importance of investigating for underlying risk factors in IMD patients due to unusual meningococcal serogroups, non-groupable strains or recurrent meningococcal disease has been highlighted and Appendix 5 added for communication with GPs concerning this.</p>
September 2024	<p>Document transferred from PHE to UKHSA branding.</p> <p>Updated epidemiology section.</p> <p>Consideration for Rifampicin prophylaxis following cases arising after travel to the Middle East and Asia, section added.</p> <p>Updated laboratory investigations and national surveillance sections.</p> <p>Contact details for all nations updated where appropriate.</p> <p>Aligning recommendations with NICE guidance (NG240, 19 March 2024).</p> <p>Isolation of meningococci from non-sterile sites – specifically covering urogenital and anorectal infections.</p> <p>Close contact definitions, including shisha sharing.</p> <p>Updated considerations for cluster/outbreak scenarios and recommended MenB outbreak vaccine schedules.</p> <p>Guidance on vaccination in pregnancy.</p>
October 2024	<p>Consideration of shisha sharing updated.</p>
November 2024	<p>Updated ciprofloxacin section under '8.3 Choice of agent for chemoprophylaxis'.</p>
December 2025	<p>Emphasis on the need to offer prophylaxis without delay, for hospitals to support this and antibiotic chemoprophylaxis should not be delayed for young children because of lack of availability or difficulties in obtaining liquid ciprofloxacin suspension see Recommendation 4: Indications for antibiotic prophylaxis.</p> <p>For close contacts of IMD cases with confirmed or likely ciprofloxacin resistance (for example, recent travel to the Middle East or Asia), the priority continues to be use of rifampicin for prophylaxis. Advice on recommended alternatives when rifampicin is not available is covered in the section on Rifampicin and travel.</p>

Date	Change details
	Definitions of clusters/ outbreaks have been updated for clarity, including in a household and educational setting under Section 10 Cluster and outbreak management, sections 10.1 and 10.2.

1. Background

Neisseria meningitidis is a major cause of septicaemia and meningitis worldwide and is associated with significant mortality as well as serious long-term sequelae among survivors ([1](#)). Six meningococcal capsular groups (A, B, C, W, X and Y) distinguished by their polysaccharide capsule cause almost all invasive infections in humans. Epidemiology differs by serogroup and clonal complex combination. The meningococcus commonly colonises the nasopharynx and carriage prevalence increases through childhood and peaks at 18 to 20 years before declining ([2](#), [3](#)). Carriage episodes may last for many months ([4](#)). Invasive disease is a rare outcome of acquisition but onward transmission from cases to close contacts can rarely result in secondary cases, as well as clusters of infection ([5](#), [6](#), [7](#)). Fewer than 2% of invasive meningococcal disease (IMD) cases, however, are considered to result from close contact with a primary IMD case ([8](#)). In 2014, prior to the introduction of routine MenB infant and MenACWY teenage vaccination, annual IMD incidence across all age groups was approximately 1.2 per 100,000 in the UK ([9](#)) with incidence in 2023/24 at 0.6 per 100,000ⁱ.

Systemic immunity, as measured by serum bactericidal antibodies, usually develops within 14 days of acquisition of meningococci ([10](#)). Rarely, acquisition may progress to invasive disease before immunity develops. This incubation period is usually 3 to 5 days, based on data from studies of laboratory-acquired infection ([11](#)), from occasional clusters where the date of exposure is known ([12](#)) and from carriage studies among military recruits ([13](#)). Established meningococcal carriers do not usually develop invasive disease ([13](#)). The risk of invasive disease following acquisition is likely to vary with environmental and host factors, but will also depend critically on the characteristics of the strain acquired. Only a very small proportion of carried strains are responsible for invasive disease ([14](#)).

Conjugate vaccines against group C meningococci (MenC) have been available since the late 1990s and quadrivalent conjugate vaccines against groups A, C, W and Y (MenACWY) have been licensed in Europe for almost 2 decades. In early 2013, a new vaccine developed specifically to prevent disease caused by group B meningococci (MenB), was licensed in Europe (4CMenB, Bexsero®, GSK Biologicals, Belgium). This vaccine was unlike the pre-existing MenC and MenACWY conjugate vaccines in that it is protein-based and, therefore, has a different mechanism of action compared with conjugate vaccines along with different safety, reactogenicity and immunogenicity profiles in different age groups ([15](#)). In 2017, another protein-based MenB vaccine, rLP2086 (Trumenba®), was licensed in Europe for individuals aged 10 years and older. This guidance includes the potential use of MenB vaccine in the public health management of cases and contacts of IMD.

ⁱ [Meningococcal disease: laboratory-confirmed cases in England 2023 to 2024 - GOV.UK](#)

1.1 Objectives of guidelines

This guidance covers pre-admission management, investigation of suspected cases, case definitions, chemoprophylaxis and vaccination of cases and close contacts of sporadic cases, as well as management of IMD clusters and outbreaks (defined in focus 3 box).

This 2024 UK guidance update has been reviewed by the UK Health Security Agency (UKHSA) Vaccine Preventable Invasive Bacterial Infections (VaPIBI) Forum. Previous updates were made in 2019, while the 2018 update replaced the 2012 Guidance for public health management of meningococcal disease in the UK and the 2014 guidance on preventing secondary cases of MenB disease. The current update includes more recent data on disease epidemiology, together with updated advice on vaccination of cases and close contacts, including public health action after infections of the genitourinary tract.

2. Epidemiology of IMD in the UK

2.1 General epidemiology

Due to SARS-CoV-2 restrictions during 2020 and 2021, the number of laboratory-confirmed IMD cases decreased substantially across all age groups and for all serogroups ([16](#)). Following the easing of restrictions after July 2021, there was a notable increase in serogroup B IMD cases, with numbers in adolescents and young adults comparable to those observed pre-pandemic. IMD cases in other age groups and from other serogroups remained very low throughout 2021 to 2022 and 2023 to 2024. MenB cases began to increase in all age groups in 2022 to 2023 whereas non-MenB disease has remained very low. Disease levels appeared to stabilise in 2023 to 2024.

2.2 Meningococcal group C

The MenC conjugate vaccine was implemented into the UK national immunisation programme in 1999 and was highly effective in preventing MenC IMD across all age groups through direct and indirect (herd) protection ([17](#)). MenC is currently well-controlled, with only 30 to 40 cases annually prior to the pandemic ([17](#), [18](#)). Current cases are mainly diagnosed in adults born outside the UK who were unvaccinated against MenC disease and small numbers of cases arising in previously-vaccinated children.

2.3 Meningococcal group W

MenW cases increased with the emergence and rapid expansion of a hyperinvasive strain belonging to the ST-11 complex ([19](#), [20](#)) in 2010 and is now controlled following implementation of an emergency MenACWY immunisation programme for teenagers in August 2015. MenACWY vaccination is now routinely offered at 13 to 14 years and MenW IMD cases are currently very low across all age groups ([21](#)). Imported cases e.g. associated with Umrah pilgrimage remain a concern and it is vital that pilgrims get the required vaccinations including MenACWY conjugate.

2.4 Meningococcal group Y

MenY disease remains uncommon in the UK and predominantly affects older adults and those with underlying health conditions. MenY IMD cases declined further across all age groups following implementation of an emergency MenACWY immunisation programme for teenagers in August 2015 and remain low with the routine teenage MenACWY vaccine programme ([21](#)).

2.5 Meningococcal group B

MenB IMD cases fell between 2001 to 2002 and 2013 to 2014, likely because of secular trends ([22](#)). MenB accounted for nearly 90% of cases during 2006 to 2011, with an overall incidence of nearly 2 per 100,000. The implementation of 4CMenB (Bexsero®) into the national infant immunisation programme since September 2015 resulted in a large and significant reduction in MenB IMD cases in vaccinated children ([23](#), [24](#)). MenB IMD cases have predominated with the early re-emergence of disease since restrictions to control the COVID-19 pandemic were withdrawn in July 2021 ([25](#)).

2.6 Other meningococcal groups

Other meningococcal groups rarely cause invasive disease in the UK, with most cases occurring in those with underlying health conditions.

2.7 Global epidemiology

In other parts of the world different serogroups may predominate. Large epidemics of both capsular group A and W meningococcal infection have occurred in association with Hajj pilgrimages, and proof of vaccination against A, C, W and Y capsular groups is now a visa entry requirement for pilgrims and seasonal workers travelling to the Kingdom of Saudi Arabia. Epidemics arise unpredictably throughout sub-saharan Africa with MenA previously predominating before the successful introduction of MenA conjugate vaccine ([26](#)). Subsequent outbreaks have been due to serogroups C, W and X.

3. Vaccination programmes

3.1 MenC conjugate vaccines

MenC vaccine (Meningitec®, Menjugate® or NeisVac®) was included in the routine infant programme from November 1999 but is no longer part of the national immunisation programme ([27](#), [18](#), [28](#)). Currently, protection against MenC includes a single Hib/MenC combination vaccine (Menitorix®) at one year of age, but this vaccine will no longer be available in 2025.

3.2 Quadrivalent MenACWY vaccines

A MenACWY conjugate vaccine (Menveo®, Nimenrix®, MenQuadfi®) is offered at 13 to 14 years of age and those who missed vaccination remain eligible up to their 25th birthday. This includes those who missed the school's programme and those attending University for the 1st time, including students arriving in the UK from countries that may not have offered this vaccination. MenACWY conjugate vaccines confer no protection against MenB.

3.3 MenB vaccine

4CMenB (Bexsero®; GSK) was licensed in Europe in early 2013 and is a protein based vaccine containing 4 main components: factor H binding protein (fHbp) variant 1.1, Neisserial Adhesin A (NadA), Neisseria Heparin Binding Antigen (NHBA) and the New Zealand Outer Membrane Vesicle (OMV) vaccine (MenZB) which incorporates Porin A (PorA) P1.4 ([15](#)).

4CMenB (Bexsero®) was implemented in the infant programme on 1 September 2015. Infants born in May 2015 were eligible for the vaccine at 16 weeks and 1 year, those born in June were eligible for the vaccine at 12 weeks, 16 weeks and 1 year, and those born since 1 July 2015 are offered the vaccine at 8 weeks, 16 weeks and 1 year of age. 4CMenB (Bexsero®) has been highly effective in preventing MenB IMD in vaccinated children ([24](#), [23](#), [29](#)). 4CMenB (Bexsero®) may also protect against other meningococcal serogroups that possess cross-reactive vaccine antigens on their cell surface ([30](#)). A number of observational studies have also reported some cross-protection from 4CMenB (Bexsero®) against gonorrhoea, which is caused by the genetically related *Neisseria gonorrhoeae* ([31](#)).

In 2017, another MenB vaccine, using bi-valent lipided fHbp (rLP2086; Trumenba®; Pfizer), was licensed in Europe. rLP2086 (Trumenba®) is licensed for individuals aged 10 years and older, but has been found to be immunogenic in children as young as one year of age ([32](#)) and without any significant safety or reactogenicity concerns:

- in a study of 220 toddlers aged 1 to less than 2 years of age, the following adverse reactions occurred at a frequency of very common (greater than or equal to 1 out of

- 10): drowsiness, irritability (fussiness), loss of or decreased appetite, fever, and injection site pain, swelling and redness
- in a study of 294 children 2 to 9 years of age, the following adverse reactions occurred at a frequency of very common (greater than or equal to 1 out of 10): headache, diarrhoea, vomiting, muscle pain, joint pain, fever, fatigue, and injection site pain, swelling and redness
 - in clinical studies, fever (greater than or equal to 38°C) occurred more frequently as subject age decreased. Of subjects 1 to less than 2 years of age, 37.3% reported fever; of subjects 2 to 9 years of age, 24.5% reported fever; of subjects 10 to 18 years of age, 9.8% reported fever; and of subjects 18 to 25 years of age, 4.4% reported fever. Fever followed a predictable pattern after vaccination: onset occurred within 2 to 4 days, lasted 1 day, and was mild to moderate in severity. Fever rate and severity tended to decrease with subsequent rLP2086 (Trumenba®) vaccinations
 - adverse reactions following a booster vaccination in 147 subjects 3 to 5 years of age were similar to adverse reactions during the primary rLP2086 (Trumenba®) vaccination series approximately 2 years earlier

rLP2086 (Trumenba®) is licensed to be given as 2 doses (0.5 ml each) administered at a 6-month interval or as 3 doses (2 doses at least 1 month apart with a third dose at least 4 months later). rLP2086 (Trumenba®) has yet to be implemented in a national immunisation schedule.

Importantly, both 4CMenB (Bexsero®) and rLP2086 (Trumenba®) target sub-capsular, surface exposed antigens (rather than the polysaccharide capsule targeted by polysaccharide-conjugate vaccines like MenC and MenACWY vaccines). They have been developed to protect against as many common MenB strains as possible but they do not protect against all MenB strains. Continued awareness and vigilance is, therefore, essential even after completion of vaccine schedules.

4. Additional guidance

National Institute for Health and Care Excellence (NICE) published updated [guidance covering Meningitis \(bacterial\) and meningococcal disease: recognition, diagnosis and management](#) (33).

UK [guidance for universities and other higher education institutions](#) (HEIs) is also available (34).

Refer to the [Sign Grading System 1999 to 2012](#) by the Scottish Intercollegiate Guidelines Network (SIGN) (35) for Table 1 and Table 2 for levels of evidence and gradings of recommendations (see [Appendix 1](#)).

5. Pre-admission management recommendations

Updated NICE guidance was published in March 2024 and should be referred to, the following specific recommendations were made ([33](#)).

1. Transfer people with suspected bacterial meningitis or meningococcal disease to hospital as an emergency.
2. Tell the hospital that a person with suspected bacterial meningitis or meningococcal disease is being transferred and that they will need assessment by a senior clinical decision maker.
3. Do not delay transfer to hospital to give antibiotics to people with suspected or strongly suspected bacterial meningitis or meningococcal disease.
4. If there is likely to be a clinically significant delay in transfer to hospital for people with strongly suspected bacterial meningitis, give intravenous or intramuscular ceftriaxone or benzylpenicillin outside of hospital.
5. For people with strongly suspected meningococcal disease, give intravenous or intramuscular ceftriaxone or benzylpenicillin as soon as possible outside of hospital, unless this will delay transfer to hospital.
6. Do not give antibiotics outside of hospital if the person has severe antibiotic allergy to either ceftriaxone or benzylpenicillin.

Recommendation 1: Pre-admission management (British National Formulary)

Rapid admission to hospital is the highest priority when IMD is suspected.

Evidence grade C

Immediate single dose of IV/IM Ceftriaxone for suspected meningococcal infections. Derived from [Ceftriaxone | Drugs | BNFC | NICE](#):

- adults – dose 2g stat
- children with body weight 50kg and over or aged 9 years and older – dose 2g stat
- children up to 50kg body weight or aged under 9 years – dose 80 to 100 mg/kg (maximum per dose 4g)

Alternatively, immediate single dose of IV/IM benzylpenicillin for suspected meningococcal infections [Benzylpenicillin sodium | Drugs | BNF | NICE](#):

- adults and children aged 10 years or over – dose of 1.2g
- children aged 1 to 9 years – dose of 600mg
- children aged under 1 year – dose of 300mg

All suspected cases of invasive meningococcal disease are statutorily notifiable by registered medical practitioners to the responsible health protection team (see section 7).

6. Laboratory investigation of suspected cases

Patients with suspected meningococcal disease should have appropriate samples taken for bacterial culture, ideally prior to antibiotic administration, as well as polymerase chain reaction (PCR) testing. The source of sampling will depend on clinical presentation and may include blood, cerebrospinal fluid (CSF), joint fluid and/or pleural fluid. A bacterial throat swab should also be taken from all cases with suspected meningococcal disease – a positive meningococcal isolate will provide important information about the infecting strain in PCR-confirmed cases, which may be required for health protection management decisions. If there is a delay in obtaining samples for meningococcal PCR from patients, it may still be possible to retrieve suitable specimens from haematology and/ or biochemistry laboratories. When meningitis is present, CSF is the most appropriate and important sample for confirmatory testing. Whilst it may not be successful, this offers the best chance of yielding an organism for culture and importantly meningococcal DNA can be found in the CSF up to 96 hours after commencing antibiotics ([36](#)). When a patient presents with septicaemia with or without meningitis then blood culture and whole blood (EDTA) are extremely useful to support the diagnosis.

Under the [Health Protection \(Notifications\) Regulations \(2010\)](#), all diagnostic laboratories in England are required to notify UKHSA when they identify specific infections, including *Neisseria meningitidis*. The regulations state that “if the operator of the diagnostic laboratory considers that the case is urgent, the notification must be provided orally as soon as reasonably practicable”. Similar legislation applies in Wales and Scotland. Diagnostic laboratories in Northern Ireland voluntarily notify the Public Health Agency (PHA) when they identify specific infections, including *Neisseria meningitidis*. The Public Health (NI) Act 1967 is currently being updated.

The UKHSA Meningococcal Reference Unit (MRU) offers a national reference service for confirmation and characterisation of invasive meningococcal isolates in England, Wales and Northern Ireland. In Scotland this service is provided by the Bacterial Respiratory Infection Service, Scottish Microbiology Reference Laboratory, Glasgow Royal Infirmary. All invasive meningococcal isolates should be referred to the National Reference Laboratory for confirmation, serogrouping and whole genome sequencing, even if the case has already been confirmed by PCR. The National Reference Laboratories also offer a free service for meningococcal PCR of clinical samples from suspected IMD cases. If IMD is confirmed by a local diagnostic laboratory (including private laboratories), then the original sample, including extracts from PCRs, should be referred to the National Reference Laboratory to allow the capsular group to be confirmed or identified and for additional characterisation. In addition to the routine testing, further strain characterisation may be undertaken in certain situations, such as outbreaks.

Meningococci isolated from a symptomatic urogenital or anorectal infection ([37](#)) or any such isolates (from symptomatic or asymptomatic individuals) considered to be epidemiologically-linked to cases of invasive meningococcal disease should be submitted to the respective national meningococcal reference laboratories for confirmation and further characterisation. All meningococcal isolates from urogenital/ anorectal sites (from symptomatic or asymptomatic individuals) with evidence of antibiotic resistance (penicillin MICs of >0.25 mg/L and/or ciprofloxacin MICs >0.03 mg/L) should similarly be submitted.

To identify and characterise *N. meningitidis*, a combination of traditional and molecular techniques are used. In general, organisms are cultured from blood, CSF or another sterile site. Strain differentiation is performed by the National Reference Laboratory and involves characterisation of capsular polysaccharide and some outer membrane proteins using a monoclonal antibody-based internationally-recognised typing scheme. This allows phenotypic classification by capsular group, type and subtype. The MRU real-time PCR assay also allows capsular group determination for the major disease-causing serogroups (MenA, MenB, MenC, MenW and MenY). Over 50% of IMD cases in England, for example, are now confirmed by PCR only. PorA and fHbp sequencing is also applied to non-culture samples (if there is sufficient DNA) and has the potential for use in cluster/outbreak investigations.

High resolution genotypic analysis (via whole genome sequencing (WGS)) is used to identify genetic relationships between organisms during clusters/outbreaks as they evolve over time and to provide MenB vaccine antigen genotyping. MenB characterisation using Meningococcal Antigen Typing System (MATs), genetic (g)MATs or Meningococcal Deduced Vaccine Antigen Reactivity (MenDeVAR) is used for assessing vaccine strain coverage. WGS and MATs/gMATs/MenDeVAR can be expedited for use in outbreak situations but decisions may need to be made before these are completed in order to achieve protection as early as possible ([38](#), [39](#)).

Detailed descriptions of MRU tests, including specimen types required, can be found in the [Meningococcal Reference Unit \(MRU\) user manual](#).

In Scotland, the user manual is available at [Scottish Microbiology Reference Laboratories - NHSGGC](#).

Recommendation 2: Laboratory investigation

The following specimens should be collected on, or soon after, admission to hospital from all patients (ideally before initiating antibiotics) when meningococcal infection is considered in the differential diagnosis:

- blood for culture
- blood for PCR (ideally EDTA or, alternatively, other unclotted blood specimen)

- if meningitis suspected, CSF for microscopy, culture and PCR - lumbar puncture should not be done if it is contraindicated but can be delayed until it is safe to perform
- for other localised infections, aspirate from sterile site according to clinical indication (for example, joints) for microscopy, culture, PCR
- nasopharyngeal (throat) swab for meningococcal culture (**all suspected cases**); a positive meningococcal swab should **not** be used to diagnose meningococcal disease (for example, pneumonia). However, in PCR-confirmed cases, a positive nasopharyngeal swab culture provides important information about the infecting strain for the purpose of public health management and should, therefore, be submitted to the National Reference Laboratory for additional characterisation

Important: All meningococcal-positive clinical materials (including isolates, PCR-positive clinical samples and/or DNA extracts) should be referred to the National Reference Laboratory for confirmation, serogrouping and further characterisation, even if the case has already been confirmed locally.

Where laboratory confirmation is made using a point-of-care testing platform (for example, Biofire, QiaSTAT), it is important to ensure any residual positive sample is forwarded to the National Reference Laboratory for genogrouping and further characterisation.

As part of enhanced national surveillance, UKHSA previously requested acute and convalescent blood samples from English and Welsh patients with confirmed IMD to assess disease and vaccine responses. **These samples are no longer required.**

In the event of a sudden unexpected death where IMD is suspected and samples were not obtained pre-mortem, post mortem samples referred to the National Reference Laboratory may be helpful in establishing a possible cause of death. Since administering antibiotics prior to taking microbiological specimens is likely to yield negative bacterial cultures from invasive samples, additional investigations such as molecular testing, including PCR testing for IMD and other infectious diseases, should be routinely performed and throat swabs collected ([40](#)).

Laboratory investigations in Scotland:

For children and adults in Scotland a multiplex RT-PCR is available for the detection of *N. meningitidis* DNA from CSF, serum or whole blood (EDTA or other unclotted sample), or any fluid from a normally sterile site, such as joint aspirate. [Sample submission forms](#) are available to download.

Important note: other investigations should be performed according to clinical indication.

Invasive meningococcal cases due to rare serogroups, recurrent disease, with strong family history or that arise after conjugate vaccination (see section 8.5)

IMD cases with a strong family history of IMD, recurrent IMD due to any serogroup, and/ or **IMD in a case aged under 25 years** due to a rare serogroup (including MenY cases or non-encapsulated meningococci (non-serogroupable)) are unusual. Additional investigations should therefore be strongly considered, especially for presence of spleen, splenic function, complement deficiency and HIV testing. IMD due to the respective serogroup after MenC or MenACWY conjugate vaccination is rare and should be similarly assessed for possible underlying conditions. Such cases should be discussed with the UKHSA national immunisation and meningococcal reference laboratory team. A template letter for the GP is available and is used by the UKHSA national immunisation team for follow-up where appropriate ([Meningococcal public health communication templates](#) are available to download).

7. Role of public health

Health protection teams have a major role in the management of meningococcal disease by ensuring that there are adequate disease prevention and surveillance programmes in place, preventing secondary cases through contact tracing, and rapidly investigating and managing clusters and outbreaks (defined in focus 3 box).

Meningococcal meningitis and septicaemia are statutorily notifiable by registered medical practitioners under the [Health Protection Legislation \(2010\)](#), under Health Protection (Wales) Regulations (2010) upon suspicion of meningitis (acute) and meningococcal disease, under Scottish legislation as meningococcal infection and under Northern Ireland as meningococcal septicaemia or acute meningitis (bacterial). Clinicians should inform the proper officer, usually an experienced member of the local health protection team, as soon as a case of meningococcal disease is **suspected** in a patient so that appropriate public health assessment and actions can be undertaken.

In England, Scotland and Northern Ireland, health protection teams should enter all the details of the reported cases on the appropriate national web-based software for public health management of infectious diseases, including any public health actions taken. In Wales, the All Wales Acute Response (AWARe) Team (Health Protection) will enter information onto the equivalent case and incident management system (Tarian).

England

To notify a possible, probable or confirmed case in England, contact your [local HPT](#).

Scotland

In Scotland an overview can be accessed at [Overview: notifiable diseases, health risk states and infections](#).

Find Scotland health protection team details at [Health protection team contacts: general enquiries](#).

Wales

To notify a possible, probable or confirmed case in Wales: during office hours (Monday to Friday 9am to 5pm) or outside of office hours (Monday to Friday 5pm to 9am; weekends and bank holidays) please contact the All Wales Acute Response (AWARe) Team on 0300 00 300 32.

Northern Ireland

To notify a case in Northern Ireland please contact the Public Health Agency in office hours on 0300 555 0119 and out of hours on 028 90404090 (ambulance control) and request to speak with the public health specialist on call.

Recommendation 3: Role of public health

Health protection teams should ensure that policies are in place, ideally through a mechanism such as a service level agreement, which recognises the corporate responsibility of the NHS. Policies should ensure that:

- cases are referred early to hospital
- cases are reported promptly to the health protection team
- cases in hospital are investigated appropriately
- contacts are traced and given appropriate information
- appropriate chemoprophylaxis and vaccination is accessible
- information can be cascaded to others, as appropriate, including primary care, schools or universities, education authorities, National Health Service helplines, meningitis charities, employers
- communication with the media is appropriate and efficient giving due consideration to case (and family) confidentiality

All cases where IMD is suspected should be promptly notified by clinicians to the health protection team, without waiting for microbiological confirmation. Important note: notification is a legal requirement.

Evidence grade D

An experienced member of the health protection team should ensure that comprehensive information on cases is gathered to contribute to local public health management and surveillance with details set out in the [national enhanced surveillance](#) of vaccination programmes targeting invasive meningococcal disease in England. The data set should include epidemiological, laboratory and clinical information. This should be recorded on HPZone/ Case and Incident Management System (CIMS)ⁱⁱ (England, Scotland and Northern Ireland) or Tarian (Wales).

For confirmed cases, the UKHSA national surveillance form (MENSVO1) detailed in the [national enhanced surveillance for meningococcal disease in England](#) or local equivalent covering the same detail should be completed and uploaded to CIMS/ HPZone.ⁱⁱ

[Surveillance forms](#) for Scotland are available to download. In Wales, the national enhanced surveillance form should be completed for all confirmed cases. No additional surveillance forms require completion in Northern Ireland.

Data for local management and audit programmes should include:

ⁱⁱ HPZone was replaced with Case and Incident Management System (CIMS) from May 2024 in England.

Case

- name and address including post code, telephone number, details of general practitioner
- dates and times of disease onset, hospital admission or reporting
- ethnic group
- occupation or workplace
- university, school, college or nursery attended
- meningococcal vaccination history
- antibiotics given prior to admission
- name of hospital or ward
- name of consultant
- specimens and dates and types of specimens
- recent travel history and underlying risk factors (asplenia, splenic dysfunction, complement deficiency, HIV status)

Contacts

- addresses and telephone numbers
- details of antibiotics, vaccine or information given and by whom
- details of GP

Notifier

- name, address and occupation

8. Public health action after a case

8.1 Risk to close contacts

Around 97% of cases are sporadic ([41](#)). Although the risk to contacts is low, the highest absolute and relative risk is to people who live in the same household as a case of meningococcal disease ([41](#), [42](#), [36](#)). This risk is highest in the first 7 days after a case and falls rapidly after this period ([41](#)). The absolute risk (AR) of developing a second case of IMD within 30 days of an index case is 1 in 300 if chemoprophylaxis is not administered ([8](#)). Beyond this period, the risk of meningococcal disease among household contacts is near background levels ([41](#)), although later secondary cases have been observed. The increased risk to household members is thought to be due to shared exposure to meningococci in the contact group, although environmental factors and genetic susceptibility in the family may contribute.

The rationale for giving antibiotic chemoprophylaxis to close contacts of IMD cases is to eliminate established carriage from the close contact group and, thereby, to reduce onward transmission. This strategy reduces the risk of secondary cases in household contacts by up to 89% (95% CI, 42 to 98%) ([43](#)). In such circumstances, the number needed to treat (NNT) – that is, the number of close contacts receiving chemoprophylaxis to prevent one IMD case – is estimated to be 218 (95% CI, 121 to 1,135) ([5](#)).

The case is likely to have acquired the invasive strain from a close contact, typically in the same household, who is an asymptomatic carrier ([44](#), [45](#)). The incubation period is usually 3 to 5 days ([11](#), [13](#)) and cases do not usually have detectable carriage until admission to hospital or shortly beforehand ([13](#)). As the highest risk of IMD in close contacts is in the first 48 hours after disease onset in the index case ([42](#)) the source of infection in secondary cases is most likely to be from the same (or another) carrier and not from the index case.

Antibiotic chemoprophylaxis also eradicates carriage in those who have newly acquired the invasive strain and who may themselves be at risk of IMD. In this instance, individuals who have prolonged close contact with the case after the onset of illness but before the case is treated with antibiotics would also benefit from antibiotic chemoprophylaxis.

It follows that transient contact with the index case before acute illness is unlikely to be a significant risk factor for disease; therefore, mere proximity to the case (for example during travel in a plane, bus or car) does not justify prophylaxis. Although European Centre for Disease Prevention and Control (ECDC) guidance indicates flight contact tracing only where there has been intense exposure to nasopharyngeal secretions, the value of this may be limited once the individuals have dispersed ([46](#)).

Whilst the US guidance recommends that passengers seated next to the index case on a plane for more than 8 hours should be offered prophylaxis, only one such possible on-board

transmission was detected in a review by ECDC with onset of symptoms 2 and 5 days after landing in 2 passengers who had sat 12 rows apart. A further possible on-board transmission was identified during an international Scout outbreak where a Japanese couple developed symptoms 3 to 4 days after the flight. In both these scenarios, it is unlikely that the ECDC-recommended post-exposure prophylaxis (contact tracing for those sitting next to and/or directly exposed to oral secretions of the index case) would have prevented these cases ([46](#)).

As the source of the meningococcal infection may sometimes be outside of the defined population – be it a household or a school – it may not be possible to prevent the meningococci from re-entering the group and, consequently, leading to additional cases in that group. Vaccination of close contacts of an index case can help prevent secondary IMD cases occurring more than 14 days after disease onset in the index case (see section 8.5: Aim of vaccination) ([47](#)).

Focus 1. Case definitions

The following situations require public health action:

Confirmed case

Clinical diagnosis of meningitis, septicaemia or other disease (for example, orbital cellulitis, septic arthritis, pneumonia) and at least one of:

- *Neisseria meningitidis* isolated from a normally sterile site
- Gram-negative diplococci identified in a normally sterile site
- meningococcal DNA in a normally sterile site
- meningococcal antigen in blood, CSF or urine

Although not meeting the definition of a confirmed case, **meningococcal infection of the conjunctiva** is considered an indication for public health action (including treatment for the case and antibiotic prophylaxis for close contacts, but not vaccination) because of the high immediate risk of invasive disease.

Probable case

Clinical diagnosis of meningitis or septicaemia or other invasive disease where an experienced member of the health protection team, in consultation with the physician and/or microbiologist, considers that meningococcal infection is the most likely diagnosis (see: [Focus 3](#) for sources of materials on characteristic symptoms and signs of meningococcal disease in different age groups).

Some microbiological tests (for example, rising antibody levels) that are not considered sufficient to confirm the diagnosis may change the case category from 'possible' to 'probable'.

For cases of meningitis or septicaemia with clinical and laboratory evidence of bacterial infection but where the causative pathogen is not known, meningococcal disease should be considered in the differential diagnosis, especially in previously healthy children and young adults. Where *Neisseria meningitidis* could be responsible and there is no alternative diagnosis at that time, such cases should be considered as 'probable' IMD

Focus 2. Definition of close contacts

Close contact

Meningococcal bacteria that are usually carried harmlessly in the back of the nose and throat with transmission most likely to occur after close and prolonged contact or after sexual or other intimate contact that enables the spread of the bacteria. Close contact is defined as prolonged close contact with the case in a household type setting during the 7 days before onset of illness. Examples of such contacts would be those living and/or sleeping in the same household, pupils in the same dormitory, boy or girlfriends, or university students sharing a kitchen in a hall of residence ([evidence grade B](#))

The definition of close contact does not include ([evidence grade C](#)):

- staff and children attending same nursery or crèche (note – see caveat below)
- students or pupils in same school, class or tutor group (note – see caveat below)
- work or school colleagues
- friends
- residents of nursing or residential homes
- kissing on cheek or mouth (intimate kissing would be considered close, prolonged contact)
- food or drink sharing or similar low level of salivary contact including shared vapes
- attending the same social function
- travelling in next seat on same plane, train, bus, or car (in the absence of intense exposure to nasopharyngeal secretions – see section 8.1)

This list is not exhaustive and should be considered in context for any individuals when prolonged contact has taken place, for example, with multiple episodes of shared vaping, drinking or straw sharing in an enclosed and intimate setting like a club.

Smoking and exposure to second-hand smoke increase the risk of developing invasive disease after acquisition of the bacteria. Where smoking apparatus is shared in an enclosed and intimate setting over a prolonged period, as often occurs with a communal shisha, the risk of transmission is increased, likely due to associated coughing and effects of the smoke on the epithelium of the respiratory tract rather than bacterial transfer via the apparatus itself ([48](#)).

Sharing of a communal shisha may not in itself identify close contacts but the duration, frequency and intimacy of the setting are important factors to be considered.

Caveat: contact in an educational setting

Educational settings include pre-schools, primary schools, secondary schools, colleges and universities. The term 'pre-school' is used synonymously with child-minders, playgroup, nursery, day care or crèche. Within an educational setting, however, it may be possible to define a group that fulfils the definition of a close contact (for example, in a child-minder setting) and, therefore, have a higher risk of developing secondary IMD. Such groups might benefit from public health action ([evidence grade D](#)).

Situations that do not require public health action:

Possible case

Clinical diagnosis of meningitis or septicaemia or other invasive disease where an experienced member of the health protection team, in consultation with the clinician and microbiologist, considers that another diagnosis, such as a viral illness, is more likely than IMD. Information dissemination after a possible case may still be considered (see Recommendation 8).

Isolation of meningococci from non-sterile sites

- isolation of meningococci from sputum, nasopharynx or bronchoalveolar lavage is not by itself an indication for public health action because asymptomatic carriage is common
- non-bacteraemic meningococcal pneumonia, is not an indication for public health action but may carry a small risk of transmission in healthcare settings, especially to the immunocompromised ([49](#), [50](#))

Where meningococci are isolated from a symptomatic urogenital/ anorectal infection, standard treatment for gonorrhoea or non-specific urethritis is expected to clear the meningococci. In asymptomatic individuals, treatment is not recommended because asymptomatic carriage is common

additionally, no further public health management of contacts of individuals with symptomatic or asymptomatic urogenital/anorectal meningococcal infection is required, but isolates should be referred to the meningococcal reference laboratories for confirmation and further characterisation

Contacts that do not meet the close contact definition

After a single IMD case, the risk of additional linked IMD cases outside of the close contact group is low because of the low likelihood of exposure to the responsible strain ([45](#)). In England and Wales during 1995 to 2001, after one case in a pre-school group, a primary school or a

secondary school, the absolute risks to each child or pupil in the same institution of becoming a case within the next 28 days were approximately one in 1,500, one in 18,000 and one in 33,000, respectively (7).

Antibiotic prophylaxis of other contacts is not recommended in educational settings after a single case because the benefits in this setting are largely unknown. The potential for risk reduction is limited by the interval between disease confirmation in the case and time to antibiotic administration within the institution; moreover, harm may arise from drug side-effects, development of antibiotic resistance, and eradication of naturally immunising strains from the nasopharynx. This particularly applies in young children who are more likely to be carrying the commensal *Neisseria lactamica* than *Neisseria meningitidis* (51).

Reports of clusters/outbreaks in other settings (for example, the workplace) are rare and the level of risk is considered to be much lower than in educational settings. As explained previously, transient contact with the index case before acute illness is unlikely to be a significant risk factor for disease, so that mere proximity to the case may not justify prophylaxis. Low-level salivary contact is also not considered to be a risk factor (52). No cases have been reported following post-mortem contact with an IMD case. Embalming is not considered a hazard for transmission.

For recommendations following a case in a healthcare worker see section 9 (Recommendation 9).

8.2 Risk reduction through chemoprophylaxis

A 2014 systematic review suggested an 84% reduction in the risk of subsequent cases of IMD within 30 days, among household contacts given chemoprophylaxis, with 200 household contacts needing to be treated to prevent one subsequent case of IMD within 30 days (53). A review of retrospective observational studies found a significantly reduced risk of additional cases in the household during the month after a case among household members given rifampicin prophylaxis (43). Two randomised controlled trials found no difference in the protection afforded by ciprofloxacin compared to rifampicin (54). In relatively small studies, a single dose of intramuscular ceftriaxone was more effective in eradicating pharyngeal carriage than 4 doses of rifampicin over 2 days, while other studies found oral cefixime and azithromycin to be as effective as rifampicin (55, 56, 57).

In an ECDC review (58), rifampicin, ciprofloxacin, ceftriaxone, cefixime and azithromycin were all recommended for preventing secondary cases of meningococcal disease. In the UK, ciprofloxacin is the recommended chemoprophylaxis of choice and rifampicin is a suitable alternative (59). Ceftriaxone must be given by injection.

In the past, ciprofloxacin was not recommended in children due to induced arthropathy in juvenile animals, but abundant evidence of lack of joint damage has been found in young children given ciprofloxacin. In one RCT on carriage eradication, ciprofloxacin when compared to rifampicin did not lead to a higher rate of side effects ([60](#)). Multiple controlled prospective and retrospective studies, using higher doses of ciprofloxacin, showed that the rate of adverse events of ciprofloxacin in children was similar to that seen using other antibiotics, and that long-term cartilage damage was not seen in humans ([61](#), [62](#)). In all studies, the risk of arthropathy due to ciprofloxacin was very low; arthralgia was transient and most cases were coincidental. A controlled study of 116 neonates receiving ciprofloxacin also showed similar clinical growth compared to 100 controls, even at one year of follow-up ([63](#)). The risk of tendon disorders in a large retrospective study involving 4,531 children given ciprofloxacin was similarly low compared to children given azithromycin (0.8%) ([64](#)). In all studies, side effects resolved after cessation of therapy.

Recommendation 4: Indications for antibiotic prophylaxis

Prophylaxis indicated

Chemoprophylaxis should be offered to close contacts, irrespective of vaccination status, of cases that require public health action (see case definitions [Focus 1](#)) in the following categories:

- those who have had prolonged close contact with the case (including conjunctivitis) during the 7 days before onset of illness (See [Focus 2](#) for definitions)
- those who have had transient close contact with a case only if they have been directly exposed to large particle droplets or secretions from the respiratory tract of a case around the time of admission to hospital

Rapid administration of chemoprophylaxis, ideally as soon as possible after IMD is suspected/confirmed in a case, is key to minimising any imminent risk of meningococcal disease in close contacts. To optimise early chemoprophylaxis, HPTs should first work with hospital clinicians as a matter of urgency to arrange the necessary prescriptions for antibiotic chemoprophylaxis for close contacts, especially for those who are physically present in the hospital with the case (for example, parents of a childhood case) and ideally for all household contacts (especially children) who may be at increased risk of secondary IMD ([Clinical response to local incidents and outbreaks of infectious disease: Commissioning guidance for ICBs March 2025](#)). Alternatively, if it is not possible to arrange urgent antibiotic prophylaxis for close contacts through the hospital, then arrangements should be made through primary care as a matter of urgency (for example, GP out of hours, NHS 111).

Note that antibiotic chemoprophylaxis should not be delayed for young children because of lack of availability or difficulties in obtaining liquid ciprofloxacin suspension in the community. If liquid ciprofloxacin suspension is not available in a timely manner, then crushed tablets may be a suitable alternative for young children. Note that children as young as 3 to 4 years of age can

learn to swallow small tablets and capsules^{iii,iv} with most children being able to [swallow tablets or capsules from 10 years of age](#). Although not ideal, for those who are unable to swallow tablets, ciprofloxacin tablets (except for extended-release forms) can be crushed and mixed with small amounts of food (such as jam, or yoghurt) or liquid (such as water or juice) for ingestion^{v,vi,vii,viii,ix}, to reduce any possible bitter or unpleasant taste.

[Evidence grade B](#)

Prophylaxis for the case

Cases treated with intravenous or intramuscular cephalosporins (for example, ceftriaxone, cefotaxime) do not require antibiotic chemoprophylaxis ([65](#), [66](#)). If the case (including conjunctivitis cases) is treated with any other antibiotic, chemoprophylaxis should be offered when the case is able to take oral medication and, ideally, before discharge from hospital.

[Evidence grade C](#)

Prophylaxis uncertain

The division between those who do and do not receive prophylaxis can be arbitrary as evidence on risk and benefit is limited outside of the household setting. The health protection team will need to use their judgement to decide whether or not to advise prophylaxis for those who do not clearly fall into the close contact or excluded categories in [Focus 2](#). For example, when a case occurs in a group of children looked after by the same child minder or among a circle of close friends, an assessment should be made as to whether these exposures meet the definitions of a close contact.

Timing

Antibiotic prophylaxis should be given as soon as possible (ideally within 24 hours) after the diagnosis of the index case.

ⁱⁱⁱ Rashed AN, Terry D, Fox A, et al. Feasibility of developing children's Pill School within a UK hospital. *Arch Dis Child* 2021;106:705–708.

^{iv} <https://www.nenc-healthiertogether.nhs.uk/parentscarers/medicine-children/pill-swallowing-kidzmed>

^v [Ciprofloxacin - can you crush Cipro 500 tablets and mix with food?](#)

^{vi} Bauters T, Buyle F, Schelstraete P, Dhooge C. Administration of ciprofloxacin through a nasogastric tube in pediatric oncology and stem cell transplantation patients. *J Oncol Pharm Pract*. 2023;29(2):413-415. doi:10.1177/10781552221147891 [PubMed 36572994]

^{vii} Klang MG. Developing guidance for feeding tube administration of oral medications. *JPEN J Parenter Enteral Nutr*. 2023;47(4):519-540. doi:10.1002/jpen.2490 [PubMed 36847617]

^{viii} Lubart E, Berkovitch M, Leibovitz A, et al. Pharmacokinetics of ciprofloxacin in hospitalized geriatric patients: comparison between nasogastric tube and oral administration. *Ther Drug Monit*. 2013;35(5):653-6. doi:10.1097/FTD.0b013e3182940426 [PubMed 23942542]

^{ix} Mimos O, Binter V, Jacolot A, et al. Pharmacokinetics and absolute bioavailability of ciprofloxacin administered through a nasogastric tube with continuous enteral feeding to critically ill patients. *Intensive Care Med*. 1998;24(10):1047-1051. doi:10.1007/s001340050714 [PubMed 9840238]

Recording of antibiotic administration

When antibiotics are prescribed outside general practice for contacts, the GP practice of each recipient of antibiotic prophylaxis should be informed so that an up-to-date medical record can be retained for their registered patient.

Other situations

Dispersal settings

In settings where close contacts have been identified and where contact has now finished, for example those sleeping in the same room on holiday or at university, attempts should be made to arrange chemoprophylaxis within one week of dispersal if practicable.

Evidence grade D

Post-mortem contact with a case

Prophylaxis is not indicated. Kissing the body is not considered to be a risk. Body bags are not necessary, and transport to other countries for burial or cremation does not pose a risk. There is no restriction on embalming.

Evidence grade D

Contacts of possible cases

Contacts of possible cases do not need prophylaxis unless or until further evidence emerges that changes the diagnostic category to confirmed or probable.

Evidence grade D

Delayed diagnosis or notification

If the health protection team receives a delayed report of the case, close contacts (as defined above) should be offered chemoprophylaxis, and vaccine if appropriate, up to 28 days after onset of illness (low risk of further cases after this period).

Evidence grade D

Cases in contacts who have received prophylaxis

If further cases occur within a group of close contacts in the 28 days after receiving prophylaxis, an alternative agent should be used for repeat prophylaxis.

8.3 Choice of agent for chemoprophylaxis

There have been recent EU-wide restrictions and precautions on the use of systemic fluoroquinolone antibiotics (including ciprofloxacin), due to very rare reports of serious side-effects. In view of the most recent UK Medicines and Healthcare products Regulatory Agency

(MHRA) position on their use in January 2024,^x the current chemoprophylaxis recommendations were reassessed and the risk of complications after a single stat dose of ciprofloxacin for meningococcal prophylaxis, if any, was considered to be extremely small whilst the benefits of preventing secondary cases of meningococcal disease are very high.^{xi}

Ciprofloxacin, therefore, remains the recommended choice for meningococcal chemoprophylaxis because it has a number of advantages over rifampicin (54). It is given as a single dose, does not interact with oral contraceptives, and is more readily available in community pharmacies; it is licensed for this indication in adults. It is contraindicated in cases of known ciprofloxacin hypersensitivity. In meningococcal disease and conjunctivitis cases who have travelled outside UK in the 7 days prior to disease, it is important that antibiotic sensitivity testing is completed. High levels of ciprofloxacin resistance have been reported in Asia and the Middle East (67, 68) (see [Rifampicin and travel](#)). Note that levels of resistance remain low in the UK (69).

Rifampicin is a suitable alternative if ciprofloxacin is contraindicated, not available or if the infecting strain is confirmed or likely to be ciprofloxacin resistant (see [Rifampicin and travel](#)). Disadvantages of rifampicin include; selection of resistant bacterial subpopulations, inhibition of contraceptives, requirement for multiple doses over 2 days and availability usually only from hospital pharmacies. Both products are available in preparations suitable for children.

Although benzylpenicillin suppresses meningococcal growth in the throat, it does not reliably eradicate carriage. Around 5% of cases treated with benzylpenicillin still carry the invasive strain after completing treatment and before discharge from hospital (70, 71, 72). Convalescent cases may then pose a risk to household contacts unless given a course of antibiotic treatment to eradicate carriage.

Recommendation 5: Choice of agent for antibiotic prophylaxis

Ciprofloxacin

Recommended for use in all age groups and in pregnancy.

[Evidence grade B](#)

The administration of ciprofloxacin may rarely be followed by anaphylactic reactions (73, 74). Healthcare staff should give out information sheets that include the risk of side effects^{xii} and be prepared to deal with allergic reactions. Ciprofloxacin can also interact with other drugs but a single dose is unlikely to have a significant effect. It has an unpredictable effect on epilepsy but may be preferable to rifampicin if the patient is on treatment with phenytoin (see notes below).

^x [Fluoroquinolone antibiotics: must now only be prescribed when other commonly recommended antibiotics are inappropriate](#)

^{xi} See [EC final decision Quinolone and fluoroquinolone Article 31 referral](#): PRAC recommends restrictions on use (updated 19 March 2019) page 24, table 11.

^{xii} See [Meningococcal public health communication templates](#).

If contacts can take tablets, then this is the preferred route of administration. For young children, liquid ciprofloxacin suspension is available. However, antibiotic chemoprophylaxis should not be delayed for young children because of lack of availability or difficulties in obtaining liquid ciprofloxacin suspension. Children as young as 3 to 4 years of age can learn to swallow small tablets and capsules, with most children being able to [swallow tablets or capsules from 10 years of age](#). Although not ideal, for those who are unable to swallow tablets, ciprofloxacin tablets (except for extended-release forms) can be crushed and mixed with small amounts of food (such as jam, or yoghurt) or liquid (such as water or juice) for ingestion.^{xiii,vi,vii,viii,ix}

Ciprofloxacin dosage (for one dose) [note 1]

All to be given as a single dose:

adults and children aged 12 years and over	500 mg stat
children aged 5 to 11 years	250 mg stat
children aged 1 to 4 years	125 mg stat
infants under 1 year [note 2]	30 mg/kg to a maximum 125mg stat

[note 1] Ciprofloxacin suspension contains 250 mg/5ml.

[note 2] prescribed off-label.

Rifampicin

For close contacts of IMD cases with confirmed or likely ciprofloxacin resistance, (see section travellers) the priority continues to be use of rifampicin for prophylaxis, even in periods of shortage, given the small number of prophylactic doses required. This is important to eradicate any ciprofloxacin-resistant meningococcal carriage and prevent onward transmission. This is particularly the case for infant contacts aged under one year who should be prioritised for rifampicin prophylaxis as there is no clear alternative.

Alternative to Ciprofloxacin in all age groups.

[Evidence grade B](#)

Rifampicin is contraindicated in the presence of jaundice or known hypersensitivity to rifampicin. Interactions with other drugs, such as anticoagulants, phenytoin, and hormonal contraceptives should be considered. Side effects should be explained including staining of urine and contact lenses. Written information for patients should be supplied with the prescription. This is the responsibility of the prescriber.

^{xiii} [Ciprofloxacin - can you crush Cipro 500 tablets and mix with food?](#)

Rifampicin dosage

All doses below to be given twice daily for 2 days:

Adults and children aged 12 years and over	600mg
Children aged 1 to 11 years	10 mg/kg (maximum dose of 600mg)
Infants (under 12 months of age)	5 mg/kg

Suitable Rifampicin doses in children based on average weight for age are:

0 to 2 months	20 mg (1 ml) [note 1]
3 to 11 months	40 mg (2 ml) [note 1]
1 to 2 years	100 mg (5 ml) [note 1]
3 to 4 years	150 mg (7.5 ml) [note 1]
5 to 6 years	200 mg (10 ml) [note 1]
7 to 12 years	300 mg (as capsule/or syrup)

[note 1] Rifampicin syrup contains 100 mg/5 ml.

If it is not possible to offer rifampicin for meningococcal prophylaxis, please contact the UK Health Security Agency, Colindale, telephone: 020 8200 4400, and ask to speak to the duty doctor for further advice. Alternative antibiotic prophylaxis may include a treatment course of oral azithromycin (10mg/kg once daily for 3 days for children up to 17 years (up to 45kg weight), and 500mg once daily for 3 days for adults or weighing more than 45kg). A treatment course of azithromycin is recommended instead of a single prophylactic dose to ensure eradication of nasopharyngeal carriage given some evidence that a stat dose may not generate sufficient concentrations in the saliva to achieve eradication.^{xiv} Alternatively, a single dose of intramuscular (IM) ceftriaxone (125mg <12 years, 250mg 12+ years) may be considered. In neonates, ceftriaxone is contraindicated in babies with hyperbilirubinaemia, premature neonates ≤ 40 weeks corrected gestational age and those receiving calcium-containing intravenous fluids,^{xv} although the risks associated with a single, prophylactic dose is likely to be very low.

Rifampicin and travel

It is important that recent travel history, including place and dates of travel, are established for all cases of invasive meningococcal disease and meningococcal infection of the conjunctiva reported to HPTs and that this information is recorded on HPZone\CIMS.

Because of increasing reports of ciprofloxacin resistance among invasive meningococcal strains in the Middle East and Asia, rifampicin is recommended for first-line prophylaxis in close contacts of both invasive disease and conjunctivitis cases with recent travel to the Middle East

^{xiv} Francisco Soriano and Violeta Rodriguez-Cerrato. Pharmacodynamic and kinetic basis for the selection of pneumococcal resistance in the upper respiratory tract. *Journal of Antimicrobial Chemotherapy* (2002) 50, Suppl. S2, 51–58. DOI: 10.1093/jac/cdf510

^{xv} <https://bnfc.nice.org.uk/drugs/ceftriaxone/>

or Asia^{xvi}. Rifampicin should also be offered for such cases not treated with cephalosporins. If ciprofloxacin has already been given, then HPTs should determine the antibiotic susceptibilities available from the NHS laboratory ^{xvii} and ensure the strain is referred to the UKHSA Meningococcal Reference Unit, Manchester promptly. If the strain is confirmed as being due to a non-groupable meningococcus and/or if ciprofloxacin resistance is identified, even if they have already received ciprofloxacin chemoprophylaxis, close contacts of both invasive disease and conjunctivitis cases should receive rifampicin prophylaxis.

8.4 Pregnancy and breastfeeding

The safety of antibiotic regimens for chemoprophylaxis in pregnant and lactating women is poorly described. Animal reproduction studies have failed to demonstrate a risk to the foetus but there are no well-controlled studies in pregnant women. Because animal reproductive studies are not always predictive of human response, antibiotic chemoprophylaxis should be used during pregnancy only if clearly needed. A single dose of ciprofloxacin can be used for the prevention of a secondary case in pregnancy [see British National Formulary], because short duration treatment for other indications appears to be safe ([75](#), [76](#), [77](#), [78](#)). Of the alternative antibiotics, rifampicin teratogenicity has been reported in animals receiving high doses, but epidemiological studies have not identified any risk in humans when administered for tuberculosis treatment ([79](#)). Another clinical trial involved 176 pregnant and lactating women, administered ceftriaxone (2g) via the intra-muscular route, and only 5 subjects reported mild side effects; there was, however, no control group ([60](#)). For breastfeeding infants, a systematic review of antibiotic use in lactation considered ciprofloxacin and rifampicin as compatible with breastfeeding; other antibiotics were not studied ([80](#)).

Recommendation 6: Antibiotic chemoprophylaxis for pregnant women

Either ciprofloxacin, ceftriaxone or azithromycin can be used as chemoprophylaxis in pregnancy.

Evidence grade C

Ciprofloxacin has the advantage of being easy to access in the community and in short duration usage appears to be safe in pregnancy. Ceftriaxone can only be given by injection and can be painful. Potential side effects include diarrhoea, allergies, hepatic and blood disorders. A treatment course of oral azithromycin is recommended (500mg once daily for 3 days for adults).

8.5 Aim of vaccination

^{xvii} EUCAST [Guidelines for antimicrobial susceptibility](#).

Meningococcal vaccination is offered to those at close prolonged contact with the index case to reduce the risk of late cases through longer-term direct protection. The risk of late cases may be due to increased exposure to virulent meningococci, to environmental factors or to increased susceptibility in the family. In cases caused by vaccine preventable strains, vaccination would be expected to reduce the long-term risk of disease in close contacts. The estimated number of unimmunised close contacts needed to vaccinate to prevent a case was estimated to be approximately 1,000 cases based on confirmed MenC IMD cases ([47](#)).

For MenB, the numbers needed to vaccinate to prevent a single case are substantially higher because at least 2 doses are required for protection ([81](#)). More importantly, the protein-based MenB vaccines are unlikely to afford adequate protection rapidly enough after a single dose (especially for young children who are at highest risk) and most secondary cases occur within a few days of onset in the index case ([7](#)). MenB vaccines do not cover all MenB strains.

Vaccine is not indicated for those who received chemoprophylaxis for transient contact, in dispersal settings or for close contacts of conjunctivitis cases.

Meningococcal cases provide an opportunity to complete the national vaccination schedule in cases and contacts who are eligible according to current [national recommendations](#) for their age. Health protection teams should ensure that all unimmunised and partially immunised cases receive meningococcal vaccination according to national recommendations for their age (Recommendation 7).

Recommendation 7: Vaccination

Vaccination of the index case

Unimmunised or partially immunised index cases should receive all their missed meningococcal vaccinations according to the nationally recommended schedule for their age cohort and national guidance, when they have recovered from their illness. Individuals who have missed MenACWY vaccine at 13 or 14 years are eligible up to their 25th birthday and children aged under 2 years who have not completed MenB vaccination should be brought up to date. See information available at [Vaccination of individuals with uncertain or incomplete immunisation](#). Fully immunised cases do not require additional vaccination as they are expected to have developed an immune response (but see paragraph below for at-risk cases).

[Evidence grade D](#)

At-risk index cases (for example asplenia, complement-deficiency) who are unimmunised or partially immunised against IMD should be appropriately immunised. Current recommendations include the MenACWY conjugate vaccine (2 doses at least 4 weeks apart if aged less than 1 year; 1 dose after first birthday) and the MenB vaccine, 4CMenB (Bexsero®) (2 doses, 8 weeks apart with a booster at 12 months for children aged less than 1 year-olds; 2 doses 8 weeks apart for 1 to 10 year olds; 2 doses 4 weeks apart for older children and adults).

Evidence Grade C

The importance of daily penicillin prophylaxis, as prescribed by the clinician responsible for their care, should be emphasised for at-risk individuals.

Evidence Grade B

Vaccination of close contacts

MenACWY

For confirmed serogroup A, C, W or Y infections, close contacts of any age should be offered the MenACWY conjugate vaccine, unless they are confirmed to have been immunised against the relevant meningococcal serogroup within the preceding 12 months (2 doses 4 weeks apart if aged less than 1 year; 1 dose after first birthday). For close contacts of MenC IMD cases, another MenC-containing conjugate vaccine (for example Menitorix®, NeisVac®, MenQuadfi®) would be a suitable alternative.

Unknown serogroup in a case with recent travel

Where the serogroup is not known in a case who has travelled abroad in the 7 days before onset (or was diagnosed abroad), close contacts of any age should be offered the MenACWY conjugate vaccine, unless the contacts are confirmed to have been immunised against the relevant meningococcal serogroup within the preceding 12 months (2 doses 4 weeks apart if aged less than one year; one dose after first birthday).

Evidence grade D

MenB

After a single case of confirmed or probable serogroup B infection, vaccination against MenB is not recommended for close contacts, even if the strain is identified as vaccine-preventable.

At-risk close contacts

Eligible at-risk close contacts (for example asplenia, complement-deficiency) who are unimmunised or partially immunised should be appropriately immunised for their age. Current recommendations include the MenACWY conjugate vaccine (2 doses 4 weeks apart if aged less than 1 year; 1 dose after first birthday) and MenB vaccine (2 doses 8 weeks apart with a booster at 12 months for less than 1 year-olds, 2 doses 8 weeks apart for 1 to 10 year-olds and 2 doses 4 weeks for older children and adults).

For confirmed serogroup A, C, W or Y infections, fully immunised at-risk close contacts should be offered the MenACWY conjugate vaccine, unless they have received a MenACWY vaccine in the previous 12 months (2 doses 4 weeks apart if aged less than 1 year; 1 dose after first birthday).

The importance of daily penicillin prophylaxis for at-risk individuals should be emphasised.

Evidence Grade B.

National recommendations for vaccination against meningococcal disease are available in the [Meningococcal: the green book, chapter 22](#) and in the [vaccination of individuals with uncertain or incomplete immunisation status](#).

Other contacts (who do not meet the definition for close contact – see [Focus 2](#))

After a single confirmed or probable IMD case, vaccination is not recommended for other contacts, including those who received chemoprophylaxis for transient contact or in a dispersal setting.

8.5.1 Previously immunised IMD cases

The term 'vaccine failure' should be used cautiously in previously immunised IMD cases. No vaccine is 100% effective and vaccine-induced antibodies will wane with time since vaccination. The duration of protection offered by conjugate vaccines, especially in infants and toddlers, is shorter than originally estimated ([17](#)). The vast majority of children and adults who develop IMD, including those who are immunised with any of the meningococcal vaccines, are healthy and have no underlying medical problems. However, IMD after teenage meningococcal conjugate vaccination (for example Men A, C, W, or Y disease after MenACWY conjugate vaccine) is uncommon and such cases should therefore be assessed for possible underlying risk factors, including asplenia and complement deficiency ([82](#)). HIV – undiagnosed or treated - is also a rare but important risk factor for IMD ([83](#)). Those with 2 or more IMD episodes and those with IMD due to unusual capsular groups are also more likely to have underlying risk factors and should be similarly investigated.

Since 4CMenB (Bexsero®) only protects against 73 to 88% of invasive MenB strains, 4CMenB (Bexsero®) failure can only be confirmed in a fully immunised individual if the responsible isolate is identified to be vaccine covered by MATS (MATS-positive). Sequence based methods (gMATS/MenDeVAR), based on MATS, may serve to indicate probable coverage ([84](#), [85](#)). 4CMenB (Bexsero®) coverage cannot be definitively determined for cases confirmed by PCR only, unless the strain possesses the PorA P1.4.

8.6 Disseminating information

Following a single IMD case, it is important to give out information as early diagnosis and treatment should improve outcome. There is a small but real risk of further linked cases ([41](#)). Vigilance for signs and symptoms among close contacts is important especially in the immediate high-risk period (one week) after a case. It is important that this information makes it clear that there are many meningococcal strains that can cause IMD and current vaccines do not protect against all the strains, therefore awareness of symptoms and signs remains critical. Accurate and timely information should help to limit the spread of false rumours and anxiety and may help early identification should a further case arise.

Recommendation 8: Disseminating information

Leaflets or other information about IMD should be widely available and quickly distributed after reporting of a confirmed or probable case. This may also be helpful after a possible case depending on levels of concern, and is a matter for local judgment. See [Focus 3](#) for useful resources for example free leaflets and [template letters](#) are available to download.

[Evidence grade D](#)

An experienced member of the health protection team should ensure that information about an IMD case is shared with other NHS colleagues and external agencies as necessary. It is important to inform any appropriate general practitioners and out-of-hours services, so that they know what public health action has been taken and to promote early recognition of any further cases. An experienced member of the health protection team may also wish to inform NHS helplines and the meningitis charities.

[Evidence grade D](#)

Cases in educational institutions

Heads of pre-school groups, schools, colleges and universities should be informed when there is a confirmed or probable IMD case in someone attending their institution. With the advice of an experienced member of the health protection team, letters are usually sent to other parents/students to inform them of the situation ([template letters](#) are available to download). It is recommended to inform and seek support for this action from the case or their relatives, as the letters may result in identification of the case. The purpose of the letter is to give information about IMD, assist parents and others in the early detection of the disease, allay anxiety and prevent uninformed rumours. It should also provide serotype-appropriate information on vaccination. If it is a MenB case, for example, it is helpful to highlight that current vaccines do not protect against all MenB infection even if a child has received MenB vaccination as part of the national immunisation programme (aged at least 8 weeks and born from 1 May 2015). It is therefore important to remain vigilant as an individual and as a parent even after vaccination.

The information given should be sufficient to ensure that parents are aware of the situation whilst preserving the confidentiality of the patient. It is usually helpful to explain what public health action has been taken.

If a **possible** case attends an educational institution, consider informing the head of the institution at an early stage. The head will then be in a good position to respond immediately to local concerns and will be able to access advice from the HPTs. Letters to other parents or students may be considered.

See also [Meningitis and septicaemia prevention and management in higher education institutions](#). In Wales, there are contingency plans for communicable disease cases or clusters in educational establishments available to the AWARE Team (Health Protection).

Dispersal

If a case is reported within one week of date of last attendance at the institution, distributing information should be considered where practical. This is consistent with chemoprophylaxis in dispersal settings.

Evidence grade D

9. Chemoprophylaxis in healthcare settings

The risk of IMD in healthcare workers is very low (86). Healthcare workers who were more heavily exposed to nasopharyngeal secretions of cases around the time of admission to hospital were considered to be at higher risk (87, 88, 89, 90). UK guidelines for preventing hospital-acquired infections recommend wearing face masks and eye protection when there is a risk of secretions splashing into face and eyes (88, 89). Laboratory studies suggest that surgical masks can protect the wearer against droplet transmission (91, 92).

Meningococcal pneumonia may carry a low risk of transmission in healthcare settings especially to the immunocompromised (41, 49). Meningococcal pneumonia cannot be diagnosed from a sputum sample or from a nasopharyngeal (throat) swab culture because carriage is relatively common in the community. Diagnosis is usually made after the meningococcus is identified in a normally sterile site (typically, blood) – such cases of bacteraemic pneumonia should be managed as invasive disease.

Recommendation 9: Prophylaxis in healthcare settings

In Scotland please refer to [National Infection Prevention Control Manual for Scotland](#). Healthcare workers should reduce the possibility of exposure to large particle droplets (for example by wearing surgical masks, using closed suction) especially when carrying out airway management procedures, so that chemoprophylaxis is not needed.

[Evidence grade D](#)

Chemoprophylaxis is recommended only for those healthcare workers whose mouth or nose is directly exposed to large particle droplets or secretions from the respiratory tract of a probable or confirmed case of meningococcal disease during acute illness until completed 24 hours of systemic antibiotics. This type of exposure will only occur among healthcare staff who are working close to the face of the case without wearing a mask or other mechanical protection. In practice this implies a clear perception of facial contact with droplets or secretions and is unlikely to occur unless using suction during airway management, inserting an airway, intubating, or if the patient coughs in your face. General medical or nursing care of cases is not an indication for prophylaxis. Ciprofloxacin 500mg as a single dose (or, alternatively, rifampicin 600mg orally twice daily for 2 days) is recommended for prophylaxis.

[Evidence grade D](#)

Exposure of the eyes to respiratory droplets is not considered an indication for prophylaxis. Such exposure may, however, carry a low risk of meningococcal conjunctivitis and subsequent invasive disease. Staff should be counselled about this risk and advised to seek early treatment if conjunctivitis should develop within 10 days of exposure.

Evidence grade D

Routine vaccination of healthcare workers with meningococcal vaccines is not recommended because the exposure is invariably transient and those at increased risk will be offered chemoprophylaxis.

Evidence grade D

Vaccination after contact with a confirmed or probable IMD case is also not recommended because the exposure is invariably transient and those at increased risk will be offered chemoprophylaxis.

Evidence grade D

The above recommendations also apply to contacts of cases in healthcare workers (including dentists), and to contacts of cases on a hospital ward where the diagnosis is initially unsuspected and not treated with systemic antibiotics. Chemoprophylaxis is not usually indicated for patient or staff contacts of such cases. A hospital ward is not equivalent to a household setting. However, the threshold for giving prophylaxis should be lower for immunocompromised contacts who may be at increased risk of invasive disease. Risk assessment is advised.

Evidence grade D

10. Cluster and outbreak management

Focus 3. Cluster and outbreak definitions

Cluster

A cluster is 2 or more confirmed or probable IMD cases linked by time and place, which have not yet been shown to be caused by different strains.

Outbreak

An outbreak is a cluster with cases that are epidemiologically linked and that are identified as being caused by the same strain.

Note: If there are 2 or more confirmed or probable cases linked in time and place this should trigger a public health review (see main text).

IMD clusters/outbreaks occur most commonly within households. In a systematic review assessing the effectiveness of vaccinating household contacts in addition to chemoprophylaxis in outbreaks caused by Men A, C, W or Y, 6 eligible studies reporting a total of 4,730 primary cases and 30 household clusters/outbreaks with 40 secondary cases were identified ([47](#)). The attack rate using a fixed effects Poisson model for meta-analysis was 1.08 per 1,000 contacts (95% CI, 0.7 to 1.7) in the 14 to 365 days after disease onset in the index case. Using data from the 4 studies with a follow-up period of more than 31 days, the secondary attack rate after chemoprophylaxis was 20 to 90 per 100,000 household contacts ([47](#)). The authors estimated that between 640 and 1,680 household contacts would need vaccinating to prevent a secondary case ([7](#), [47](#)).

IMD clusters/outbreaks can also occur in a variety of community settings, particularly in institutions such as pre-schools, schools and residential halls of colleges/universities. Enhanced national surveillance indicated that there were approximately 16 IMD clusters/outbreaks annually in school settings and a further 3 in pre-school settings in England and Wales during 1995 to 2001 during a period of relatively high incidence ([7](#)). Over the same period (prior to and during the introduction of universal MenC conjugate vaccination), the overall risk of a cluster/outbreaks was similar for MenB and MenC disease. Most MenC cases occurred in secondary schools, while MenB clusters/outbreaks were more common in primary schools ([7](#)).

An increase in the relative risk (RR) and absolute risk (AR) of a cluster/outbreak due to any capsular group following an initial case in these educational settings has been reported, with the risk being highest in pre-school (RR, 27.6; AR, 70 per 100,000) and lowest in secondary school (RR 3.6; AR, 3 per 100,000) settings ([7](#)). In most clusters/outbreaks, secondary cases occurred

within one week of the index case (29% within 2 days, 68% within 7 days) and, by the end of the third week, the RR of a secondary case was similar to baseline. The majority of clusters/outbreaks (89%) had only 2 cases and, where third cases did occur, 93% were diagnosed within 6 days of a secondary case (although, in one cluster/outbreak, a third case occurred 21 days after the second case).

A case-control analysis of school children in the USA (1989 to 1994) estimated the secondary incidence of IMD as 2.5 per 100,000 in school children aged 5-18 years, with relative risk of 2.3 (8). One third of cases occurred within 48 hours of the index case and 75% within 2 weeks. In secondary schools, where 75% of clusters/outbreaks occurred, 73% of secondary cases occurred within 2 weeks of the index case. When more than 2 cases were identified, in school-based clusters/outbreaks, the mean time between second and third cases was 1.6 days (range 0 to 5 days). No attempt was made in these studies to estimate any additional benefit of vaccination over chemoprophylaxis in preventing further cases.

In educational settings, once a second linked case has occurred, the risk of a third case has been reported to be as high as 30 to 50%, with the risk of a third case being highest in the week after the second case (7, 93). These studies were, however, conducted more than 2 decades ago when IMD incidence was higher. Where there is no direct connection between the 2 cases, the risk of a third case remains low and is influenced by meningococcal circulation in the wider community rather than within the educational-setting itself. It is, therefore, important to gather as much information as possible on potential links if one is not immediately apparent.

The risk to staff in educational clusters/outbreaks is not known. However, of 6 clusters/outbreaks that contained confirmed cases among both staff and children in educational settings in England and Wales from 1995 to 2001, 5 involved pre-school groups or primary schools (7), suggesting a greater risk to teachers of young children.

Relative risk of additional cases in other settings has not been formally assessed, but outbreaks in definable social groups, civilian communities and military recruits are well-described (94). Although one trial of mass chemoprophylaxis in a closed community (military barracks) showed a significant effect on disease reduction (95), whether such interventions work in schools or civilian communities is not known (96, 97). The aim of such interventions is to eradicate carriage of the outbreak strain from a population at high IMD risk (98).

If a cluster/outbreak is caused by strains of a serogroup for which an effective vaccine exists, vaccination should be considered. Data from England and Wales showed that if the serogroup of one case had been identified and another case was diagnosed within 4 weeks in the same school, the second case was likely to be of the same strain as the first case (7). In the USA, vaccination of whole communities in community MenC IMD outbreaks is considered when a defined threshold is reached (93).

Assessment of the likely benefits and costs of interventions must then lead to a decision on public health action. External factors such as availability of staff, antibiotics, vaccine and

feasibility of action (such as holidays just started) may well influence the decisions made ([99](#)). More evidence is needed on the effectiveness of such interventions.

10.1 Clusters or outbreaks in a household setting

An IMD household cluster is defined as 2 or more probable or confirmed cases with symptom onset within 28 days in the same household. Household clusters are almost always caused by the same strain and would therefore constitute an outbreak. An outbreak in a family may indicate increased susceptibility of family members to IMD and/or on-going transmission within the household setting. Such outbreaks are rare and occasionally may occur after longer intervals possibly due to re-introduction into the household. Following the introduction of the infant MenB immunisation programme in September 2015, the recommendations for MenB vaccination in a outbreak were revised as set out in Recommendation 10.

Recommendation 10: vaccination following a household outbreak

- all close contacts (including the case) are recommended to receive the appropriate vaccine (MenACWY or MenB) in addition to antibiotic prophylaxis unless they are confirmed to have been immunised against the relevant meningococcal serogroup within the preceding 12 months
- if additional cases occur within 28 days of receiving antibiotic prophylaxis, then an alternative agent should be used for repeat antibiotic prophylaxis unless antibiotic sensitivities are available for the additional case (see Recommendation 4: Indications for antibiotic prophylaxis)

[Evidence Grade D](#)

10.2 Management of clusters/outbreaks in an educational or residential setting

Two or more cases of probable or confirmed IMD occurring with symptom onset within 28 days in an educational or residential setting should be considered as a cluster. Cases in this type of cluster are more likely to constitute an outbreak if a common social or other network can be identified, alongside a close relationship in time and place. The following should be considered:

1. Are the infecting strains indistinguishable?
 - many clusters are likely to be caused the same strain that is circulating in the local community – therefore, identification of indistinguishable strains may not necessarily indicate the need for prophylaxis or vaccination, especially if there are no identified epidemiological links between the cases (see point 2 (below) and ‘Linked cases with no identifiable group for public health action’ (bottom of section 10.2))

- clear differences between strains can, however, rule out a cluster or outbreak and therefore the need for prophylaxis or vaccination
- because of the need for speed, public health action may need to be initiated even if the second case is probable or before full strain characterisation is available for either or both cases, especially if a common social network can be identified

2. Is there a clearly identifiable group at increased risk of IMD that may benefit from public health action such as wider antibiotic prophylaxis and or vaccination?

Evidence suggests that increased risk of a second case arising in an educational setting following the first case persists for up to around 3 weeks ([7](#)). Cases arising more than 30 days apart in an educational setting are most likely to be due to different capsular groups or strains of the meningococcus and, therefore, unrelated. Thus where intervals exceed the 28-day cluster/outbreak definition, it is more than likely to represent 2 separate introductions into the population or indicate that circulation of that particular meningococcal strain is occurring more widely in the local community.

It is known, however, that MenB carriage can persist over a prolonged period. Where strain characterisation of 2 or more IMD cases suggests an identical strain and the cases are linked through a distinct social network or other clearly defined network, therefore, further action may be considered by an incident management team (IMT) even when a period exceeding 28 days has elapsed between cases ([100](#)).

These considerations should be discussed by an IMT and will inform the choice of public health interventions that should be undertaken:

- antibiotic prophylaxis leads to short-term meningococcal clearance and, therefore, offers the greatest benefit if given as soon as possible after an outbreak is defined and the risk group is identified
- vaccination provides longer-term direct protection to individuals in defined outbreak networks. Conjugate vaccines (MenC, MenACWY) can also interrupt transmission of the respective serogroups within a network; the current MenB vaccines, which are protein-based, do not have any impact on carriage ([101](#))

Important note: Irrespective of whether the cases constitute a cluster or an outbreak, early dissemination of information should be undertaken to raise awareness of signs and symptoms of IMD because of the importance of being informed and seeking early medical advice if symptoms arise ([template letters](#) are available to download).

Different educational settings are considered; pre-schools, primary schools, secondary schools and universities. Residential setting includes, for example military barracks, asylum centres, nursing or residential homes. If indicated, vaccination of contacts in the educational or

residential setting should be offered as early as possible because the attack rates are much higher within the first week after the index case is diagnosed.

It may be possible to identify a clearly defined social network where further intervention could be beneficial. If there is no clear group at increased risk of disease, identifying a potential group to be vaccinated becomes difficult. In the absence of a closed or semi-closed network, wider chemoprophylaxis is also unlikely to be beneficial because chemoprophylaxis provides short-term clearance of the nasopharynx but the bacteria can be reintroduced into the network from those outside the network as soon as protection from chemoprophylaxis declines. Decisions related to public health action in such circumstances should generally be made by experienced members of the health protection team or by the IMT.

Protective immune response after conjugate vaccines (MenC or MenACWY) is usually rapid, typically within a week after a single dose of vaccine. Conjugate vaccines also prevent acquisition of carriage, which may help control the spread of infection within the local setting. On the other hand, not all MenB strains in the UK are predicted to be preventable by 4CMenB (Bexsero®) or rLP2086 (Trumenba®) and at least 2 doses are required for protection (4). Additionally, 4CMenB (Bexsero®) and rLP2086 (Trumenba®) do not reduce carriage – this needs to be taken into consideration when assessing the potential benefit of vaccination to control outbreaks.

Recommendation 11: Managing clusters/outbreaks in educational and residential institutions

Expert advice is available for managing clusters/outbreaks from:

- UK Health Security Agency, Colindale, telephone: 020 8200 4400
- AWARe (all Wales Acute Response team) telephone: 0300 00 300 32
- Public Health Scotland, telephone: 0141 300 1100
- Northern Ireland Public Health Agency Health Protection Duty Room (telephone 0300 555 0119) and out-of-hours on 028 90404090 (ambulance control) and request to speak with the public health specialist on call

Please alert the appropriate organisation to any cluster/outbreak situation.

Assess the information

When 2 or more cases are reported in an educational or residential setting, careful and rapid assessment should be made. This should include a review of:

- clinical features of the cases
- microbiological data (serogroup and sequence-based typing)
- dates of onset of illness and of last attendance
- links between cases by age, school year, home address, social activities, and friends

- the type of setting (pre-school, primary school, secondary school or university)
- numbers of students in the school and in each school year

Consider the public health management options

The usual course of action should include dissemination of information to raise awareness of symptoms and signs of IMD because of the need for early medical intervention. Information should be distributed widely using all available platforms to parents and students, as appropriate ([template letters](#) are available to download).

Evidence grade D

The main decision to be taken by an IMT is whether a high-risk group can be identified that might benefit from public health action, including antibiotic prophylaxis and vaccination to reduce that risk.

The target group should be a discrete group that contains the cases and makes sense to staff, parents or students. For example, children and staff of the same preschool group, children of the same school year, children or students who share a common social activity, or a group of friends. The evidence on risk indicates a need to act promptly with the agreed public health action to prevent additional cases.

Evidence grade D

Make a decision on antibiotic prophylaxis

For clusters/outbreaks in an educational or residential setting, if a clear high-risk group can be defined that contains the cases, antibiotic prophylaxis should be offered to that group. If a subgroup cannot be defined, then a decision may be needed on offering prophylaxis to the whole institution. This will depend on factors, for example, such as the size of the population, the time interval and age difference between cases, whether the cases are confirmed or not and any strain information.

For clusters/outbreaks among children at pre-school groups and primary schools, both children and staff should normally be included in the group offered chemoprophylaxis and vaccination (some evidence of increased risk) but not in clusters/outbreaks among students at secondary schools, colleges, or universities (no evidence of increased risk).

Where ciprofloxacin is recommended, patient group directions by [UKHSA](#) and [Public Health Scotland](#), may be helpful.

Evidence grade D

Make a decision on vaccination

Template patient group directions by [UKHSA](#) and Public Health Scotland ([MenACWY](#) and [MenB](#) vaccines) are available for meningococcal vaccines.

For a cluster/outbreak involving confirmed serogroup A, C, W or Y cases: the MenACWY conjugate vaccine should be offered to all individuals of any age who were offered antibiotics unless they are confirmed to have been immunised against the relevant meningococcal serogroup within the preceding 12 months. In the case of a MenC cluster/outbreak, another MenC-containing conjugate vaccine (for example Menitorix®, NeisVac®, MenQuadfi) would be a suitable alternative.

Evidence grade D

For a cluster/outbreak involving confirmed MenB cases: vaccination against MenB should be considered and would usually be offered to the same group that would receive antibiotic chemoprophylaxis, including cases once recovered from their illness, as soon as practically possible based on the schedule table below. However, vaccination should target those in the group identified as potentially being at **ongoing** increased risk of disease; for example, if there are 2 MenB cases in a nursery, then nursery contacts may be offered MenB vaccination, but the household contacts of each case would not be considered as have ongoing increased risk.

Evidence grade D

Choice of MenB vaccine

Two vaccines are licensed against MenB; 4CMenB (Bexsero®) and rLP2086 (Trumenba®). The vaccination dosing and schedule for 4CMenB (Bexsero®), as well as the licensed age indication, is in general more suitable for cluster/outbreak control than rLP2086 (Trumenba®). 4CMenB (Bexsero®) also has proven efficacy in the field. Therefore, until more data become available, 4CMenB (Bexsero®) is the vaccine of choice **unless**:

- there is a case with complete 4CMenB (Bexsero®) vaccination (3 doses) thus indicating the strain is more likely to not be covered by 4CMenB (Bexsero®)

or:

- where the outbreak strain is predicted not to be prevented by 4CMenB (Bexsero®) using for example, MATS (if isolates or assay are available, or gMATS/MenDeVar (if a sequence-based approach is used))

Where additional testing is required, results may not be timely and should not delay public health decisions. Please discuss with the UKHSA Meningococcal Reference Unit or, in Scotland, Public Health Scotland.

[Where 4CMenB \(Bexsero®\) is recommended in a cluster/outbreak situation](#); the 2 doses should be offered with a 4-week interval for those aged one year and older because of the need for early protection which, in these circumstances, would outweigh the need for longer-term protection.

It is important to emphasise to vaccinees or parents of vaccinees that the vaccine will only start protecting around 2 weeks after the second dose and, therefore, they should remain vigilant for symptoms and signs of meningococcal disease. Additionally, if 4CMenB (Bexsero®) is administered without confirmation of potential strain coverage by the vaccine, vaccinees/parents of vaccinees should be made aware that the vaccine may not protect against the circulating strain even after 2 doses.

[Evidence grade D](#)

In a cluster or outbreak situation, the cases of serogroups A, B, C, W and Y IMD should also be vaccinated as part of the social network when the cluster/outbreak is declared, unless they have received the vaccine in the previous 12 months.

[Evidence grade D](#)

In line with Recommendation 7, in a cluster/outbreak situation, vaccine is not indicated for those who received chemoprophylaxis for transient contact in dispersal settings as they would not usually be considered to have ongoing increased risk.

[Evidence grade D](#)

Table of 4CMenB (Bexsero®) vaccination schedule for MenB cases and contacts

Age	4CMenB (Bexsero®) Vaccination Status	Schedule for secondary prevention of MenB disease
Under 8 weeks	Unvaccinated	Vaccinate in accordance with routine vaccination schedule at the appropriate ages
At least 8 weeks and less than 1 year old	Unvaccinated	Give 2 doses 8 weeks apart with a booster at 1 year of age at least 4 weeks after the last 4CMenB (Bexsero®) dose
Less than 1 year old	Vaccinated	Ensure up to date with nationally recommended immunisation schedule. Continue and complete.
1 year and older	Completed course (2 doses under 1 year and a third dose on or after first birthday, or 2 doses after first birthday) and: where the most recent dose was more than 12 months ago	Single dose of 4CMenB (Bexsero®) to boost immunity.
	Completed course (2 doses under 1 year and a third dose on or after first birthday, or 2 doses after first birthday) and: where the most recent dose was less than 12 months ago	No additional vaccination
	Received an incomplete course (see above for definition of complete course) and: the most recent dose was at least 4 weeks ago	Single dose of 4CMenB (Bexsero®) to complete the schedule
	Received one 4CMen B dose in the last 4 weeks	Single 4CMenB (Bexsero®) doses after 4 weeks to complete the schedule
	Not received any prior doses of 4CMenB (Bexsero®) vaccine	Give 2 4CMenB (Bexsero®) doses 4 weeks apart [note 1]

[note 1] There is no accelerated immunisation schedule for 4CMenB (Bexsero®) but the interval between doses for 1 to 10 year olds should be reduced to 4 weeks for secondary prevention of MenB disease because of the need for rapid protection.

[Evidence grade D](#)

Use of rLP2086 (Trumenba®) vaccine in clusters/outbreaks

When there is a fully 4CMenB (Bexsero®) vaccinated case in a child younger than 5 years of age, irrespective of strain coverage, or where the typing profile indicates its unlikely to be covered by Bexsero, then rLP2086 (Trumenba®) at the following schedule is recommended, after discussion with the UKHSA MRU or, in Scotland, Public Health Scotland:

- anyone aged one year or over should receive 2 doses of rLP2086 (Trumenba®) at least 4 weeks apart (irrespective of prior 4CMenB (Bexsero®)), with the first dose given as soon as possible – this is to ensure early and rapid protection against IMD and, as such, the third dose of rLP2086 (Trumenba®) at 6 months is not required
- in children aged one year to less than 2 years, any outstanding 4CMenB (Bexsero®) doses, as recommended according to the national immunisation schedule, should be given at least 4 weeks after the second additional rLP2086 (Trumenba®) dose administered as part of the public health response
- although rLP2086 (Trumenba®) is unlicensed in children aged 1 to 9 years, studies have shown the vaccine is safe and immunogenic in this age group
- infants aged 11 months until their first birthday should receive an extra dose of 4CMenB (Bexsero®) immediately as part of the public health management if they have not had a dose in the previous 4 weeks – they can then be given 2 doses of rLP2086 (Trumenba®) at least 4 weeks apart from their first birthday and at least 4 weeks after their last 4CMenB (Bexsero®) dose
- infants under 11 months should receive an extra dose of 4CMenB (Bexsero®) if they have not had a dose in the previous 4 weeks (rLP2086 (Trumenba®) has not been used in infants under one year of age) – this extra dose should be considered additional to the nationally recommended 2+1 schedule at 8 weeks, 16 weeks and one year of age. Any further 4CMenB (Bexsero®) doses as recommended according to the national immunisation schedule should be given at least 4 weeks after the additional 4CMenB (Bexsero®) dose given as part of the public health response. Maximising 4CMenB (Bexsero®)-induced antibodies with an additional vaccine dose should provide the infant with some additional cross-protection against IMD in the context of a cluster/outbreak

rLP2086 (Trumenba®) is not currently included in the national childhood programme and it may be problematic to purchase a timely supply (see [Emergency vaccine supply for use in an cluster/outbreak situation](#)). If rLP2086 (Trumenba®) is indicated but cannot be made available within 2 weeks of antibiotic chemoprophylaxis administration, then 4CMenB (Bexsero®) should be used according to the [4CMenB \(Bexsero®\) recommended cluster/outbreak schedule](#). This is expected to provide additional cross-protection during the outbreak period.

For a cluster/outbreak involving 2 or more “probable” cases: every attempt should be made to determine the meningococcal capsular group for at least one case before any decision to offer vaccination is made. If this is not possible, then a vaccine with broad coverage such as

4CMenB (Bexsero®) or rLP2086 (Trumenba®) may be considered as these vaccines may also help protect against other capsular groups.

If antibiotics and/or vaccine are to be offered, make urgent arrangements in line with local procedures with:

- community medical or nursing staff to deliver medicines, vaccine or information
- head of the institution to inform parents/students and seek consent
- pharmacists to supply antibiotics (in correct formulation, dosage and information sheets) and vaccines ([102](#))

Important note: Closing the school is not advised as no reduction in risk would be expected (levels of contact among social networks are unlikely to be reduced and may increase with school closure; also, application and success of public health actions will be assisted if school attendance is high).

Swabbing to measure carriage of cluster/outbreak strains is not usually recommended in acute clusters/outbreaks because decisions have to be taken before results are available and because carriage rates often bear no relationship to risk of further cases.

Important note: If 2 or more cases occur within a clearly defined social group outside an educational setting, the same principles as for a school cluster/outbreak apply.

Recording of antibiotic and vaccine administration

When antibiotics are prescribed or vaccinations given outside general practice in a cluster/outbreak setting, the GP practice of each recipient of antibiotic prophylaxis and vaccination (with batch and product information) should be informed so that an up-to-date medical record can be retained for their registered patient.

Emergency vaccine supply for use in an outbreak situation

In England and Wales, if there is no vaccine stock available from the manufacturers or if there is any delay, then discuss with colleagues in the Immunisation Division at UKHSA Colindale or email vaccinesupply@ukhsa.gov.uk for advice.

In Scotland, vaccine supply should be obtained from local hospital pharmacy departments.

Vaccine supply for use in clusters/outbreaks in Northern Ireland should be discussed with the Public Health Agency Health Protection Duty Room 0300 555 0119 or out of hours on 028 90404090 by requesting to speak with the public health specialist on call.

Linked cases with no identifiable group for public health action (see top of section 10.2)

In such circumstances (for example, 2 students at the same university with no contact and no common social links identified), a broad 'warn and inform' approach would be indicated and further public health action will need to be decided based on the specific circumstances.

10.3 Vaccination in pregnancy

In the context of a cluster/outbreak, chemoprophylaxis is recommended in close contacts who are pregnant to reduce their immediate risk of invasive disease. MenACWY conjugate vaccination is also recommended to close contacts of cases with these 4 serogroups (A, C, W, Y), including those who are pregnant, given their low reactogenicity.

MenB vaccination is not advised for close contacts who are pregnant in an outbreak setting because of the lack of data on administration of licensed MenB vaccines during pregnancy and their known reactogenicity profile which includes pyrexia. For those who are pregnant and in a high risk group (such as asplenia or on treatment inhibiting complement activation, such as eculizumab, ravulizumab, or zilucoplan) specialist advice must be sought.

10.4 Management of clusters/outbreaks in the wider community

One of the major difficulties in targeting a wider community for intervention is deciding on the population boundaries, often defined by age group and geography. Such boundaries will of necessity be arbitrary. As far as possible, use existing administrative boundaries that make sense to the people who live within and without them. In any case, there are likely to be people living on the other side of the boundary who may feel unjustifiably excluded. The extent of public concern and press interest can be extensive. There have been examples of extended clusters/outbreaks of disease within socially-related groups over a poorly defined geographical area. Such clusters/outbreaks may be difficult to define as there may be more than one link by recreational activity (for example, sports club) or through regular social groups. In a university setting where cases are not limited to a well-defined student population or student residence, for example, it may be more helpful to consider the university in the context of a community setting.

Although school clusters/outbreaks must be handled quickly in order to control alarm and reduce immediate risk of further cases, wider community clusters/outbreaks usually build up more slowly and by their nature are more diffuse. The same principles and management steps apply (see recommendations 11, 12 and 13).

In such situations, age-specific attack rates should be calculated.

10.4.1 Calculating age-specific attack rates

The numerator would be the number of confirmed cases in the population at risk caused by strains of the same capsular group and that are not distinguishable by standard molecular typing. Multiple cases in the same household or in the same institutional setting would be considered (if this setting is considered to be the focus of a separate cluster/outbreak) as a single case.

The denominator would be the population at risk, which must be clearly defined if meningococcal vaccination is to be offered, and make sense to the people who live within and outside the selected boundaries (for example a rural town or village, a secondary school with its feeder schools). It may not be easy to define such a population. If the cluster/outbreak is mainly in children, the denominator should be based on the age range of children at risk (for example, 2 to 4 year olds, 2 to 16 year olds) in whom the vaccine should be effective.

Vaccine should only be considered if the age-specific attack rate (number of confirmed cases due to the outbreak strain [suggested minimum of 4] divided by the number in target age-group) in a 3-month period is 'high'. Although a precise threshold for intervention has not been set, age-specific attack rates among 2 to 16 year olds targeted for intervention in 2 community outbreaks during the winter of 1995 and 1996 caused by MenC were greater than 40 per 100,000.

Recommendation 12: Managing clusters/outbreaks in the wider community

Any decision to offer meningococcal vaccines to wider communities will require careful assessment of all the available epidemiological information, such as the number of confirmed and probable cases, molecular information on infecting meningococcal strains, dates of onset, links between cases, size of the community, and routine vaccination uptake rates.

Evidence grade D

Vaccination against clusters/outbreaks caused by the same serogroup of IMD may be considered in the community if the age-specific attack-rate (for a vaccine preventable strain in the case of MenB) within a defined geographical boundary over a 3 month period exceeds 40 per 100,000.

10.5 Disseminating information in cluster/outbreak management

It is essential that clear, consistent and accurate information is provided to parents, students and staff, and the wider community. The target group should be clearly identified and information to this group should emphasise the importance of early recognition of symptoms and prompt access to medical services ([template letters](#) are available to download).

Local general practitioners and out-of-hours services should be advised to be on the alert for any new cases associated with the cluster/outbreak. It may also be helpful to alert receiving Accident and Emergency departments and admitting clinicians.

As far as possible, information that may need to be disseminated should be prepared in advance. In pre-school and school settings an experienced member of the health protection team should liaise closely with the manager or head teacher. In college or university settings liaison will usually be with a member of the senior management team (see [higher education institution guidance](#)) and representation from the educational setting would usually be invited to join the IMT. It is advisable for one person within the college/university to coordinate operations, and to receive and disseminate all information. Registry departments can aid in tracing students and getting information to them, and personnel or occupational health departments can help disseminate information to staff groups. In Wales there are institution-specific contingency plans for educational settings available to the AWARe team and Out Of Hours teams.

A communication strategy will be required. If high levels of interest are anticipated or already evident, consider; telephone helplines (see Focus 4 for helpline contact details), allowing controlled media access to vaccination sites, and regular coordinated press briefings and to hold press conferences ([83](#)).

Focus 4. Helplines and leaflets

Meningitis charities and NHS111, NHS 24, NHS Inform

The meningitis charities may be contacted when there is a case of meningococcal disease. They need to have sufficient information so that they can support callers with appropriate advice. The information given to these bodies should include anonymised details of the case and of public health action taken.

NHS inform is Scotland's national health information service [Scottish health information you can trust | NHS inform](#).

Leaflets and posters available from Health and Social Care Publications orderline

Ordering from the Health and social care order line for England is easy and the service is free of charge. Anyone can [register for an account](#). Once you have registered you will then be allocated an account and can place orders. You will need your full postal address and an email address.

In Scotland, if you wish to order vaccine leaflets or posters, please contact your local Health Board Resources Officer who can place an order via the online portal. Vaccine leaflets can also be ordered directly from Public Health Scotland by emailing p hs.generalpublications@p hs.scot

Meningitis Research Foundation

For enquiries and to request resources: 080 8800 3344 (UK) or 1800 41 33 44 (Republic of Ireland)

Helplines

Meningitis Now nurse-led Helpline 0808 80 10 388 (Freephone) email: helpline@meningitisnow.org and on social media [Facebook](#) [@meningitis_now](#)

[Leaflets and posters](#) can be ordered or downloaded from the website or by calling the Meningitis Now office.

Meningitis Research Foundation 0808 800 3344 in the UK (Freefone), 9am to 5pm Monday to Friday. One-to-one advice and support is available on the phone and via online chat, email and social media: meningitis.org/get-support/get-support.

Help and support resources can be accessed at any time at meningitis.org/get-support/help-and-support-resources.

- NHS 111 (England)
- NHS 24 (Scotland) Dial 111
- NHS Direct Wales 0845 46 47

Websites

- [Meningitis Research Foundation](#)
- [Meningitis Now](#)
- [Meningitis on NHS.UK](#)
- [Meningococcal disease \(UKHSA\)](#)
- [Public Health Agency, Northern Ireland](#)
- [Meningococcal disease](#), [Public Health Scotland](#)
- [Scotland NHS Inform](#)
- [Immunisation Scotland](#)
- [Meningococcal disease](#), Public Health Wales
- [Meningococcal green book chapter 22](#)

11. Appendix 1

Table 1. Levels of evidence ([35](#))

1++	High quality meta analyses, systematic reviews of RCTs, or RCTs with a very low risk of bias.
1+	Well conducted meta analyses, systematic reviews of RCTs, or RCTs with a low risk of bias.
1-	Meta analyses, systematic reviews of RCTs, or RCTs with a high risk of bias.
2++	High quality systematic reviews of case-control or cohort studies. High quality case-control or cohort studies with a very low risk of confounding, bias, or chance and a high probability that the relationship is causal.
2+	Well conducted case control or cohort studies with a low risk of confounding, bias, or chance and a moderate probability that the relationship is causal.
2-	Case control or cohort studies with a high risk of confounding, bias, or chance and a significant risk that the relationship is not causal.
3	Non-analytic studies, for example case reports, case series.
4	Expert opinion.

Table 2. Grades of recommendation

A	At least one meta-analysis, systematic review, or RCT rated as 1++, and directly applicable to the target population; or a systematic review of RCTs or a body of evidence consisting principally of studies rated as 1+, directly applicable to the target population, and demonstrating overall consistency of results.
B	A body of evidence including studies rated as 2++, directly applicable to the target population, and demonstrating overall consistency of results; or extrapolated evidence from studies rated as 1++ or 1+.
C	A body of evidence including studies rated as 2+, directly applicable to the target population and demonstrating overall consistency of results; or extrapolated evidence from studies rated as 2++.
D	Evidence level 3 or 4; or extrapolated evidence from studies rated as 2+.

References

1. Halperin SA, Bettinger JA, Greenwood B, Harrison LH, Jelfs J, Ladhani SN and others. 'The changing and dynamic epidemiology of meningococcal disease' Vaccine 2011: volume 30, supplement 2 (1873 to 2518 (Electronic)) B26 to B36
2. Christensen H, May M, Bowen L, Hickman M, Trotter CL. 'Meningococcal carriage by age: a systematic review and meta-analysis' Lancet Infectious Diseases 2010: volume 10, issue 12, pages 853 to 861
3. MacLennan JM, Rodrigues CMC, Bratcher HB, Lekshmi A, Finn A, Oliver J, Wootton M, Ray S, Cameron C, Smith A, Heath PT, Bartolf A, Nolan T, Hughes S, Varghese A, Snape MD, Sewell R, Cunningham R, Stolton A, Kay C, Palmer K, Baxter D, Suggitt D, Zipitis CS, Pemberton N, Jolley KA, Bray JE, Harrison OB, Ladhani SN, Pollard AJ, Borrow R, Gray SJ, Trotter C, Maiden MCJ. 'Meningococcal carriage in periods of high and low invasive meningococcal disease incidence in the UK: comparison of UKMenCar1-4 cross-sectional survey results' Lancet Infectious Diseases 2021: volume 21, issue 5, pages 677 to 687. Epub 2021 January 19. Erratum in: Lancet Infectious Diseases May 2021 May: volume 21, issue 5, e122. Erratum in: Lancet Infectious Diseases 2021: volume 21, issue 7, e182
4. Trotter CL, Gay NJ. 'Analysis of longitudinal bacterial carriage studies accounting for sensitivity of swabbing: an application to *Neisseria meningitidis*' Epidemiological Infections 2003: volume 130, issue 2, pages 201 to 205
5. Ladhani SN, Cordery R, Mandal S, Christensen H, Campbell H, Borrow R and others. 'Preventing secondary cases of invasive meningococcal capsular group B (MenB) disease using a recently-licensed, multi-component, protein-based vaccine (Bexsero®)' The Journal of Infection July 2014
6. Granerød J, Davison K, Stuart J, Crowcroft N. 'Clusters of meningococcal disease in educational establishments in the UK April 2001 to March 2002' Communicable Disease and Public Health 2004: volume 7, issue 1, pages 51 to 55
7. Davison KL, Andrews N, White JM, Ramsay ME, Crowcroft NS, Rushdy AA and others. 'Clusters of meningococcal disease in school and preschool settings in England and Wales: what is the risk?' Archives of Disease in Childhood 2004: volume 89, issue 3, pages 256 to 260
8. Zangwill KM, Schuchat A, Riedo FX, Pinner RW, Koo DT, Reeves MW and others. 'School-based clusters of meningococcal disease in the United States: descriptive epidemiology and a case-control analysis' Journal of the American Medical Association 1997: volume 277, issue 5, pages 389 to 395
9. Whittaker R, Dias JG, Ramliden M, Kodmon C, Economopoulou A, Beer N and others. 'The epidemiology of invasive meningococcal disease in EU/EEA countries 2004 to 2014' Vaccine 2017: volume 35, issue 16, pages 2,034 to 2,041
10. Goldschneider I, Gotschlich EC, Artenstein MS. 'Human immunity to the meningococcus. II. Development of natural immunity The Journal of Experimental Medicine 1969: volume 129, issue 6, pages 1,327 to 1,348

11. Boutet R, Stuart JM, Kaczmarek EB, Gray SJ, Jones DM, Andrews N. 'Risk of laboratory-acquired meningococcal disease' *Journal of Hospital Infection* 2001: volume 49, issue 4, pages 282 to 284
12. Orr H, Kaczmarek E, Sarangi J, Pankhania B, Stuart J. 'Cluster of meningococcal disease in rugby match spectators' *Communicable Disease and Public Health* 2001: volume 4, issue 4, pages 316 to 318
13. Edwards EA, Devine LF, Sengbusch GH, Ward HW. 'Immunological investigations of meningococcal disease. III. Brevity of group C acquisition prior to disease occurrence' *Scandinavian Journal of Infectious Diseases* 1977: volume 9, issue 2, pages 105 to 110
14. Bygraves JA, Urwin R, Fox AJ, Gray SJ, Russell JE, Feavers IM and others. 'Population genetic and evolutionary approaches to analysis of *Neisseria meningitidis* isolates belonging to the ET-5 complex' *Journal of Bacteriology* 1999: volume 181, issue 18, pages 5,551 to 5,556
15. Bai X, Findlow J, Borrow R. 'Recombinant protein meningococcal serogroup B vaccine combined with outer membrane vesicles' *Expert Opinion on Biological Therapy* 2011: volume 11, issue 7, pages 969 to 985
16. Clark SA, Campbell H, Ribeiro S, Bertran M, Walsh L, Walker A, Willerton L, Lekshmi A, Bai X, Lucidarme J, Ladhani SN, Borrow R. 'Epidemiological and strain characteristics of invasive meningococcal disease prior to, during and after COVID-19 pandemic restrictions in England' *Journal of Infections* 2023: volume 87, issue 5, pages 385 to 391
17. Campbell H, Borrow R, Salisbury D, Miller E. 'Meningococcal C conjugate vaccine: the experience in England and Wales' *Vaccine* 2009: volume 27 supplement 2: B20 to B29
18. Findlow Helen, Campbell Helen, Lucidarme Jay, Andrews Nick, Linley Ezra, Ladhani Shamez, Borrow Ray. '[Serogroup C *Neisseria meningitidis* disease epidemiology, seroprevalence, vaccine effectiveness and waning immunity, England, 1998/1999 to 2015/2016](#)' *Euro Surveillance* 2019: volume 24, issue 1, pii=1700818
19. Ladhani SN, Beebeejaun K, Lucidarme J, Campbell H, Gray S, Kaczmarek E and others. 'Increase in endemic *Neisseria meningitidis* capsular group W sequence type 11 complex associated with severe invasive disease in England and Wales' *Clinical Infectious Diseases* 2015: volume 60, issue 4, pages 578 to 585
20. Lucidarme J, Hill DM, Bratcher HB, Gray SJ, du Plessis M, Tsang RS and others. 'Genomic resolution of an aggressive, widespread, diverse and expanding meningococcal serogroup B, C and W lineage' *Journal of Infection* 2015: volume 71, issue 5, pages 544 to 552.
21. Campbell H, Andrews N, Parikh SR, White J, Edelstein M, Bai X, Lucidarme J, Borrow R, Ramsay ME, Ladhani SN. 'Impact of an adolescent meningococcal ACWY immunisation programme to control a national outbreak of group W meningococcal disease in England: a national surveillance and modelling study' *Lancet Child Adolescent Health* 2022: volume 6, issue 2, pages 96 to 105
22. Parikh SR, Campbell H, Gray SJ, Beebeejaun K, Ribeiro S, Borrow R, Ramsay ME, Ladhani SN. '[Epidemiology, clinical presentation, risk factors, intensive care admission and outcomes of invasive meningococcal disease in England, 2010 to 2015](#)' *Vaccine* April 2018: pii: S0264-410X(18)30212-3

23. Parikh SR, Andrews NJ, Beebeejaun K, Campbell H, Ribeiro S, Ward C and others. 'Effectiveness and impact of a reduced infant schedule of 4CMenB vaccine against group B meningococcal disease in England: a national observational cohort study' *Lancet* 2016: volume 388, issue 10,061, pages 2,775 to 2,782
24. Ladhani SN, Andrews N, Parikh SR and others. 'Vaccination of infants with meningococcal group B vaccine (4CMenB) in England' *New England Journal of Medicine* 2020: volume 382, pages 309 to 317
25. Stephen A. Clark, Helen Campbell, Sonia Ribeiro, Marta Bertran, Lloyd Walsh, Andrew Walker, Laura Willerton, Aiswarya Lekshmi, Xilian Bai, Jay Lucidarme, Shamez N. Ladhani, Ray Borrow. 'Epidemiological and strain characteristics of invasive meningococcal disease prior to, during and after COVID-19 pandemic restrictions in England' *Journal of Infection* 2023: volume 87, issue 5, pages 385 to 3391
26. Trotter CL, Lingani C, Fernandez K, Cooper LV, Bitá A, Tevi-Benissan C, Ronveaux O, Préziosi MP, Stuart JM. 'Impact of MenAfriVac in 9 countries of the African meningitis belt, 2010 to 2015: an analysis of surveillance data' *Lancet Infectious Diseases* 2017: volume 17, pages 867 to 872
27. Miller E, Salisbury D, Ramsay M. 'Planning, registration, and implementation of an immunisation campaign against meningococcal serogroup C disease in the UK: a success story' *Vaccine* 2001: supplement 20, 1:S58 to S67
28. Trotter CL, Andrews NJ, Kaczmarski EB, Miller E, Ramsay ME. 'Effectiveness of meningococcal serogroup C conjugate vaccine 4 years after introduction' *Lancet* 2004: volume 364, issue 9,431, pages 365 to 367
29. Ladhani SN, Andrews N, Parikh SR and others. 'Vaccination of infants with meningococcal group B vaccine (4CMenB) in England' *New England Journal of Medicine* 2020: volume 382, pages 309 to 317
30. Ladhani SN, Campbell H, Andrews N, Parikh SR, White J, Edelstein M, Clark SA, Lucidarme J, Borrow R, Ramsay ME. 'First real-world evidence of Meningococcal group B vaccine, 4CMenB, protection against meningococcal group w disease: prospective enhanced national surveillance, England' *Clinical Infectious Diseases* 2021: volume 73, issue 7, pages e1,661 to e1,668
31. Ladhani SN, White PJ, Campbell H, Mandal S, Borrow R, Andrews N, Bhopal S, Saunders J, Mohammed H, Drisdale-Gordon L, Callan E, Sinka K, Folkard K, Fifer H, Ramsay ME. 'Use of a meningococcal group B vaccine (4CMenB) in populations at high risk of gonorrhoea in the UK' *Lancet Infectious Diseases* 2024: volume S1473-3099, issue 24, 00031-8
32. Helen S Marshall, Timo Vesikari, Peter C Richmond, Jacek Wysocki, Leszek Szenborn, Johannes Beeslaar, Jason D Maguire, Paul Balmer, Robert O'Neill, Annaliesa S Anderson, Jean-Louis Prégaldien, Roger Maansson, Han-Qing Jiang, John L Perez. 'Safety and immunogenicity of a primary series and booster dose of the meningococcal serogroup B-factor H binding protein vaccine (MenB-FHbp) in healthy children aged 1 to 9 years: 2 phase 2 randomised, controlled, observer-blinded studies' *The Lancet Infectious Diseases* 2023: volume 23, issue 1, pages 103 to 116
33. National Institute of Clinical Excellence. 'Overview: Meningitis (bacterial) and meningococcal disease: recognition, diagnosis and management'

34. Public Health England (PHE). 'Guidance on the prevention and management of meningococcal meningitis and septicaemia in higher education institutions: raising awareness, promoting immunisation and planning ahead' 2017
35. [SIGN sign grading system 1999 2012.pdf](#)
36. Ragunathan L, Ramsay M, Borrow R, Guiver M, Gray S, Kaczmarski EB. 'Clinical features, laboratory findings and management of meningococcal meningitis in England and Wales: report of a 1997 survey' *Journal of Infection* 2000: volume 40, 0163 to 4453, 74 to 79
37. Brooks A, Lucidarme J, Campbell H, Campbell L, Fifer H, Gray S, Hughes G, Lekshmi A, Schembri G, Rayment M, Ladhani SN, Ramsay ME, Borrow R. 'Detection of the United States *Neisseria meningitidis* urethritis clade in the UK, August and December 2019: emergence of multiple antibiotic resistance calls for vigilance' *Euro Surveillance* 2020 volume 25, issue 15, page 2000375
38. Rodrigues CMC, Jolley KA, Smith A, Cameron JC, Feavers IM, Maiden MCJ. 'Meningococcal deduced vaccine antigen reactivity (MenDeVAR) index: a rapid and accessible tool that exploits genomic data in public health and clinical microbiology applications' *Journal of Clinical Microbiology* 2020: volume 59, issue 1, pages e02161 to e02120
39. Rodrigues CMC, MacDonald L, Ure R, Smith A, Cameron JC, Maiden MCJ. '[Exploiting real-time genomic surveillance data to assess 4CMenB meningococcal vaccine performance in Scotland, 2015 to 2022](#)' *mBio* 2023: 14:e00499-23
40. Campbell H, Borrow R, Arumugam C, Ramsay M, Ladhani S. 'Invasive meningococcal disease as a cause of sudden and unexpected death in a teenager: the public health importance of confirming the diagnosis'
41. Hastings L, Stuart J, Andrews N, Begg N. 'A retrospective survey of clusters of meningococcal disease in England and Wales, 1993 to 1995: estimated risks of further cases in household and educational settings' *Communicable Disease Report review* 1997: volume 7, issue 13, R195 to R200
42. De Wals P, Hertoghe L, Borlee-Grimee I, De Maeyer-Cleempoel S, Reginster-Haneuse G, Dachy A and others. 'Meningococcal disease in Belgium: secondary attack rate among household, day-care nursery and pre-elementary school contacts' *Journal of Infection* 1981: volume 3, 1 Supplement, pages 53 to 61
43. Purcell B, Samuelsson S, Hahné SJM, Ehrhard I, Heuberger S, Camaroni I and others. 'Effectiveness of antibiotics in preventing meningococcal disease after a case: systematic review' *British Medical Journal* 2004: volume 328, issue 7,452, page 1,339
44. Cartwright KA, Stuart JM, Robinson PM. 'Meningococcal carriage in close contacts of cases' *Epidemiology and Infection* 1991: volume 106, issue 1, pages 133 to 141
45. Kristiansen B-E, Tveten Y, Jenkins A. 'Which contacts of patients with meningococcal disease carry the pathogenic strain of *Neisseria meningitidis*? A population-based study' *British Medical Journal* 1998: volume 317, issue 7,159, page 621
46. Leitmeyer K. 'European Risk Assessment Guidance for Infectious Diseases transmitted on Aircraft – the RAGIDA project' *Euro Surveillance* 2011: volume 16, issue 16, pii=19845
47. Hoek MR, Christensen H, Hellenbrand W, Stefanoff P, Howitz M, Stuart JM. 'Effectiveness of vaccinating household contacts in addition to chemoprophylaxis after a

- case of meningococcal disease: a systematic review' *Epidemiology and Infection* 2008: volume 136, issue 11, pages 1,441 to 1,447
48. Communicable diseases weekly report, week 48 2022, New South Wales Government
 49. Cohen MS, Steere AC, Baltimore R, von Graevenitz A, Pantelick E, Camp B and others. 'Possible nosocomial transmission of group Y *Neisseria meningitidis* among oncology patients' *Annals of Internal Medicine* 1979: volume 91, issue 1, pages 7 to 12
 50. Riewerts Eriksen NH, Espersen F, Laursen L, Skinhøj P, Høiby N, Lind I. 'Nosocomial outbreak of group C meningococcal disease' *British Medical Journal* 1989: volume 298, issue 6,673, pages 568 to 569
 51. Gold R, Goldschneider I, Lepow ML, Draper TF, Randolph M. 'Carriage of *Neisseria meningitidis* and *Neisseria lactamica* in infants and children' *Journal of Infectious Diseases* 1978: volume 137, issue 2, pages 112 to 121
 52. Orr HJ, Gray SJ, Macdonald M, Stuart JM. 'Saliva and meningococcal transmission' *Emerging Infectious Diseases* 2003: volume 9, issue 10, pages 1,314 to 1,315
 53. Telisinghe L. 'Systematic review of the effect of antibiotics and/or vaccination in preventing subsequent disease among household contacts of cases of meningococcal disease' World Health Organization May 2014 Report number 1.7
 54. Fraser A, Gafer-Gvili A, Paul M, Leibovici L. 'Antibiotics for preventing meningococcal infections' *The Cochrane database of systematic reviews* 2006, issue 4, Cd004785
 55. Schwartz B, Al-Tobaiqi A, Al-Ruwais A, Fontaine RE, Aashi J, Hightower AW and others. 'Comparative efficacy of ceftriaxone and rifampicin in eradicating pharyngeal carriage of group A *Neisseria meningitidis*' *Lancet* 1988: volume 1, issue 8,597, pages 1,239 to 1,242
 56. Podgore JK, Girgis N, L-Refai E, Abdel-Moneim A. 'A double-blind randomized trial of cefixime compared to rifampin in the eradication of meningococcal pharyngeal carriage in a closed population' *Journal of Tropical Medicine* 1993: volume 2, issue 5, pages 41 to 45
 57. Girgis N, Sultan Y, Frenck RW, Jr., El-Gendy A, Farid Z, Mateczun A. 'Azithromycin compared with rifampin for eradication of nasopharyngeal colonization by *Neisseria meningitidis*' *Pediatric Infectious Disease Journal* 1998: volume 17, issue 9, pages 816 to 819
 58. 'Public health management of sporadic cases of invasive meningococcal disease and their contacts' European Centre for Disease Prevention and Control (ECDC) 2010
 59. British National Formulary, NICE 2023
 60. Cuevas LE, Kazembe P, Mughogho GK, Tillotson GS, Hart CA. 'Eradication of nasopharyngeal carriage of *Neisseria meningitidis* in children and adults in rural Africa: a comparison of ciprofloxacin and rifampicin' *Journal of Infectious Diseases* 1995: volume 171, issue, 3, pages 728 to 731
 61. Drew TM, Altman R, Black K, Goldfield M. 'Minocycline for prophylaxis of infection with *Neisseria meningitidis*: high rate of side effects in recipients' *Journal of Infectious Diseases* 1976, volume 133, issue 2, pages 194 to 198
 62. Burkhardt JE, Walterspiel JN, Schaad UB. 'Quinolone arthropathy in animals versus children' *Clinical Infectious Diseases* 1997: volume 25, issue 5, pages 1,196 to 1,204
 63. Drossou-Agakidou V, Roilides E, Papakyriakidou-Koliouka P, Agakidis C, Nikolaides N, Sarafidis K and others. 'Use of ciprofloxacin in neonatal sepsis: lack of adverse effects up

- to one year' *Pediatric Infectious Disease Journal* 2004: volume 23, issue 4, pages 346 to 349
64. Yee CL, Duffy C, Gerbino PG, Stryker S, Noel GJ. 'Tendon or joint disorders in children after treatment with fluoroquinolones or azithromycin' *Pediatric Infectious Disease Journal* 2002: volume 21, issue 6, pages 525 to 529
 65. Clark J, Lakshman R, Galloway A, Cant A. 'Does cefotaxime eradicate nasopharyngeal carriage of *N. meningitidis*?' *Archives of Disease in Childhood* 2002: volume 87, issue 5, page 449
 66. Simmons G, Jones N, Calder L. 'Equivalence of ceftriaxone and rifampicin in eliminating nasopharyngeal carriage of serogroup B *Neisseria meningitidis*' *Journal of Antimicrobial Chemotherapy* 2000: volume 45, issue 6, pages 909 to 911
 67. Al-Abri SS, Abuhasan MY, Albayat SSA, Bai X, Bastaki H, Borrow R, Caugant DA, Dbaiibo G, Deghmane AE, Dinleyici EC, Ghuneim N, Sheek-Hussein M, Lucidarme J, Leng S, Koliou MG, Sáfadi MAP, Salman JA, Al-Sanouri T, Smith V, Taha MK, Vázquez J, Wright C, Yezli S. 'Meningococcal disease in the Middle East: a report from the Global Meningococcal Initiative' *Journal of Infections* 2024: volume 88, issue 2, pages 71 to 76
 68. MM, Robinson JL, Romeu B, Sadarangani M, Sáez-Llorens X, Sáfadi MAP, Stephens DS, Stuart JM, Taha MK, Tsang RSW, Vazquez J, De Wals P. 'Meningococcal disease in North America: updates from the Global Meningococcal Initiative' *Journal of Infections* 2022: volume 85, issue 6, pages 611 to 622
 69. Willerton L, Lucidarme J, Walker A, Lekshmi A, Clark SA, Walsh L, Bai X, Lee-Jones L, Borrow R. 'Antibiotic resistance among invasive *Neisseria meningitidis* isolates in England, Wales and Northern Ireland (2010/2011 to 2018/2019)' *PLoS One* 2021: volume 16, issue 11, article :e0260677
 70. Barroso D. '*Neisseria meningitidis* nasopharynx colonization of diseased patients on presentation and on discharge' *Tropical Doctor* 1999: volume 29, issue 2, pages 108 to 109
 71. Alvez F, Aguilera A, Garcia-Zabarte A, Castro-Gago M. 'Effect of chemoprophylaxis on the meningococcal carrier state after systemic infection' *Pediatric Infectious Disease Journal* 1991: volume 10, issue 9, page 700
 72. Abramson JS, Spika JS. 'Persistence of *Neisseria meningitidis* in the upper respiratory tract after intravenous antibiotic therapy for systemic meningococcal disease' *Journal of Infectious Diseases* 1985: volume 151, issue 2, pages 370 to 371
 73. Davis H, McGoodwin E, Reed TG. 'Anaphylactoid reactions reported after treatment with ciprofloxacin' *Annals of Internal Medicine* 1989: volume 111, issue 12, pages 1,041 to 1,043
 74. Burke P, Burne SR. 'Allergy associated with ciprofloxacin' *British Medical Journal* 2000: volume 320, issue 7,236, page 679
 75. Connell WR. 'Safety of drug therapy for inflammatory bowel disease in pregnant and nursing women' *Inflammatory Bowel Diseases* 1996: volume 2, issue 1, pages 33 to 47
 76. Ferrero S, Ragni N. 'Inflammatory bowel disease: management issues during pregnancy' *Archives of Gynecology and Obstetrics* 2004: volume 270, issue 2, pages 79 to 85

77. Cassina M, Fabris L, Okolicsanyi L, Gervasi MT, Memmo A, Tiboni GM and others. 'Therapy of inflammatory bowel diseases in pregnancy and lactation' *Expert Opinion on Drug Safety* 2009: volume 8, issue 6, pages 695 to 707
78. Lin KJ, Mitchell AA, Yau WP, Louik C, Hernandez-Diaz S. 'Safety of macrolides during pregnancy' *American Journal of Obstetrics and Gynecology* 2013: volume 208, issue 3, pages 221.e1 to e8
79. Dautzenberg B, Grosset J. 'Tuberculosis and pregnancy' *Revue des maladies respiratoires* 1988: volume 5, issue 3, page 279
80. Nahum GG, Uhl K, Kennedy DL. 'Antibiotic use in pregnancy and lactation: what is and is not known about teratogenic and toxic risks' *Obstetrics and Gynecology* 2006: volume 107, issue 5, pages 1,120 to 1,138
81. Ladhani SN, Cordery R, Mandal S, Christensen H, Campbell H, Borrow R and others. 'Preventing secondary cases of invasive meningococcal capsular group B (MenB) disease using a recently-licensed, multi-component, protein-based vaccine (Bexsero®)' *Journal of Infection* 2014: volume 69, issue 5, pages 470 to 480
82. Ladhani SN, Campbell H, Lucidarme J, Gray S, Parikh S, Willerton L and others. 'Invasive meningococcal disease in patients with complement deficiencies: a case series (2008 to 2017)' *BMC Infectious Diseases* 2019: volume 19, issue 1, page 522
83. Simmons RD, Kirwan P, Beebeejaun K, Riordan A, Borrow R, Ramsay ME and others. 'Risk of invasive meningococcal disease in children and adults with HIV in England: a population-based cohort study' *BMC Medicine* 2015: volume 13, issue 1, page 297
84. 'Meningococcal deduced vaccine antigen reactivity (MenDeVAR) index: a rapid and accessible tool that exploits genomic data in public health and clinical microbiology applications' *Journal of Clinical Microbiology*
85. 'Exploiting real-time genomic surveillance data to assess 4CMenB meningococcal vaccine performance in Scotland, 2015 to 2022' *mBio*
86. Ricco M, Vezzosi L, Odone A, Signorelli C. 'Invasive meningococcal disease on the workplaces: a systematic review' *Acta bio-medica: Atenei Parmensis* 2017: volume 88, issue 3, pages 337 to 351
87. Gilmore A, Stuart J, Andrews N. 'Risk of secondary meningococcal disease in health-care workers' *Lancet* 2000: volume 356, issue 9,242, pages 1,654 to 1,655
88. Coia JE, Ritchie L, Adisesh A, Makison Booth C, Bradley C, Bunyan D and others. 'Guidance on the use of respiratory and facial protection equipment' *Journal of Hospital Infection* 2013: volume 85, issue 3, pages 170 to 182
89. Bunyan D, Ritchie L, Jenkins D, Coia JE. 'Respiratory and facial protection: a critical review of recent literature' *Journal of Hospital Infection* 2013: volume 85, issue 3, pages 165 to 169
90. Loveday HP, Wilson JA, Pratt RJ, Golsorkhi M, Tingle A, Bak A and others. 'epic3: national evidence-based guidelines for preventing healthcare-associated infections in NHS hospitals in England' *Journal of Hospital Infection* 2014: volume 86, supplement 1:S1 to S70
91. Weber A, Willeke K, Marchioni R, Myojo T, McKay R, Donnelly J and others. 'Aerosol penetration and leakage characteristics of masks used in the health care industry' *American Journal of Infection Control* 1993: volume 21, issue 4, pages 167 to 173

92. Chen CC, Willeke K. 'Aerosol penetration through surgical masks' American Journal of Infection Control 1992: volume 20, issue 4, pages 177 to 184
93. CDC. 'Control and prevention of meningococcal disease and control and prevention of serogroup C meningococcal disease: evaluation and management of suspected outbreaks' Morbidity and Mortality Weekly Report 1997: volume 46, RR-5, pages 1 to 21
94. Stuart JM. 'Managing outbreaks the public health response' In: 'Meningococcal disease: methods and protocols' (editors: Pollard AJ, Maiden MCJ) Humana Press Inc 2001, pages 257 to 272
95. Kuhns DM, Nelson CT, Feldman HA, Kuhn LR. 'The prophylactic value of sulfadiazine in the control on meningococcic meningitis' Journal of the American Medical Association 1943: volume 123, issue 6, pages 335 to 339
96. Jackson LA, Alexander ER, DeBolt CA, Swenson PD, Boase J, McDowell MG and others. 'Evaluation of the use of mass chemoprophylaxis during a school outbreak of enzyme type 5 serogroup B meningococcal disease' Pediatric Infectious Disease Journal 1996: volume 15, issue 11, pages 992 to 998
97. Shehab S, Keller N, Barkay A, Leitner L, Leventhal A, Block C. 'Failure of mass antibiotic prophylaxis to control a prolonged outbreak of meningococcal disease in an Israeli village' European Journal of Clinical Microbiology and Infectious Diseases 1998: volume 17, issue 11, pages 749 to 753
98. Irwin DJ, Miller JM, Milner PC, Patterson P, Richards RG, Williams DA and others. 'Community immunization programme in response to an outbreak of invasive *Neisseria meningitidis* serogroup C infection in the Trent region of England 1995 to 1996' Journal of Public Health Medicine 1997: volume 19, issue 2, pages 162 to 170
99. Ardern K, Bowler S, Hussey RM, Regan CM. 'Managing meningococcal disease case clusters: art or science?' Journal of Epidemiology and Community Health 1999: volume 53, issue 9, pages 565 to 571
100. Olof S, Lorraine E, Berhane A I, Alexander P, Anders M, Sara TH, Martin S, Per O, Hans F, Bianca S, Eva S, Paula M, Susanne J. 'Prevalence and persistence of *Neisseria meningitidis* carriage in Swedish university students' Epidemiology and Infection 2023: volume 151, e25
101. Marshall HS, McMillan M, Koehler AP, Lawrence A, Sullivan TR, MacLennan JM, Maiden MCJ, Ladhani SN, Ramsay ME, Trotter C, Borrow R, Finn A, Kahler CM, Whelan J, Vadivelu K, Richmond. 'Meningococcal B vaccine and meningococcal carriage in adolescents in Australia' New England Journal of Medicine 2020: volume 382, issue 4, pages 318 to 327
102. Barker RM, Shakespeare RM, Mortimore AJ, Allen NA, Solomon CL, Stuart JM. 'Practical guidelines for responding to an outbreak of meningococcal disease among university students based on experience in Southampton' Communicable Disease and Public Health 1999: volume 2, issue 3, pages 168 to 173

About the UK Health Security Agency

UKHSA is responsible for protecting every member of every community from the impact of infectious diseases, chemical, biological, radiological and nuclear incidents and other health threats. We provide intellectual, scientific and operational leadership at national and local level, as well as on the global stage, to make the nation's health secure.

[UKHSA](#) is an executive agency, sponsored by the [Department of Health and Social Care](#).

Prepared by: Helen Campbell, Mary Ramsay and Shamez Ladhani of the UKHSA Immunisation Division together with Ray Borrow of the UKHSA Meningococcal Reference Unit, NHS National Services Scotland, Public Health Wales and Public Health Agency in Northern Ireland. We are grateful for the additional contributions made by the UKHSA Vaccine Preventable Invasive Bacterial Infections Forum and UKHSA Health Protection Teams in particular Surrey and Sussex Health Protection Unit and North East Health Protection Team.

For queries relating to this document, please contact:

Shamez Ladhani
Immunisation and Vaccine-Preventable Diseases Division
UKHSA Colindale
61 Colindale Avenue
Colindale
London NW9 5EQ
Email: shamez.ladhani@ukhsa.gov.uk

© Crown copyright 2025

Published: December 2025

Publishing reference: GOV-19795



You may re-use this information (excluding logos) free of charge in any format or medium, under the terms of the Open Government Licence v3.0. To view this licence, visit [OGL](#). Where we have identified any third party copyright information you will need to obtain permission from the copyright holders concerned.



UKHSA supports the
Sustainable Development Goals

