



Department  
for Environment  
Food & Rural Affairs

# Risk assessment on the likelihood of spread of highly pathogenic avian influenza H5Nx associated with bird fairs, shows, markets, sales and other gatherings

Qualitative risk assessment

February 2026

We are responsible for improving and protecting the environment. We aim to grow a green economy and sustain thriving rural communities. We also support our world-leading food, farming and fishing industries.

Defra is a ministerial department, supported by 34 agencies and public bodies.



© Crown copyright 2026

This information is licensed under the Open Government Licence v3.0. To view this licence, visit [www.nationalarchives.gov.uk/doc/open-government-licence/](http://www.nationalarchives.gov.uk/doc/open-government-licence/)

Where we have identified any third party copyright information you will need to obtain permission from the copyright holders concerned.

This publication is available at [www.gov.uk/defra](http://www.gov.uk/defra)

Any enquiries regarding this publication should be sent to us at [iadm@apha.gov.uk](mailto:iadm@apha.gov.uk)

## Contents

Qualitative risk assessment .....	1
Executive summary .....	5
Background.....	7
Trends and risk levels in the current epizootic .....	7
Captive bird orders glossary .....	8
Hazard identification .....	9
Risk question .....	9
Scope .....	9
Terminology related to the assessed level of risk.....	9
Entry assessment .....	9
Likelihood of captive bird being infected prior to being taken to gathering .....	10
Psittaciformes and Passeriformes .....	11
Columbiformes .....	11
Birds of prey (Accipitriformes and Falconiformes) .....	12
Ratites .....	13
Galliformes .....	13
Anseriformes .....	13
Likelihood of HPAIV H5Nx not being detected prior to gathering .....	18
Psittaciformes .....	18
Columbiformes .....	18
Birds of prey .....	18
Passerines .....	19
Galliformes .....	20
Anseriformes .....	20

Ratites .....	20
Exposure assessment .....	20
Spread of disease within the gathering .....	21
Qualitative risk assessment .....	21
Consequence assessment.....	23
Spread of disease from the gathering .....	23
Mitigation measures.....	24
Conclusions .....	25
Uncertainties.....	26
References .....	26
Annex 1 .....	27

## Executive summary

The hazard is high pathogenicity avian influenza (HPAI) virus H5Nx. Here, a rapid risk assessment (RRA) is undertaken to reassess the risk of HPAI H5Nx introduction from poultry and captive birds taken to bird fairs, shows, markets, sales and other gatherings (henceforth referred to as “gatherings”). The risk of other birds subsequently becoming infected at a gathering should an infected bird be introduced into that gathering and then spreading through dispersing to different establishments has been assessed in previous versions of this RRA as medium and this risk assessment uses the same methodology as previous iterations. Critical to this RRA therefore is the likelihood that captive birds are infected prior to their transport to a gathering, together with the likelihood that any infected birds are not detected (and hence not removed) before reaching the gathering.

The main route of exposure of captive birds prior to transport is through contact with wild birds (either direct or indirect contact). At the present time (26 February 2026) the wild bird risk for HPAI H5Nx is at **very high** in Great Britain. Since the beginning of the 2025 to 2026 season (1 October 2025) wild bird cases have been reported across all of England, most of Wales and parts of Scotland with many cases inland. Resident waterbird species, namely mute swans, Canada and greylag geese are still badly affected. Since the increase in IPs reported in early November 2025 across England and Wales, the number of IPs each week has fallen through January and into February such that the risk level for poultry with suboptimal biosecurity was reduced from very high to **high** with medium uncertainty.

In terms of the bird species considered in this RRA, those poultry and bird species that are most likely to be taken to gatherings include 7 bird orders, namely Psittaciformes (parrots), Columbiformes (doves and fancy pigeons including racing pigeons if taken to a gathering), birds of prey (Accipitriformes and Falconiformes), Passeriformes (finches), Galliformes (poultry and game birds), Anseriformes (ducks and geese) and Ratites (ostriches, emus and rheas). The baseline risk of those birds being infected prior to the gathering for this RRA is based on that of poultry in Great Britain taking into account whether the biosecurity is more in line with sub-optimal biosecurity (Galliformes, Anseriformes and Ratites) or stringent biosecurity (Psittaciformes, Columbiformes, birds of prey, and Passeriformes). As with the previous gatherings RRA, that risk for each of the 7 bird orders is then refined based on specific differences in access to the environment where residual infectivity from wild birds may be present. Captive ducks and geese (Anseriformes) kept outside or in un-netted ponds are likely to have contact with wild ducks, geese and swan species particularly if kept at premises near to where wild waterfowl gather. The risk of infection of captive Anseriformes prior to attending the gathering or show is therefore **very high** (with medium uncertainty) due to the ongoing detections in wild waterfowl which could visit water bodies utilised by or frequented by captive Anseriformes. There is also a high probability of not detecting infected Anseriformes prior to their being taken to the gathering because of reduced likelihood of clinical signs. The risk of infection for captive Galliformes across Great Britain prior to attending the gathering is assessed to be **high** (with medium

uncertainty) in line with the risk for poultry with suboptimal biosecurity. The risk of infection prior to the gathering for Psittaciformes, Columbiformes, birds of prey and Passeriformes is assumed to be the same as that for poultry with stringent biosecurity, namely **low** (having been lowered from medium in the last week). However, birds of prey (if used for hunting) and ratites (being kept outside and not housed) are allocated a **high** risk of infection prior to the gathering.

The results of the risk assessment are set out in the table below.

**Summary table: Qualitative risk assessment for entry and subsequent spread of HPAI H5Nx at bird gatherings according to bird group (for February 2026)**

Captive bird group taken to gathering	Risk of entry and subsequent spread	Uncertainty
Psittaciformes	Low	Medium
Columbiformes	Low	Medium
Birds of Prey	Low	Medium
Passerines	Low	Medium
Galliformes	Medium	Medium
Anseriformes	High	Medium
Ratites	Low	Medium

For most captive bird groups, the risk of entry and subsequent spread of HPAI H5 at a gathering is **low**. Since our previous risk assessment in November 2025, the risk of entry and subsequent spread at bird gatherings has decreased from medium to low for Columbiformes, and Passerines. This reflects the recent lowering of the risk level for poultry in Great Britain with stringent biosecurity. However, the risk from captive Anseriformes is still assessed to be **high** due to the likely exposure to wild water birds at water bodies where the captive birds may be kept. This in effect reflects the direct exposure of captive Anseriformes across Great Britain to potentially infected wild waterbirds (both resident species and migratory species) for which the risk level is still very high. The risk of entry and subsequent spread of HPAI H5 at a gathering through Galliformes is still assessed to be **medium** (despite the high risk level for poultry with suboptimal biosecurity) because most infected Galliformes would be detected prior to being taken to the gathering. Similarly the risk of entry and spread from Psittaciformes, ratites and birds of prey is **low** because infected birds would be detected prior to being taken to the gathering.

The uncertainty in the risk levels is medium mainly reflecting the uncertainty for the likelihood of infected birds not being detected in each group. It should be noted that **any suspicion of infection being reported by the owner prior to the gathering is a critical protective barrier for gatherings**.

Given this range of risks, there are options around the licensing of gatherings which include:

- to ban for the period of the AI prevention zone
- to allow but only for certain species of birds
- to allow but with improved requirements for biosecurity or
- captive birds could be housed for 7 days prior to moving to the gathering

If the available biosecurity measures are considered to be insufficient to mitigate the risks identified, or if there is insufficient assurance that they will be complied with, then a ban on some or all categories of gathering would be justified.

## Background

Here, a rapid risk assessment (RRA) is undertaken to reassess the risk of high pathogenicity avian influenza (HPAI) H5N1 introduction and spread from poultry and captive birds taken to bird fairs, shows, markets, sales and other gatherings (henceforth referred to as “gatherings”). Gatherings of birds involve the coming together and subsequent dissemination of live kept birds (as well as people, vehicles and equipment) and for this reason can facilitate the introduction and spread of avian notifiable disease including HPAI to different locations across Great Britain. The magnitude of this risk is influenced by the number of different groups of birds brought together and the likelihood of their being already infected at their point of origin. Movements out of an SZ or PZ around a confirmed infected premises are not permitted but other than this, traceability of poultry is lightly regulated so most moves outside a restriction zone are of uncertain origin and status.

Previous Defra risk assessments (Defra, 2016) were used as a basis for the general licence allowing bird gatherings to take place while minimising the risk of introduction of avian notifiable diseases to these events and mitigating the likelihood and impact of any subsequent spread. Here the risk assessment is updated to accommodate the risk levels for HPAI H5N1 in wild birds and poultry in February 2026.

## Trends and risk levels in the current epizootic

Please see our updated outbreak assessment for details: [Avian influenza \(bird flu\) in Europe, Russia and the UK - GOV.UK](#). It is interesting to note (Table 1) from the point of view of gatherings that resident Anseriformes (mallard ducks, Canada and greylag geese) together with the resident (mainly mute) swans account for almost two thirds of the wild bird cases in terms of wild bird group with migratory whooper swan cases and migratory geese together accounting for almost 15%. Resident swans (mainly mute swans but some black swans) represent the most frequently detected species (Table 1). Also of note is the detection of 7 cases in pigeons in Great Britain suggesting that the very high wild bird infection pressure has even impacted on less susceptible species present in high abundance.

**Table 1: Numbers (and percentages) of wild bird cases of HPAI H5N1 and H5Nx according to bird group in Great Britain collected from 1 October 2025 to 17 February 2026.**

Wild Bird Group	Number of Positive Cases	Percentage
Resident swans	233	39.1%
Resident goose or duck	144	24.2%
Raptor	61	10.2%
Migrant goose or duck	55	9.2%
Gull	32	5.4%
Migrant swan	30	5.0%
Gamebird	21	3.5%
Pigeon	7	1.2%
Owl	6	1.0%
Seabird	3	0.5%
Wader	3	0.5%
Heron	1	0.2%
Total	596	100%

## Captive bird orders glossary

The bird orders of captive birds considered are set out in Table 2.

**Table 2:2 Glossary of captive bird orders considered here with examples.**

Order	Examples
Psittaciformes	Parrots
Columbiformes	Pigeons and doves
Birds of prey (Accipitriformes and Falconiformes)	Hawks and falcons
Passeriformes	Perching birds (finches and canaries)
Galliformes	Turkeys, pheasants, chickens, guineafowl.
Anseriformes	Ducks and geese
Ratites	Ostriches, emus and rheas

# Hazard identification

The hazard identified is highly pathogenic avian influenza virus, (HPAI) H5Nx.

## Risk question

- 1) *What is the risk of the introduction of highly pathogenic avian influenza H5Nx onto bird fairs, shows, markets, sales and other gatherings?*
- 2) *What, if any, management options are available to reduce the likelihood and the impact of introduction and subsequent spread of avian notifiable disease through the abovementioned gatherings?*

## Scope

This qualitative risk assessment covers the risk of introduction and subsequent spread of avian notifiable disease to and from bird gatherings organised in Great Britain that were legally moved to the event from within the UK.

This risk assessment does not assess the risk related to illegal movements, failure to report clinical disease, false certification, breaches in biosecurity etc. Any risks potentially presented by (or to) wild birds are also not assessed here.

## Terminology related to the assessed level of risk

For the purpose of the risk assessment, the following terminology will apply (OIE, 2004):

- **Negligible:** So rare that it does not merit to be considered
- **Very low:** Very rare but cannot be excluded
- **Low:** Rare but does occur
- **Medium:** Occurs regularly
- **High:** Occurs very often
- **Very high:** Event occurs almost certainly

It should be noted, however, that the risk terminologies here do not represent how often an event will occur but more indicate the probability of the event occurring.

## Entry assessment

The Defra (2016) assessments concluded that the overall risk of the introduction of avian notifiable diseases including HPAI to a bird gathering is related to the ongoing disease situation in Great Britain, both in domestic poultry and in wild birds.

As adopted in previous RRAs for gatherings, the risk of those birds which may attend gatherings being infected with HPAIV H5Nx is based on the official risk levels for poultry in Great Britain being infected through background environmental contamination from infected wild birds (see Table 3). **This risk is currently (26 February 2026) high (with medium uncertainty)** for poultry where biosecurity is sub-optimal and **low (with medium uncertainty)** for poultry where biosecurity is stringent. The Avian Influenza Prevention Zone (AIPZ) is still in place (26 February 2026). A mandatory housing order was put in place across the whole of England on 6 November 2025, Wales on the 13 November 2025 and Northern Ireland on 6 November 2025 ([National Housing Order declared to protect poultry from Avian Influenza - GOV.UK](#)). An AIPZ is in place in Scotland but there is no mandatory housing.

It is assumed that keepers of any birds participating at a gathering are doing so in full compliance with the legal requirements for movements of live birds, and that birds are not coming from areas under disease control restrictions.

## Likelihood of captive bird being infected prior to being taken to gathering

Captive birds, such as those held in collections, zoos or approved bodies are already semi-housed and should be kept separate from wild waterfowl. For some, it will be difficult to prevent access to their water environment (ducks, geese, swans, penguins, pelicans, flamingos etc) and it is unlikely that it will be possible to house indoors, so every effort should be made to prevent wild waterfowl access.

Due to the varying level of biosecurity (and considered lower than stringent), those birds being taken to gatherings and shows will be of more uncertain infection status than commercial poultry with stringent biosecurity. Also monitoring and data for commercial production mean that commercial birds can be tracked more easily than for those birds at gatherings and shows. Birds from non-commercial settings carry greater uncertainty. The baseline risk of those birds being infected prior to the gathering for this RRA is based on that of poultry in Great Britain taking into account whether the biosecurity is more in line with sub-optimal biosecurity (Galliformes, Anseriformes and Ratites) or stringent biosecurity (Psittaciformes, Columbiformes, birds of prey, Passeriformes).

Currently (26 February 2026) risk to poultry with sub-optimal biosecurity is **high with medium uncertainty** and the risk to poultry with stringent biosecurity is **low with medium uncertainty**. The risks for the 7 bird orders considered here (Table 2) are refined in Table 3 based on specific differences in access to the environment where residual infectivity from wild birds may be present. It has already been noted above that the risk levels exist as bands, though those at the higher end are much narrower than the lower risk levels.

## Psittaciformes and Passeriformes

Passerine species infected with HPAI H5Nx globally include several species of finch, sparrow, thrush and myna ([Bird & mammal species affected by H5Nx HPAI](#)). In March 2022 there were 6 cases of HPAI H5/H5N1 in pied wagtails in Crewe. In the 2022 to 2023 season in Great Britain there were 4 cases of HPAI H5 in passerines, namely a reed warbler and 3 carrion crows. However, since the beginning of the 2023 to 2024 season, there have been no detections of HPAI H5Nx in found dead wild passerines in Great Britain. This may partly reflect the character of this citizen-science based surveillance scheme, where dead small passerines are relatively hard to find and may not remain available for collection and analysis. In Europe, according to WOAHA, passerine cases that have been reported since 1 October 2025 to 17 February 2026 include one thrush in the Netherlands, 4 blackbirds (Belgium, France and Poland), one redwing in Belgium, a Sylvia warbler in Austria and single unidentified passerines in Germany and Poland. In addition, some 19 cases have been reported in corvid species in Europe since 1 October 2025 (to 17 February 2026). Psittaciformes and collections of ornamental passerines are kept in aviaries and some may even be kept indoors where there is less likelihood of direct wild bird contact. The risk of infection of Psittaciformes and kept passerines prior to being taken to the gathering is considered to be the same as that for the risk to poultry with stringent biosecurity, namely **low**. It is noted that those Psittaciformes and passerines kept in outside aviaries will be at higher risk and at the higher end of the low risk level range. The uncertainty is medium due to the variation in how these birds are kept, i.e ranging from a single bird kept indoors to outdoor aviaries with large collections. Further species susceptibility across these species is poorly defined.

## Columbiformes

Racing pigeons competing within Great Britain or Europe are not considered here (they are the subject of a separate and specific risk assessment). However, racing pigeons taken to other gatherings are included here together with related Columbiformes of other species or breeds. There have been relatively few reports of cases in wild Columbiformes on WOAHA over the last few years of the epizootic.

In a study of domestic pigeons, inoculated oculo-nasally with HPAI H5N8 (Clade 2.3.4.4 sub-group icA3) of Korean origin, Kwon et al, (2017) concluded that, though pigeons have lower susceptibility than some other species, they can be infected with HPAI H5N8 when exposed to high doses and could excrete the virus in sufficiently high doses to infect other species of birds. Pigeons could also be fomite transmitters of the virus. A study in which 18 pigeons were inoculated intranasally with HPAI H5N8 (clade 2.3.4.4 subgroup B) from South Africa reported viral shedding in medium and high-dose pigeons for up to 8 days (Abolnik et al, 2018). Infected pigeons successfully transmitted virus to contact pigeons and seroconversion was observed in 2 of the high-dose group chickens (Abolnik et al, 2018).

More recent experimental research has provided further evidence for the low level of susceptibility of these birds to HPAI H5N1 (Di Genova et al. 2025). Thus, following direct inoculation, pigeons did not develop clinical signs, and only those inoculated with the highest dose shed viral RNA or seroconverted to H5N1, revealing a 50% minimum infectious dose of  $10^5$  50% egg infectious dose (EID<sub>50</sub>). Even in the high-dose group, only low-level shedding and environmental contamination were observed, and low-level viral RNAs were present in the tissues of directly inoculated pigeons, with no distinct pathological lesions. A single pigeon in the medium-dose group ( $10^4$  EID<sub>50</sub>) exhibited a single subthreshold level of viral RNA in the oropharyngeal cavity. In contrast, viral RNA was shed by 7 of the 8 pigeons in the high dose group ( $10^6$  EID<sub>50</sub>) between 1 and 7 days post infection. In the case of HPAI H5N1, pigeons did not transmit the virus to other pigeons or chickens placed in direct contact (Di Genova et al. 2025).

Columbiformes may be kept in aviaries outside where direct contact with waterbirds might occur. Since October 2025 there have been 6 cases of HPAI H5N1 in collared doves, 5 cases in unidentified Columbiformes and 18 cases of HPAI H5/H5N1 in woodpigeons in Europe according to IZS<sub>Ve</sub> (2026). Similarly on WOA<sub>H</sub> from 1 October 2025 to 14 February 2026 there have been 32 cases of HPAI H5/H5N1 in pigeons, including 15 woodpigeons, 4 feral pigeons and 7 Eurasian collared doves in Europe. In Great Britain from 1 October 2025 to 9 February 2026 there have been 4 wood pigeon cases, 3 rock doves and 1 unidentified dove. These cases may reflect the very high wild bird infection pressure in Great Britain and Europe in the 2025 to 2026 season together with the abundance of pigeons as wild birds. It should be noted that just because in an experimental setting the “infectious dose” for pigeons is high (see Di Genova et al. 2025), does not mean that lower doses do not cause infection in at least one pigeon when large enough numbers of pigeons are each exposed to low doses. Comorbidities and health status also likely impact upon infection outcome. While infected pigeons may only shed at low levels, low levels may be sufficient to infect more susceptible species at a gathering including Galliformes or Anseriformes.

However, given the recent (February 2026) reduction in the risk level for infection of poultry with stringent biosecurity to low together with the relatively low susceptibility of Columbiformes to HPAI H5Nx and the relatively few cases in wild Columbiformes (discussed above), the likelihood of Columbiformes being infected prior to being taken to the gathering is assumed to be **low** with medium uncertainty.

### **Birds of prey (Accipitriformes and Falconiformes)**

There are many reports of wild raptors being infected both in Great Britain and globally with HPAI H5N1 and also with HPAI H5N5, perhaps because they are exposed to very high viral doses when scavenging infected birds. Most captive birds of prey will be fed on commercial feed considered to be low risk for infection (such as day-old chicks, reared small rodents) but some are fed shot game, including wild duck, which increases risk of infection through feeding. Given the very high risk level in wild birds, including cases in ducks which captive falcons may be used to hunt, the overall risk of captive birds of prey

having disease prior to being taken to a gathering is still assumed to be **high** (with medium uncertainty). The risk would be lower for those falcons not contacting wild birds including waterfowl hence the medium uncertainty.

## Ratites

Given the small number of ratites (compared to other poultry species) in Great Britain, it was previously considered here that the likelihood of an infected bird being taken to a gathering was **low**. Ratites are always kept outside and it may not be possible to house them given their large size. In 2025 there were 3 reported cases of HPAI H5N1 in ratites including a captive emu in Brazil, a captive ostrich in Mexico and most recently in a captive rhea species in Portugal according to WOA. HPAI H5N1 has also been detected in a captive Greater Rhea in Great Britain (Coombes et al 2025). With the current very high wild bird infection pressure, the likelihood that a ratite is infected prior to a gathering is maintained at **high** (with high uncertainty).

## Galliformes

The likelihood of captive Galliformes being infected prior to being taken to the gathering is assumed to be the same as that currently (26 February 2026) for poultry with suboptimal biosecurity namely **high** with medium uncertainty. The medium uncertainty reflects the fact that the captive Galliformes' risk level could be the same as that currently for wild birds, namely very high, if they are allowed to mix with wild birds on ranges.

## Anseriformes

Since October 2025 the number of cases of HPAI H5N1 in wild resident Anseriforme species (including mute swans, greylag geese, Canada geese and mallard ducks) has remained high across Great Britain through the winter and into February 2026 with smaller numbers of cases in migratory waterbirds (including pink-footed geese, barnacle geese and whooper swans). Captive ducks and geese kept outside or in un-netted ponds are likely to have contact with wild duck, geese and swan species at premises near where wild waterfowl gather. The risk level for captive Anseriformes (ducks and geese) prior to being taken to gatherings is therefore assessed to be **very high** with medium uncertainty across Great Britain (Table 3) reflecting the current (26 February 2026) very high risk level for wild birds and the medium uncertainty for poultry with suboptimal biosecurity.

It is important to note that the wild bird surveillance figures for Great Britain are based on passive surveillance of found dead birds reported to Defra by the general public and as such, may be affected by several factors including frequency of visitors accessing areas with bird populations, the potential for immunity in the wild bird population (which may result in fewer birds developing clinical disease and or dying with HPAI), variable surveillance system sensitivity, as well as the size, location and accessibility of carcasses. Thus, this wild bird surveillance does not necessarily capture all of the cases that occur,

although the current number of cases reported fully supports the very high-risk level for wild birds.

**Table 3: Likelihood of HPAI H5Nx infection in each group of captive birds prior to attending the gathering**

Order	Examples	Likelihood of being infected currently (26 February 2026) prior to going to gathering	Uncertainty
Psittaciformes	Parrots	Low - reflecting risk for poultry with stringent biosecurity is now at low. Mostly kept in aviaries or indoors. Risk may be higher if keep in outdoor aviaries.	Medium
Columbiformes	Pigeons and doves	Low – based on the low risk level for infection of poultry with stringent biosecurity together with the relatively low susceptibility of Columbiformes to HPAI H5Nx and the relatively few cases in wild Columbiformes (discussed in text).	Medium
Birds of prey	Hawks and falcons	High – particularly if allowed to catch wild waterfowl at the current time (26 February 2026) when wild bird risk is very high.	Medium

Order	Examples	Likelihood of being infected currently (26 February 2026) prior to going to gathering	Uncertainty
Passerines	Finches and canaries	Low - kept in aviaries or indoors. Risk may be higher if keep in outdoor aviaries, although risk for poultry with stringent biosecurity is now at low.	Medium
Galliformes	Turkeys, pheasants, chickens, guineafowl	<sup>a</sup> High- based on current risk to poultry with poor biosecurity. Captive Galliformes likely to be outdoors with greater opportunity of exposure to wild birds hence suboptimal biosecurity.	<sup>b</sup> Medium

Order	Examples	Likelihood of being infected currently (26 February 2026) prior to going to gathering	Uncertainty
Anseriformes	Ducks and geese	Very high – same as that for wild birds assuming that captive ducks and geese have greater contact than other captive bird groups with wild birds including resident ducks, geese, swans and gulls. Captive ducks and geese likely to be outdoors with access to ponds where HPAI H5N1 could be introduced if present in local wild resident ducks, geese and swans particularly at current time of year when wild waterfowl aggregate.	<sup>b</sup> Medium
Ratitias	Ostriches, emus and rheas.	High – kept outdoors with access to wild birds	High

<sup>a</sup>Based on the current risk level (26 February 2026) to poultry with suboptimal biosecurity.

<sup>b</sup>In the case of some captive Galliformes and many Anseriformes the risk level will be the same as that currently for wild birds, namely very high, if they are allowed to mix with wild birds on ranges and ponds respectively, hence medium uncertainty.

## Likelihood of HPAIV H5Nx not being detected prior to gathering

The level of awareness of avian notifiable diseases in Great Britain is thought to be generally high and suspicions of clinical disease in poultry and other captive birds would be reported reasonably quickly, generally within a few days, particularly with the current awareness in the sector of HPAI H5N1 in wild birds. Movement restrictions for disease control purposes would be uniformly implemented based on domestic and retained EU Community legislation. The length of the virus incubation period as well as the possibility of virus shedding during this time is an important factor to be considered while assessing these risks. However, no official incubation period for avian influenzas is established for bird species other than poultry and the actual length of the incubation period is affected by numerous factors including the disease, the virus load, the actual virus strain, the species, and immune status.

### Psittaciformes

Cases of HPAI H5N1 infection in wild parrots have been reported on WOAHP mainly in South America in recent seasons. A range of species have been reported, including blue-and-yellow macaw, budgerigar, burrowing parrot, Mealy parrot, red-and-green macaw, scarlet macaw, slender-billed parakeet, white-winged parakeet, and yellow-headed Amazon parrot. There has been a case where budgerigars were infected at a premises in Norfolk in mid-December 2024. The likelihood of HPAI infection being undetected for Psittaciformes is unknown, but given the multiple detections in wild parrots in South America and the cases in zoos, including the captive bird establishment in Great Britain, is assumed to be **low** with medium uncertainty.

### Columbiformes

In the study of Abolnik et al (2018) with HPAI H5N8 there were no clinical signs observed in any of the pigeons involved despite viral shedding in medium and high-dose pigeons for up to 8 days. The study of Kwon et al. (2017) with domestic pigeons inoculated oculonasally with HPAI H5N8 (Clade 2.3.4.4 sub-group icA3) of Korean origin, showed no clinical signs or mortality even though, relatively high levels of shedding were observed in half of the pigeons. Pigeons do not show clinical signs when infected with H5N8 (Abolnik et al, 2018, Kwon et al, 2017). Similarly, 6 of 6 pigeons given high doses of HPAI H5N1 survived infection to 14 days post infection despite shedding virus (Di Genova et al. 2025). Therefore it is assumed here that there is a **high** likelihood (medium uncertainty) of not detecting Columbiformes infected with HPAI H5.

### Birds of prey

The likelihood of disease not being detected prior to the gathering is **low** for birds of prey. This is because birds of prey seem particularly susceptible to morbidity and mortality from

HPAI H5Nx with many affected in the wild in both Great Britain and globally. It is known that birds of prey generally develop overt clinical signs if infected with the H5N8 HPAI virus, and HPAI H5N1 positive found-dead birds of prey have been a feature in Great Britain in recent weeks and previous seasons (along with captive birds of prey in previous seasons). The infected birds of prey would show clinical signs within 2-3 days of feeding and are likely to be detected prior to taking to a gathering, hence the low risk. However, this may not be the case of all birds of prey, for example white-tailed eagle and the uncertainty is medium.

## Passerines

Passerines, including canaries and finches, are known to be susceptible to low pathogenicity avian influenza (LPAI) H5 and that they can shed large amounts of viral RNA through the respiratory route (Marché et al 2018). While they do not show clinical signs or mortality with LPAI, if infected with HPAI H5 then a proportion would be expected to show mortality and there have been reports of mortality of wild passerines both globally and in Great Britain from HPAI H5N1 strains. The likelihood of infected passerines not being detected is therefore assumed to be **medium** with medium uncertainty.

Ellis et al. (2021) have studied the susceptibility, infection and transmission of LPAI in starlings. All starlings were infected if inoculated experimentally and shed viral RNA for around 7 days. Furthermore, starlings were infected by drinking water used by an experimentally infected mallard duck. However, 4 naive starlings placed in a cage with 5 starlings infected by inoculation were not infected (in 4 replicate experiments) suggesting starling to starling transmission is not efficient and that starlings do not act as maintenance hosts. Those starlings infected by the mallard duck water did shed viral RNA at high levels and it must be assumed could therefore infect poultry and Anseriformes. Thus, starlings could act as a bridging species to poultry.

The key points from the paper by Ellis et al. (2021) for assessing the risks from passerines at gatherings are that captive passerines would be infected if exposed to infected wild ducks for example if kept outside in an aviary where flooding could cause ingress of water containing virus shed by wild birds. Furthermore, those starlings infected through exposure to water from infected ducks shed sufficient quantities of viral RNA to infect poultry and Anseriformes at a gathering (although the authors have not demonstrated this and suggest this as further work). The fact that starling to starling spread did not occur efficiently means that only a small proportion of passerines in an aviary would be infected if one were to be infected, so reducing the chances of the owner realising HPAI was present in the flock. So if passerine to passerine transmission were efficient the whole aviary would go down and the owner would notice. This further supports our conclusion here that the probability of not detecting infected passerines prior to going to a gathering is **medium**.

It's worth noting that there is considerable uncertainty here. We have assessed the uncertainty as medium in the RRA for all 7 bird groups. However, within the passerine family there is a range of species so there could also be considerable variation in the risk

depending on the passerine species. In addition, there will be variation between holdings, husbandry and biosecurity that we cannot assess.

It is considered here that infection of captive birds is more likely to be detected in parrots and birds of prey than in passerines. This is because there are often more passerines in a cage than parrots and birds of prey, so if one or 2 passerines in a large group die they may not be noticed so much as if a single parrot or bird of prey on its own in a cage dies.

## Galliformes

Galliformes show high mortality in the poultry outbreaks. Similarly, pheasants are susceptible to H5 HPAI infection and rapidly show clinical signs although those birds infected recently would still be in the incubation period. Therefore, the likelihood of infected pheasants or chickens not being detected prior to taking to a gathering would be **low**. However, partridges may not show clinical signs and could be missed ([Seekings et al. 2024](#)). Furthermore, although Galliformes show high mortality it is considered that detection or reporting could be low in backyard poultry. Therefore, for the purpose of this risk assessment it is assumed the likelihood of disease not being detected or reported in Galliformes on backyard premises which could be taken to gatherings is **medium** with medium uncertainty.

## Anseriformes

While there have been many dead Anseriformes-positive wild bird cases reported both in Great Britain and globally, many wild ducks and geese have survived, and given the level of exposure and virus circulation in recent years there is a possibility of immunity though this is unknown. Furthermore, some ducks may not show clinical signs suggesting that the likelihood of infected ducks and geese not being detected prior to the gathering event is **high**. The uncertainty is medium.

## Ratites

Elsayed et al. (2022) reported mortality rates of 90% in ostrich flocks in South Africa infected with HPAI H5N8. The birds suffered loss of appetite, dropped production, and oculo-nasal discharges with bleeding from natural orifices. The probability of not being detected is therefore **low**, with medium uncertainty.

# Exposure assessment

This section deals with the risk that an infected bird entering a gathering will spread disease to other birds at that gathering. If infection were to be introduced to a bird gathering, the likelihood of its spread depends on a number of factors such as the pathogenicity and transmissibility of the virus, the amount of virus being shed by the infected bird, the nature and layout of the gathering such as the housing and proximity of

the participating birds, whether the birds are mixing, access to common water sources and whether they are in direct contact with visitors.

## Spread of disease within the gathering

Spread through a gathering is based on the aggregated risk from multiple contacts between the infected bird introduced to the gathering and the other birds from different origins and hence going to different destinations after mixing at the gathering. From an epidemiological point of view, the probability of one or more birds actually being exposed to infection at the gathering is defined as  $1-(1-p)^n$ , where  $p$  is the probability that an infected bird introduced to the gathering infects another bird given a contact, and  $n$  is the number of such contacts. There are no data on  $p$  and  $n$  at gatherings. However, even if  $p$  were low then just ten contacts with the infected bird at the gathering would be sufficient to give an aggregated probability of medium that at least one exposed bird at the gathering would be infected. If  $p$  were medium, then just one contact would be sufficient to give a medium aggregated probability that at least one bird would be exposed. Therefore, the risk of disease spread, if introduced to a gathering would be at least medium. This is increased to high in the case of Anseriformes and Galliformes because more individual birds of these 2 groups may be taken to gatherings.

As discussed above there is a low likelihood of spread between pigeons and between passerines. However, this may reflect their lower susceptibilities and they could transmit to more susceptible species such as Galliformes at the gathering through fomite routes for example. Also the low risks from a single passerine or pigeon escalates with increasing number of birds, justifying the medium aggregated risk assigned to these 2 groups.

## Qualitative risk assessment

For the purpose of this RRA a simplified risk pathway is used comprising three steps each described by a qualitative risk as set out below:..

1. Probability captive bird is infected prior to gathering;
2. Probability infected bird is not detected prior to entering gathering; and
3. Probability transmission occurs at gathering given infected bird introduced.

The 3 qualitative risks in the pathway for each bird order are combined using the matrix of Gale et al. (2009) to give the overall risk of spread of HPAIV H5Nx from bird gatherings in Great Britain currently.

Incursions at gatherings from wild birds are not considered here.

The risk assessment for the current situation (26 February 2026) is set out in Table 4.

**Table 4: Qualitative risk assessment for entry and subsequent spread of HPAI H5Nx at bird gathering according to bird group based on risk levels for 26 February 2026. Uncertainty in parentheses.**

	Psittaciformes	Columbiformes	Birds of Prey	Passerines	Galliformes	Anseriformes	Ratites
Risk of infection prior to gathering (20 February 2026) see Table 3	Low (Medium)	Low (Medium)	High (Medium)	Low (Medium)	High (Medium)	Very high (Medium)	High (High)
Likelihood of infected bird not being detected prior to gathering	Low (Medium)	High (Medium)	Low (Medium)	Medium (Medium)	Medium (Medium)	High (Medium)	Low (Medium)
Risk of spread of disease at gathering	Medium	Medium	Medium	Medium	High	High	Medium
<sup>a</sup> Overall risk	Low ( <sup>c</sup> Medium)	Low ( <sup>c</sup> Medium)	Low ( <sup>b</sup> Medium)	Low ( <sup>c</sup> Medium)	Medium ( <sup>b</sup> Medium)	High ( <sup>b</sup> Medium)	Low ( <sup>b</sup> Medium)

<sup>a</sup>Overall risk of spread of HPAIV H5Nx at bird gathering calculated as lowest probability in the column according to matrix of Gale et al. (2009).

<sup>b</sup>The medium uncertainty reflects the medium uncertainty in the probability of detection of an infected bird prior to the gathering.

<sup>c</sup>The medium uncertainty reflects the medium uncertainty in the probability the captive bird is infected prior to the gathering.

# Consequence assessment

## Spread of disease from the gathering

If undisclosed avian notifiable disease were to be introduced to a bird gathering by live birds, there is a possibility that unless disease is confirmed during the event, it would remain undisclosed until after the gathering – and therefore there is a potential for further spread. The likelihood of this depends on the length of the event as with longer events the possibility that birds show signs of disease becomes greater, although it may also increase the number of potential contacts between birds at the gathering.

The extent of onward geographical spread depends on the extent of contact and spread between birds at the gathering itself and also where the birds are transported to following the event. The most effective way of preventing such spread would be to detect suspicion of disease at the time of the gathering, while the birds are still together. However, if the source of infection is a (group of) birds with subclinical infection, this increases the risk of onward spread. The size of the gathering, levels of biosecurity and length of the gathering would directly affect the number of potential contacts between infected and susceptible birds.

The consequence of avian influenza being detected in birds either at or having attended a gathering during the risk period is a serious matter for not only industry but also for the competent authorities. This could lead to a multi-focal outbreak in birds which have moved to different parts of the country, which are difficult to trace.

Any outbreak of notifiable avian disease has a significant impact on the UK poultry industry, through the trade and economic impacts on the producer. This is the same for any notifiable avian influenza virus. Average costs to government may be between £2 million and £4 million per outbreak, depending on the number of birds involved and complexity of the investigation.

If disease is detected at a gathering before it concludes and before the birds are dispersed, Government would face a complex challenge relating to disease control at the gathering, including dealing with a large number of owners who may be resistant to the need to cull their birds.

Whilst spread from a gathering may not lead to widespread disease into the commercial sector and may be restricted to small producers, the case in 2007 in the UK involving a market showed that there is a potential scenario for this occurrence. While for the majority of shows and gatherings involve birds classified as not destined for the food chain (as breeders or producers) it is important to note that even one outbreak in backyard premises would still lead to implementing disease control measures, as specified in the current regulations.

The risk assessment presented here addresses the risk of transmission at gatherings and does not consider how many other establishments could be infected, which would depend on the size of the gathering and individual bird keepers who attend them. In the 2020 to 2021 epizootic of HPAIV H5Nx in Europe, there was evidence that many captive bird and poultry outbreaks could be traced to a single dealer in southern Germany emphasizing the potential consequence that gatherings could have.

A reasonable worst-case outcome for multiple outbreaks to occur would be for an infected but apparently healthy bird to be taken to a gathering where it infects some, but not all, of the other birds present, but disease is not detected. The birds at the gathering are then taken to widely distributed premises. One or more of the infected birds is then detected through passive surveillance leading to at least one outbreak being confirmed with consequent disease control zones, impacts on industry and a costly tracing exercise. There has been one comparable case in recent years but this involved LPAI so the consequences were limited and again, for certain species of birds this is less likely.

Pigeons infected experimentally with HPAI H5N1 did not transmit the virus to  $n = 8$  pigeons or  $n = 8$  chickens placed in direct contact (Di Genova et al. 2025) although pigeons with HPAI H5N8 did infect contact pigeons (Abolnik et al, 2018). Although Columbiformes and Passerines may have lower susceptibilities and do not transmit HPAI readily to each other (as shown for starlings with LPAI (Ellis et al 2021), they could transmit to more susceptible species such as Galliformes at the gathering through fomite routes for example. Also the low risks from a single passerine or pigeon escalates with increasing number of birds so justifying the medium aggregated risk assigned to birds from these 2 groups spreading HPAI within a gathering.

## Mitigation measures

Measures to mitigate the risk of disease entering a gathering and the potential impact include disease vigilance and prompt reporting of any suspicion, high levels of biosecurity and accurate record keeping to assist in any possible tracing exercise following the event. A table has been provided in previous versions of the document (ANNEX1). The risk of further (cross) contamination and onward spread occurring at and beyond the gathering could be mitigated by maintaining high levels of biosecurity, including reducing the number of potential contacts between infected and susceptible birds and informing livestock keepers about the need for vigilance for clinical signs of avian notifiable disease. However, there is a significant infection pressure currently, with HPAI widely circulating in wild bird populations and poultry premises with varying levels of biosecurity implemented. A quarantine or standstill period on holdings prior to attendance at gatherings and also after return of birds from gatherings could also be considered, although may be impractical, particularly for backyard premises and against the backdrop of the current infection pressure and anticipated trajectory which is expected to continue until after aggregations disperse and environmental conditions are less suitable for survivability.

# Conclusions

Currently (26 February 2026) the risk from gatherings is predicted to be **low** for Psittaciformes, Columbiformes, Passerines, captive birds of prey and Ratites and **medium** for Galliformes. The risk is still **high** for captive Anseriformes (ducks and geese).

The risk for captive Anseriformes is **high** because kept ducks and geese, if they have access to the outdoors, would have greater contact with water bodies which could attract infected resident waterbirds such as Canada geese, greylag geese, mute swans and mallards. The **medium** risk from captive Galliformes is dependent on most infected Galliformes showing clinical signs and being detected prior to being taken to the gathering. The same applies to birds of prey and ratites.

The uncertainty in the risk levels is medium mainly reflecting the uncertainty for the likelihood of infected birds not being detected in each group. It should be noted that the **detection of infected birds by the owner prior to the gathering is an important protective barrier for gatherings**. This reflects the very high wild bird infection pressure currently across Great Britain.

Therefore in response to the risk questions:

- 1) What is the risk of the introduction of HPAI H5Nx into bird fairs, shows, markets, sales and other gatherings?

**The risk currently (at 26 February 2026) is low for Psittaciformes, kept birds of prey, Ratites, Columbiformes and passerines, but medium for Galliformes and high for Anseriformes (ducks and geese).**

- 2) What, if any, management options are available to reduce the likelihood and the impact of introduction and subsequent spread of avian notifiable disease through the above mentioned gatherings?

**Options are to ban, allow only certain species, where the risk is considered to be lower, or allow everything with stricter controls and this is in order of increasing risk. Given the disperse locations of cases in found dead wild birds and confirmed IPs, regionalisation is not recommended to be considered. When effective biosecurity measures are in place, housing is a barrier between poultry and pathogen. However, there is a continued infection pressure currently and this may be difficult to achieve. Additionally, the passive nature of the wild bird surveillance cannot inform of where background risk may be highest or lowest or indeed may be changing.**

# Uncertainties

For all 7 groups of captive birds the overall uncertainty in the predicted risk of entry and spread at a gathering is medium. Since our previous risk assessment in November 2025 the risks of entry and spread for Psittaciformes, Columbiformes and passerines have fallen from medium to low reflecting the decrease in the risk level for poultry with stringent biosecurity to low albeit with medium uncertainty. The medium uncertainty in the low risk for Psittaciformes, Columbiformes and passerines reflects the medium uncertainty in the probability the captive bird is infected prior to the gathering. The medium uncertainty for birds of prey, Anseriformes, Galliformes and Ratites reflects the medium uncertainty in the probability that a keeper actually detects infected birds and does not take them to the gathering.

# References

- Abolnik, C., Stutchbury, S. and Hartman, M. J. (2018) Experimental infection of racing pigeons (*Columba livia domestica*) with highly pathogenic Clade 2.3.4.4 sub-group B H5N8 avian influenza virus. *Veterinary Microbiology* 227:127-132.
- Coombes, H.A., et al (2025) [Infection of ratites with clade 2.3.4.4b HPAIV H5N1: Potential implications for zoonotic risk | bioRxiv](#)
- Ellis JW, Root JJ, McCurdy LM, Bentler KT, Barrett NL, VanDalen KK, et al. (2021) Avian influenza A virus susceptibility, infection, transmission, and antibody kinetics in European starlings. *PLoS Pathog* 17(8): e1009879. <https://doi.org/10.1371/journal.ppat.1009879>. [Avian influenza A virus susceptibility, infection, transmission, and antibody kinetics in European starlings | PLOS Pathogens](#)
- Defra (2016) [Risk assessment on the likelihood of spread of avian notifiable disease associated with bird fairs, shows, markets, sales and other gatherings \(publishing.service.gov.uk\)](#).
- Di Genova C, Warren CJ, Johnson S, Riccio S, Roper K, Thomas SS, Schlachter AL, Jorge D, Ralh K, Hassan J, Billington E, Nunez A, Brown IH, Slomka MJ, Banyard AC, James J. Pigeons exhibit low susceptibility and poor transmission capacity for H5N1 clade 2.3.4.4b high pathogenicity avian influenza virus. *J Gen Virol.* 2025 Sep;106(9):002156. doi: 10.1099/jgv.0.002156.

Coombes, H.A. et al. (2025) Infection of ratites with clade 2.3.4.4b HPAIV H5N1: Potential implications for zoonotic risk. bioRxiv 2025.09.08.674895;  
doi: <https://doi.org/10.1101/2025.09.08.674895>.

Gale, P., Brouwer, A., Ramnial, V., Kelly, L., Kosmider, R., Fooks, A.R. and Snary, E.L. (2009) Assessing the impact of climate change on vector-borne viruses in the EU through the elicitation of expert opinion. *Epidemiology and Infection* 138, 214-225.

Kwon, J.-H., Noh, Y.K., Lee, D.-H., Yuk, S.-S., Erdene-Ochir, T.-O., Noh, J.-Y., Hong, W.-T., Jeong, J.-H., Jeong, S., Gwon, G.-B., Song, C.-S., Nahm, S.-S. (2017). Experimental infection with highly pathogenic H5N8 avian influenza viruses in the Mandarin duck (*Aix galericulata*) and domestic pigeon (*Columba livia domestica*). *Veterinary Microbiology* 203: 95-102.

Seekings AH, Liang Y, Warren CJ, Hjulsager CK, Thomas SS, Lean FZX, Nunez A, Skinner P, Selden D, Falchieri M, Simmons H, Brown IH, Larsen LE, Banyard AC, Slomka MJ. Transmission dynamics and pathogenesis differ between pheasants and partridges infected with clade 2.3.4.4b H5N8 and H5N1 high-pathogenicity avian influenza viruses. *J Gen Virol*. 2024 Jan;105(1). doi: 10.1099/jgv.0.001946 ([jgv001946.pdf](#)).

## Annex 1

**Table 3. Comparison of measures available to regulate poultry gatherings and those applied in the current General Licence**

Measure	In force
General licence permitting gatherings	Yes
Specific licences	No
Withdraw licence (ie ban)	No
Notify APHA at least 7 days in advance including: a) date the gathering will take place b) location of the gathering c) details of the licensee (including full name, contact address and telephone number) d) anticipated numbers and type(s) of birds	Yes
No sales allowed unless additional conditions are met (see below)	No

<p>The licensee must make a record of all people who bring poultry or other captive birds to a gathering or take such birds from a gathering and keep the record for at least 3 months following the end of the gathering. The record must include at least the following information:</p> <ul style="list-style-type: none"> <li>a) full name</li> <li>b) home address</li> <li>c) telephone number</li> <li>d) number and type(s) of birds exhibited, raced, bought or sold</li> </ul>	Yes
<p>The licensee must not allow any cage, crate, basket or other container onto a gathering which is contaminated with bird droppings, bedding or other material of bird origin other than that from, or provided for, the birds brought to the gathering and must make this requirement known to all those bringing birds to the gathering in advertising, on entry forms or by any other means.</p>	Yes
<p>The licensee must ensure that any cages, crates, baskets or other containers not removed from the premises by those attending the gathering are cleansed and disinfected as soon as reasonably practicable after the end of the gathering and in any case before they are used again.</p>	Yes
<p>i) ensure that any feed to which poultry or other captive birds had access, and all bedding, droppings, other material of bird origin and other contaminants derived from birds at the gathering and which are left on the premises when the gathering has ended are -</p> <ul style="list-style-type: none"> <li>a) destroyed</li> <li>b) treated so as to remove the risk of transmission of disease</li> <li>c) disposed of so that birds do not have access to them; or</li> <li>d) disposed of as Category 2 products under the Animal By-Products Regulations 2005</li> </ul> <p>ii) where practicable, cleanse and disinfect those parts of the premises contaminated by such materials</p>	Yes
<p>14 days' notice to APHA</p>	No
<p>A named veterinary surgeon must be available on site during the whole time of the event for advice in case of suspect disease or a welfare problem. The veterinary surgeon should be responsible to ensuring that only clinically healthy birds in clean cages are entering the event.</p>	No
<p>Biosecurity advice must be distributed at the event.</p>	No
<p>Written contingencies, held by the nominated responsible person, must be available in the event of a disease incident at the event or nearby the event.</p>	No
<p>All cages used in the show must be cleansed and disinfected prior to and after the show.</p>	Implicitly yes
<ul style="list-style-type: none"> <li>• For sales: A record of all sales taking place at the event should be kept for at least 3 months, this should include: the name, address and telephone number of both the vendor and buyer and any identifying features or individual identification of the purchased bird(s).</li> </ul>	No

<ul style="list-style-type: none"> <li>• For sales: Buyers must isolate the purchased bird(s) from any other birds (except those purchased at the same event) for at least 1 week.</li> <li>• Any signs of ill health observed in the purchased bird(s) during this period must be reported to a veterinary surgeon and such birds must not be mixed with any other birds until the presence of an avian notifiable disease has been ruled out</li> </ul>	No
<ul style="list-style-type: none"> <li>• Restrict to birds of certain species – columbiformes, passerines and psittaciformes</li> </ul>	No
<ul style="list-style-type: none"> <li>• Restrict to regions only</li> </ul>	No