



Steeping River

Natural capital register and account

2025



Contents

Executive Summary	1
1. Introduction	4
1.1 Aim and purpose of this document	4
1.2 Usage statement	4
1.3 Catchment context	4
2. Understanding natural capital	7
2.1 An introduction to natural capital	7
2.2 The importance of considering natural capital within flood risk and water management infrastructure investment	9
2.3 Key accounting methods	10
3. Our approach to understanding the benefit and value of natural capital	12
3.1 Developing our evidence base	12
4. The natural capital register and account for the catchment	18
4.1 Materiality Assessment	18
4.2 Natural Assets	19
4.3 Ecosystem Services	33
4.4 Benefits	38
4.5 Value	44
5. Summary and next steps	50
5.1 Summary of the NCRAT results for the catchment	50
5.2 Strengths, weaknesses, opportunities and threats	52
6. Bibliography	A-1
Tables	
Table 1: Confidence rating definitions	14
Table 2: Materiality Assessment summary	18
Table 3: Drivers of environmental change considered in the Risk Register	31
Table 4: The definition of the ecosystem service significance ratings used in the NCRAT.	34
Table 5: The natural capital values calculated using the NCRAT colour coded according to the confidence interval.	48
Appendix A	A-7
Output - Benefits Statement	A-7

Executive Summary

Natural capital refers to the elements of the natural environment that provide valuable goods and services to people [1]. These include food, water, flood alleviation, thriving wildlife and places to enjoy. Natural capital underpins our wellbeing and economic prosperity, providing multiple benefits to society.

This Natural Capital Baseline Report provides an assessment of the natural capital assets within the Steeping catchment¹. The report highlights the importance of natural capital in supporting economic activity and human wellbeing through the provision of ecosystem services. The assessment was conducted using the Environment Agency's Natural Capital Register and Account Tool (NCRAT) and includes both quantitative and qualitative analysis undertaken through desk-based research, spatial analysis using Geographical Information Systems (GIS) and engagement with key stakeholders.

The catchment comprises 63.37km² (6,337 hectares (ha)) of terrestrial and coastal habitats, including urban land; agricultural land; coastal margins; freshwater habitats; mountains, moorlands and heaths; and woodlands. The catchment also contains local, national and international designated sites such as Gibraltar Point, The Wash, Saltfleet by Theddlethorpe Dune & Gibraltar Point, and Bratoft Meadows.

The key ecosystem services from natural assets in the catchment can be categorised into provisioning, regulating, cultural and supporting / bundled services, the estimated natural capital values of these are summarised below:

- Agriculture, water supply, timber, and renewable energy generation are key **provisioning services**. Agriculture alone contributes an estimated £5.9 million annually, while water supply is valued at nearly £4.2 million per year.
- Climate regulation, air quality improvement, hazard regulation, and pollination are critical **regulating services**. Coastal habitats play a significant role in carbon sequestration and climate regulation, sequestering an estimated 1940tCO₂e/yr² in the catchment. Conversely, the peatland in the catchment is likely to be in a degraded condition and consequently emits approximately 282tCO₂e/yr.
- Recreation, physical health, and aesthetic value are important **cultural services**. Despite being predominantly agricultural in land-use, it is estimated that the catchment attracts approximately 172,000 visitors annually to open spaces³, with an estimated value of £694,000 This contributes to natural capital value through improvements to physical and mental health (welfare) and can boost the local economy.
- **Supporting / bundled ecosystem services** in the catchment include water quality and biodiversity, which allow other ecosystem services to function and often involve multiple services or non-use values. The catchment hosts a large number of internationally

¹ The Steeping catchment is not strictly a hydrological catchment as there are movements of water both into and out of the area defined, that are not being considered in this report. However, the term 'catchment' has been used throughout the dialogue with the Environment Agency during establishment of the study areas and these areas are being defined for water management purposes. Therefore, the term catchment is used to describe the study area.

² Tonnes of carbon dioxide equivalent per year. This is a measure of how much carbon dioxide would have to be released to have an equivalent warming impact to the amounts of other greenhouse gases.

³ Open spaces considered within ORVal include publicly accessible green spaces within the catchment, for example, public parks, doorstep greens and nature reserves.

designated sites that support biodiversity conservation, for example, approximately 4% (248ha) of the catchment is designated as a Special Protection Area, Special Area of Conservation or Ramsar Site. Other statutory sites for nature conservation in the catchment include Sites of Special Scientific Interest and National Nature Reserves.

Ecosystem services that have not been valued but are important for the catchment include pollination and seed dispersal; control of pests and diseases; regulation of soil quality and erosion; biodiversity; and cultural services through the contribution of ecosystems to aesthetics, the historic environment, and a sense of place.

Other ecosystem services that have not been valued include volunteering and education. These ecosystem services have not been included due to a lack of data or because they are too complex to accurately determine with the information available. Since there is no marine habitat within the catchment, the provision of food from fish and shellfish landings has not been included as it is not appropriate to attribute these benefits to the study area. It should also be noted that the estimated value for agricultural activities does not include the negative impacts it can have on the other ecosystem service (for example water quality, carbon sequestration, hazard regulation and biodiversity) through pollution, biodiversity loss, erosion and peat degradation. Figure 1 provides an overview of the natural capital value within the catchment.

The natural capital assets in the catchment are at risk, which in turn creates risks for the delivery of ecosystem services. Broadly, the risks identified include climate change; intensive agricultural practices; urban development; poor water resource management; pollution; and invasive species. If not managed, these risks will degrade the quantity and quality of natural capital assets in the catchment and prevent the provision of ecosystem services that our societies economy and wellbeing rely on.

Conversely, there are many opportunities to enhance natural capital assets in the catchment, thereby increasing the provision of ecosystem services, and boosting local and global natural capital value. These opportunities include restoration of natural habitats including peatland, woodland and wetland; habitat creation and growth; implementing nature-based solutions (NbS); adopting sustainable practices in agriculture and water management; regenerative farming; and strengthening policy and governance that enables the conservation of natural capital assets and supports sustainable development. These interventions would deliver multiple benefits across a range of ecosystem service types.

The report highlights the critical role of natural capital in supporting the local community's resilience and wellbeing in the catchment, and the value of considering natural capital in strategic decision-making. Through addressing the identified risks and leveraging the opportunities, the Fens2100+ programme, and future investment projects should look to sustainably manage existing natural capital assets and look for opportunities to enhance ecosystem service delivery for future generations.




Figure 1: Summary of the NCRAT results for the catchment, showing the flow of ecosystem services and their benefits from natural assets

1. Introduction

1.1 Aim and purpose of this document

This report presents a catchment-scale natural capital baseline for the Fens2100+ Steeping River catchment (hereafter referred to as the 'catchment'). It provides an insight into the provision of ecosystem services by natural capital assets across the catchment, highlighting the wider benefits this has on society, such as the importance of natural capital in supporting economic activity and human wellbeing, as well as the risks and opportunities. It aims to provide decision makers with evidence to support future choices, investments and actions for the Fens2100+ programme.

The assessment was conducted using the Environment Agency's Natural Capital Register and Account Tool (NCRAT) suite of tools⁴ and includes an estimated total quantifiable natural capital value (£m) in the catchment, supported by qualitative evidence and analysis. More information on the importance of using a natural capital approach is provided in Section 2.2.

1.2 Usage statement

This report should be read alongside the Steeping River baseline evidence report and the Steeping River environmental and agricultural baseline appendix. This document provides a proportionate level of detail required for a strategic catchment-scale baseline. A detailed baseline analysis may be required at project level to align with the Environment Agency's Minimum Technical Requirements or Environmental Impact Assessment.

1.3 Catchment context

The Steeping River catchment covers 6337ha of east Lincolnshire, located at the far north-east of the Fens2100+ study area. It spans from the base of the chalk hills of the Lincolnshire Wolds, to the Lincolnshire Marshes, where the river outfalls at Burgh Sluice into The Wash near Gibraltar Point. The study area is shown in Figure 2.

The catchment is predominately artificially drained through a complex network of low-lying drains which are pumped into the Steeping River.

The catchment lies slightly above sea level (approximately 3 – 9m Above Ordnance Datum (AOD)) with a wide coastal plain partially consisting of drained salt marsh and priority habitats including coastal and floodplain grazing marsh and coastal saltmarsh. It is a highly productive agricultural area with approximately one-third of the land classified as Grade 1 agricultural land.

The northern area of the catchment is located within National Character Area (NCA) 42 Lincolnshire Coast and Marshes, a low-lying, broad plain bounded largely by the North Sea to the east, the Lincolnshire Wolds to the west, the Humber Estuary to the north and the entrance to The Wash to the south. Marine processes strongly influence the physical and biological character through sediment transfer. There are long, wide coastal views across this NCA, and also out to adjoining areas such as the North Norfolk coast. The land use within the catchment is primarily agricultural with areas to the north of the catchment

⁴ Environment Agency NCRAT includes: Environment Agency Natural Capital Register and Account Tool version 1.2, Workbook; and Environment Agency natural capital scorecard version 1.2

utilised for grazing and keeping of livestock. Field patterns decrease in size in the north of the catchment, where more intact hedgerows and blocks of woodlands are present.

The southern area of the catchment is located within NCA 46 The Fens. The Fens is a distinctive, historic and human influenced wetland landscape lying to the west of The Wash, the largest estuarine system in the UK. The NCA is notable for its large-scale, flat, open topography with extensive vistas to level horizons. Land use is predominately arable agriculture. Presence of irrigation reservoirs (over 1ha) in this area supports higher value cropping. Between The Wash and Skegness lies Lincolnshire Wildlife Trust Gibraltar Point National Nature Reserve (NNR). This site is recognised nationally and internationally for its species and habitats.

Key settlements include the historic port market town of Wainfleet All Saints, located near the western boundary of the catchment, and the village of Wainfleet St Mary, which is directly south of Wainfleet All Saints, separated by the Steeping River. There are several other linear settlements spread along minor roads and lanes in the north of the catchment area. Other key features include a number of small-scale renewable energy generation sites, caravan parks and camping sites, some of which also host fishing ponds/lakes.

The Local Planning Authority for the catchment is East Lindsey District Council, with the Environment Agency and the Lindsey Marsh Internal Drainage Board as Risk Management Authorities. The Environment Agency have permissive powers to carry out flood and coastal risk management activities on Main Rivers. In addition, the Environment Agency are responsible for producing a national strategy for Flood and Coastal Erosion Risk Management (FCERM). The latest strategy published in 2020 contains 'Measure 1.5.4'. This requires development of a long-term plan for managing future flood risk in the Fens.

Lindsey Marsh IDB play a vital role in managing water levels and drainage across the catchment for agricultural and environmental purposes. They are responsible for managing the complex network of drains that feed into the Main Rivers by pumps. They also regulate water levels on non-Main Rivers (ordinary watercourses). They seek to maintain and enhance biodiversity through implementation of their Biodiversity Action Plan (2010).

All organisations have a complicated task of managing flood risk whilst balancing agricultural water demands e.g. for irrigation and maintaining water supply rights to nature reserves.




Figure 2: Catchment boundary

2. Understanding natural capital

2.1 An introduction to natural capital

Natural capital refers to the renewable and non-renewable natural resources that combine to provide valuable goods and services to people [2]. All aspects of the natural environment - including soils; geological formations; habitats; and ecosystems - are natural capital assets which provide ecosystem services, from which flow benefits to society.

There is a growing awareness of the role of natural capital in delivering quality of place, and in enhancing health and wellbeing. Some benefits of natural capital can be measured and valued, but many are difficult to define and quantify. Taking a natural capital approach facilitates comparability between environmental and economic costs and benefits and can help organisation, planners and policy makers assess needs, identify dependencies on natural assets, and translate this into relevant information for decision making, which can in turn facilitate better design and achievement of policy, strategies, programmes and projects.

A natural capital framework

A natural capital framework, such as that shown in Figure 3, provides a coherent, comprehensive and consistent mechanism to analyse the key ecosystem services and benefits provided by nature. It is recognised by the HM Government's Green Book as a way to improve appraisal of a range of environmental benefits and effects [3].




Figure 3: A Natural Capital Framework, adapted from Enabling a Natural Capital Approach Guidance [4]

Natural capital / ecosystem assets

Natural capital assets are aspects of the natural environment. This includes but is not limited to:

- Habitats that serve as natural environments for living organisms such as woodland, heathland, marine and intertidal habitat, scrub, agricultural land, hedgerows, semi-natural grassland and urban trees;
- All living organisms including animals, plants, fungi, and bacteria;
- Green infrastructure, which comprises multifunctional green spaces such as parks and green walls;
- Geological features and processes including rocks, soils and minerals; and
- The water environment, including rivers, lakes, aquifers, and oceans.

The quality, quantity, location and connectivity of these natural capital assets impacts the ecosystems services and benefits they provide. These in turn can be affected by

background pressures, management practices and drivers of demand, therefore impacting their natural capital value.

Ecosystem Services

The benefit and contribution of nature to our society is a product of the various ecosystem services it provides, which can be grouped into four categories [5]:

- **Provisioning:** Material outputs from nature (e.g., seafood, water, fibre, genetic material);
- **Regulating:** Indirect benefits from nature generated through regulation of ecosystem processes (e.g., mitigation of climate change through carbon sequestration, water filtration by wetlands, erosion control and protection from storm surges by vegetation, crop pollination by insects);
- **Cultural:** Non-material benefits from nature (e.g., spiritual, aesthetic, recreational, and others) - wellbeing gains from sites and areas are often derived from a combination of the heritage, biodiversity and geodiversity aspects; and
- **Supporting/bundled services [5]:** Fundamental ecological processes that support the delivery of other ecosystem services (e.g., nutrient cycling, primary production, soil formation). Bundled services may include: Amenity, Biodiversity, Soil Quality, Water Quality, Landscape and non-use benefits.

Benefits

Ecosystem services provide a wealth of benefits to society and the economy, for example:

- Improved physical health through the supply of clean air, water and food; as well as regulation of temperature and flood protection.
- Improved mental health through provision of green and blue space for recreation;
- Improved economy through the availability and quality of commercial activities such as arable farming, aquaculture, timber production, and pastoral farming.

Each ecosystem service will have numerous benefits to society and the economy. For example, erosion control benefits human physical health by reducing the risk of dangerous landslides; mental health by improving access to nature; and benefits the economy by improving soil health for agriculture and reducing the cost of repairing infrastructure after a landslide. Sometimes, other economic inputs are required to enable ecosystem services to provide a benefit to society. For example, the health and wellbeing benefits of recreation requires providing accessibility, including functional gateways and path maintenance.

Value

The value of the benefits provided from ecosystem services in an area can be estimated using quantitative and qualitative information about the quantity, quality and location of natural capital assets, as well as information on how local natural assets are used by society – both locally and globally, and for private and public sectors.

- Quantitative valuation is best used to provide information on the marginal value of incremental changes in impacts or dependencies, either at a point in time or over a given period and is usually monetised in units (£). It can also be used to estimate the economic value of a range of ecosystem services that affect economic performance and human wellbeing - some of which are commonly unpriced and not considered in economic metrics such as Gross Domestic Product (GDP).
- Qualitative valuation, describing the natural capital impacts, is therefore also important as it can be used to take into account the importance, worth, or usefulness of the

impact and/or dependency by taking into account the context, to give equal values to any monetary figures presented.

Where products are directly or indirectly linked to ecosystem services and are traded in markets, valuation can be carried out using market prices - this is often the case for provisioning services such as timber production [4]. For other services that are not typically traded in markets, such as improved water quality or climate regulation, non-market based approaches are used. These typically involve measuring the price individuals are willing to pay to secure positive impacts or avoid negative impacts, through stated preference, revealed preference or cost-based approaches [4].

It should be noted that the representation of natural capital in monetary accounts will only ever be partial as a number of ecosystem services remain difficult to quantify and value. Some are highly location specific, for example water flow and impact on flood risk, and some are intangible or difficult to even place a monetary value on, for example cultural services, such as aesthetic experiences and sense of place.

Through valuing natural capital benefits, their importance to society and the economy can be better represented and accounted for in decision making about future place-based planning and investment.

2.2 The importance of considering natural capital within flood risk and water management infrastructure investment

The concepts of natural capital and ecosystem services are widely supported; the challenge, however, is in implementing the approach and embedding it in working practices, so that the value from the natural environment becomes an integral component of investment decision making. This is recognised in the State of Natural Capital Report for England 2024 [6], which explores the relationship between natural capital and infrastructure. Two key outcomes of the report that highlight the importance of natural capital include:

- Infrastructure, such as flood risk and water management infrastructure can deliver benefits for natural capital, such as through the provision of protected natural habitats and connecting corridors for species along linear infrastructure. In addition, the environment affects infrastructure as changes in the environment such as the climate emergency can increase the costs of infrastructure (e.g. cost of damages caused by flooding). In some cases, natural capital approaches may reduce the demand for hard engineered infrastructure, such as coastal defences, by supporting the use of green engineering or NbS.
- There are significant risks to natural capital that must be considered. The degradation of natural capital can lead to the loss of ecosystem services that are vital for human well-being and economic stability. For instance, the destruction of wetlands can increase the risk of flooding, while deforestation can exacerbate climate change by reducing carbon sequestration. Ignoring these risks can result in increased costs for disaster management and recovery, loss of biodiversity, and diminished quality of life.

The consideration and implementation of natural capital approaches in planning future flood risk and water management infrastructure investment can provide a wide range of benefits [7]:

- Reducing risk of not fully considering the value of the natural environment (whether monetised or not) in decision making.
- Enabling a holistic cost to benefit analysis and risk assessment.

- Identifying how economic and social activity is dependent on the natural environment.
- Facilitating an innovative approach to identifying policy solutions and managing risk.
- Helping to identify priorities for investment.
- Providing a basis for systematic accounting of changes to natural capital assets over time.

2.3 Key accounting methods

2.3.1 Natural capital accounting

Natural capital accounting is an approach that can be used to provide a systematic and repeatable exercise to collate information on natural capital, the ecosystem services that they provide and the associated market value (if appropriate) [4]. These accounts can be used to measure, value, monitor and communicate the stock and condition of natural assets and the flows of services that they provide. By understanding the state of natural capital this can help ensure that the value of natural assets is considered as part of the decision-making process.

There are a variety of approaches that can be used in natural capital accounting depending on the scale and purpose of the exercise, however the key considerations in each are the extent, condition, physical flow, monetary flow and asset accounts.

2.3.2 Environment Agency Natural Capital Register and Account Tool

The Environment Agency NCRAT was developed to help estimate the flow of services from natural assets and understand the value and benefits that the natural environment provides.

NCRAT requires information on the habitat extent and condition to be provided alongside other local data and uses this information to calculate flows of services and associated monetary values. This information is then summarised in a benefits statement showing the flows of ecosystem services from assets and the calculated monetary values of selected services including present value over time. Figure 4 outlines the types of input data that is required, processes that are undertaken and outputs that are provided by the NCRAT [8].




Figure 4: NCRAT Flow Chart

The NCRAT does not include valuation of supporting services (such as nutrient cycling) as this has the potential to lead to double counting as they underpin cultural, provisioning and regulating services. There are also ecosystem services that are not captured by the current version of the NCRAT due to a lack of robust data. This includes pollination, noise mitigation and soil quality and erosion. Where ecosystem services have been identified as being important, but are not captured by NCRAT, qualitative commentary has been provided to ensure that key benefits are captured.

3. Our approach to understanding the benefit and value of natural capital

3.1 Developing our evidence base

This section outlines the methodology used to assess the baseline natural capital value within the catchment.

3.1.1 Methodology

The natural capital assessment was conducted in six distinct stages, broadly aligned with the Natural Capital Protocol [1] and guidance set out within the Enabling a Natural Capital Approach guidance [4]:

1. Desk-based research on natural capital assets, ecosystem services, benefits and plans.
2. An initial materiality assessment to scope in/out relevant ecosystem services based on the wider context of the catchment, use of resources and how people interact with habitats (see section 3.1.3 for more information).
3. Baseline analysis using the Environment Agency's (EA) Natural Capital Register and Account Tool (NCRAT), qualitative analysis of additional ecosystem services not quantified within the NCRAT assessment, and spatial GIS analysis.
4. Natural Capital Scorecard and associated narrative (including ecosystem services and benefits that were not assessed in the NCRAT but were scoped in during the materiality assessment).
5. Presenting outcomes of spatial analysis, and natural capital assessment. Contextualisation of results by incorporating outcomes of stakeholder engagement and spatial analysis. Qualitative narrative on wider ecosystem services not assessed in monetary or biophysical terms to provide a holistic perspective on natural capital value generated.
6. Interpret results and identify limitations in the assessment.

3.1.2 Desk-based research: Data Sources

The data sources used for the desk-based assessment of natural capital assets and ecosystem services are listed below.

- Corine Land Cover 2018 [9]
- Provisional Agricultural Land Classification (England) [10]
- Statutory Main Rivers [11]
- OS Open Rivers [12]
- OpenStreetMap Waterways [13]
- National Forest Inventory woodland [14]
- England Peat Status GHG and C Storage [15]
- Ancient Woodland Inventory [16]
- Priority Habitat Inventory [17]
- Water Framework Directive (WFD) catchment data – waterbody classifications and extent (rivers, surface waters and groundwaters) [18]

- WFD waterbody extent (rivers, surface waters, transitional and coastal waterbodies, lakes and groundwater bodies) [19] [20] [21]
- Felling Licence Applications [22]
- Surface Water Availability for Water Resource Charging [23]
- Groundwater Management Units coloured according to water resource availability [24]
- Bathing waters data [25]
- Sites of Special Scientific Interest [26]
- Special Protected Areas [27]
- Special Area of Conservation [28]
- Ramsar Sites [29]
- National Nature Reserves [30]
- Local Nature Reserves [31]
- Natural England Green Infrastructure Map [32]
- Outdoor Recreation Valuation (ORVal) Tool [33]
- NCRAT Data Support Package⁵

3.1.3 Materiality assessment

The NCRAT requires a range of local information to be entered into the tool to support the assessment. There are instances where the level of effort required to gather the data for a specific input is disproportionate to the scale of the study and relevance of the data to the scheme. In these instances, the information is excluded from the tool. There are also ecosystem services that are not quantified or given a monetary value by the NCRAT but that may be of importance in the catchment (e.g. pollination or soil erosion/ retention) and therefore require additional analysis to capture the potential benefits and provide a more well-rounded account.

An initial materiality assessment was undertaken to determine which ecosystem services are of direct relevance to the catchment and should be considered as part of the baseline natural capital account (e.g. the ecosystem services that are provided currently). The list of ecosystem services outlined in the Natural England Accounting for National Nature Reserves report [34] was used as the basis to scope services in/out of the assessment.

The results of the materiality assessment, including an overview of the ecosystem services that have been scoped in/out is provided in section 4.1.

3.1.4 NCRAT

The Environment Agency's NCRAT has been used to develop a baseline of the condition and extent of natural capital assets and help understand and quantify the value and benefits that they provide to society. The NCRAT has been selected as it provides a consistent methodology to develop natural capital registers and accounts for each of the catchments within the Fens2100+ project. The NCRAT is also best suited for use at the strategic scale (above 10,000ha) and therefore is most suitable for the size of catchments that are considered as part of Fens2100+. An overview of the NCRAT is provided in 2.3.2 and further guidance on the tool can be found in The Natural Capital Register and Account Tool, Version 1.2 – Technical Report [8].

⁵ This package was provided by the Environment Agency and is not publicly available.

3.1.5 Confidence Ratings

The NCRAT considers thirteen key ecosystem services as shown below in Figure 5.



Figure 5: Ecosystem services quantified and valued in the NCRAT. Environment Agency © [7]

A natural capital register and account will only ever reveal a partial value of nature, however the benefits statement provides an overview of the flows of ecosystem services from assets and the calculated monetary values of selected services including present value over time. Monetary values have been provided for annual flows and for asset value over a 100-year period (minimum and maximum values also provided where data is available).

Confidence ratings have then been assigned to the monetary values presented for each of the ecosystem services to highlight where the data used to calculate a monetary value is considered to be of greater quality or more appropriate. These confidence intervals are assigned based on a Red-Amber-Green (RAG) rating in line with the NCRAT methodology and are illustrated in Table 1 [8].

As highlighted in Section 3.1.1, the number of ecosystem services and benefits provided by a place vary, and not all of them can be quantified. This means that not all services are currently represented within NCRAT.

Table 1: Confidence rating definitions

Confidence rating	Definition
Green	High confidence in results. Input data and assumptions are based on statistical reports, peer reviewed values or industry standard methodologies
Amber	Moderate confidence in results. Input data and/or assumptions from single source/not peer reviewed or based on sources that are not specifically tailored to this context. Some assumptions or estimations may be open to question. Accuracy is expected to better than plus or minus 50%.
Red	Low confidence in results. Input data and/or parameters from a single source and low level of transferability from original context. It is expected that estimate is in the right order of magnitude. Order of magnitude implies that, for example, for an estimate of 5 the 'real' figure is within the range 0.5 to 50.
White	It is not possible to provide a result that can be judged to be in the right order of magnitude. This is due to unquantifiable uncertainty in the science, valuation or the relationship between them. Currently limited to qualitative assessment only.

Confidence rating	Definition
Grey	These ecosystem services are not valued by the NCRAT, either due to a lack of data or because they are too complex for the scope of this assessment.

3.1.6 Stakeholder Engagement

Stakeholder engagement is valuable for identifying appropriate input data and interpreting the outputs from the register and account. Organisations and local stakeholder groups - such as The Wildlife Trusts - may be able to provide, or have access to, locally specific data. They can also offer diverse perspectives on qualitative assessments, including risks, pressures, and the significance of benefits in a particular area.

The qualitative aspects of the assessment, such as the significance assessment, have been based on professional judgement and understanding of the catchment. It is recommended that this be refined through engagement with stakeholders as part of any future update and project progression.

3.1.7 Limitations and assumptions

The following assumptions should be noted, in addition to those provided in the NCRAT [7].

- All values within the analysis were expressed in 2024 prices and British Pound (GBP).
- The 'Risk Register,' 'Significance Assessment,' and 'Beneficiaries Assessment' tabs in the NCRAT spreadsheet tool have been completed without detailed stakeholder engagement having been undertaken. It is recommended that this be refined through engagement with stakeholders as part of any future update and project progression. However, the inputs for these assessments do not impact the overall natural capital value calculated for the study area.
- Statutory main river widths are assumed to be 4m and ordinary watercourses, which are mapped using both the OS Open Rivers and OpenStreetMap datasets, are assumed to be 3m wide, respectively⁶. It should be noted that there are a large number of man-made drainage ditches and channels across the catchment that are not accounted for in this analysis.
- Woodland areas under management are assumed to be areas with a felling licence expiry dated after 31st March 2015, this is based on the Forest Commission's definition of 'Woodland area under management' having had a felling licence in the past ten years, being owned by government organisations, or having a management plan [35].
- Peatland condition classifications are assigned based on the estimated rate of GHG flux (named 'newghgr' in the 'England Peat Status GHG and C storage dataset') and the closest emission factor provided in the Peatland Code Field Protocol [36].
- To calculate input values for water supply and renewable energy, local authority data was used and adjusted in proportion to size of the catchment. This assumes that the distribution of renewable energy and water supply is even across the local authority.

⁶ Based on a sample of 15 river widths taken from across the catchment boundary. Measured using aerial imagery using ArcGIS.

- To estimate the number of recreational visits in the catchment and the associated value of this, ORVal data was used for each of the Middle layer Super Output Areas (MSOAs) in the catchment. Where an MSOA is only partially within the catchment, the number of visits and associated value was scaled down relative to the area within the catchment – this assumes that the distribution of recreational visits for each welfare group is even across the MSOA.

The following limitations should also be considered:

- Data resolution is generally low; with the majority of habitat data being derived from remote sensing datasets such as CORINE Land Cover or the Centre for Ecology and Hydrology Land Cover Maps being the main data sources for the assessment. Datasets such as the National Forest Inventory, EA Statutory Main Rivers, OS Open Rivers and OpenStreetMap datasets were used to supplement habitat type data and reduce this limitation.
- The provision of biodiversity benefits is currently not quantifiable within the tool (for example when comparing arable land to more biodiverse semi-natural habitats).
- The net carbon sequestration rate for arable land is based on Christie et al. 2011 at 0.107tCO₂e/ha/yr, resulting in a sequestration of carbon [37]. More recent data collated in the Natural England report 'Carbon Storage and Sequestration by Habitat 2021 (NERR094)' uses figures from a study by Muhammed et al. 2018 which showed a net emission rate of 0.290tCO₂e/ha/yr for arable land and a net sequestration rate of 0.360tCO₂e/ha/yr for improved grasslands [38]. There is variation in carbon fluxes on arable land based on a variety of factors, including production and application of fertilisers, vegetation type, catchment topography, weather, and land use. However, it is likely that the 0.107tCO₂e/ha/yr figure does not provide an accurate picture of carbon fluxes within the catchment.
- The area of freshwater habitats is likely to be underrepresented in this study due to the large number of drainage channels in the catchment that are not captured in the EA Main River, OS Open Rivers datasets or OpenStreetMap datasets.
- The 'England Peat Status GHG and C storage dataset' is expected to be an overestimate of peatland extent in the Fens2100+ study area, based on a comparison with more recent data published by Natural England [39]. This indicates the amount of carbon dioxide emitted as a result of actively eroding or degraded peatland has also been overestimated, so the monetary value of climate regulation in the catchment is expected to be greater than calculated for this baseline.
- It is likely that these limitations lead to an underestimate of the benefits provided by semi-natural habitats, and furthermore underestimate the negative impacts of arable land while overemphasising the wider benefits of arable land, beyond food production.
- The NCRAT uses national data to calculate the natural capital value for agriculture. This is based on the average of the John Nix Farm management units Editions 49-52, years 2019-2022 [40]. The natural capital value is updated for the 2024 accounting price year which takes account of inflation to the price base. Local deviations from the national average may lead to the value of local agriculture being misrepresented. In the catchment, there is a higher proportion of arable agriculture and fruit / vegetable crops than the national average, meaning that the NCRAT is likely to underestimate the value of food production.
- The NCRAT values water quality based on all the WFD/WER surface water bodies in the operational catchments that the catchment intersects (Fens East and West and

Steeping and Eaus). There are sixteen surface water bodies across these operational catchments, however only three of these fall within the catchment (Lymn/ Steeping; Cow bank drain and the Wast Outer) boundary. All of these waterbodies score 'moderate' for ecological status for the River Basin Management Plan Cycle 2, and Cycle 3. This is therefore likely to be an overestimation of the value of water quality in the catchment.

- This baseline assessment has not been informed or verified by any field surveys or site visits conducted by Arup.
- This baseline assessment is currently limited to the catchment boundary that has been defined for the Fens2100+ project by the Environment Agency. There is no official data boundary that matches the catchment study area.
- The baseline collated comprises a proportionate level of detail from publicly available data sources and it is reported at the geographical basis relevant to the data availability. Therefore, this report reviews and provides commentary of existing reports written by others. Arup accepts no responsibility for any error or omission in this report which is due to an error or omission in the original documentation we refer to. Arup have not independently verified the information and have assumed it to be accurate, complete, reliable, and current as of the date of such information.

4. The natural capital register and account for the catchment

4.1 Materiality Assessment

A summary of ecosystem services to be scoped in/out of the assessment is provided in Table 2 and the full materiality assessment is provided in Appendix A. The initial materiality assessment highlighted that the majority of ecosystem services are of significance and should therefore be scoped into the assessment. Of those services that have been scoped out, this is predominantly due to lack of marine habitat (for fish and shellfish) and woodland habitat (for foraging and noise regulation). Of those that have been scoped into the assessment, six are not quantified by NCRAT and require additional qualitative analysis. These are indicated using ‘*’ in Table 2.

Please note:

- Ecosystem services related to wild plants, algae & their outputs, wild animals & their outputs, and aquaculture have been scoped out due to lack of reliable data on gathering rates and/or lack of suitable habitat for these benefits to materialise (e.g. availability of coastal waters for fishing and/or aquaculture, presence of ports that record fish landings, availability of land used for commercial hunting, datasets on foraging rates).
- Noise regulation benefits were scoped out of the assessment, as most approaches to quantifying benefits are based on complex modelling that requires significant collection of location-specific data.

Table 2: Materiality Assessment summary

Scoped in	Scoped out
<ul style="list-style-type: none"> • Materials from plants, animals & algae • Cultivated crops • Water supply • Reared animals & their outputs • Water quality • Air quality • Mass stabilisation and control of erosion rates* • Flood protection • Global, regional & local climate regulation • Pollination & seed dispersal* • Maintenance of nursery populations and habitats* • Pest & disease control* • Global, regional & local climate regulation • Experiential and physical use • Scientific and educational use • Aesthetic* • Spiritual* 	<ul style="list-style-type: none"> • Wild animals & their outputs • Wild plants, algae and their outputs • Aquaculture • Noise regulation

4.2 Natural Assets

4.2.1 Land cover types in the catchment

This section focuses on providing an overview of the quantity, quality and location of natural capital assets within the catchment, which contains approximately 6,340ha of terrestrial and coastal habitats. Figure 6 shows land cover in the catchment according to Corine Land Cover 2018 [9].




Figure 6: Overview of Corine Land Cover for the Steeping catchment.

For the purpose of this report, the natural capital assets in the catchment have been grouped according to UK National Ecosystem Assessment (UKNEA) broad habitat types. The quantity of each habitat type in the catchment is shown in Figure 7. The Office for National Statistics UK Natural Capital Accounts methodology guide provides further detail on habitat types and the methods used to estimate habitat extent, ecosystem services and asset value in the UK [41].




Figure 7: Natural capital asset quantity within the catchment. Based on the UK Natural Capital Ecosystem Assessment broad habitat types / land cover data.

4.2.1.1 Coastal margins

The catchment has approximately 5km of coastline along the North Sea and 225ha of coastal saltmarsh. The catchment intersects with the northern coastline of The Wash, an intertidal embayment designated as a Ramsar site, Special Protection Area, Special Area of Conservation, and Site of Special Scientific Interest. The site is of significant importance due to habitats such as sublittoral sandbanks, sandflats, mudflats, shallow inlets and reefs which attract a range of wildlife including wintering waterbirds; passage birds; waders; breeding birds; harbour seal *Phoca vitulina*; and shellfish [42] [43] [44].

These coastal margins in the catchment primarily include priority habitat coastal saltmarsh, coastal sand dunes and reedbeds [45]. Coastal habitats protect against erosion and flooding by dissipating waves, trapping sediment, and creating buffer zones that store water during storms. Additionally, they provide various ecosystem services, including water quality improvement, carbon sequestration, biodiversity, recreation and tourism, cultural and historic value, and serve as nurseries for many commercial fish species.

4.2.1.2 Enclosed farmland

Enclosed farmland or agricultural land is essential for food production and provides local employment. Additionally, it can support a variety of wintering and breeding birds, as well as provide habitat connectivity through hedgerow and ditch networks for a variety of other species.

The most abundant land cover type in the catchment is enclosed farmland, covering an area of 5,894ha - 93% of the catchment. Around 3,350ha (56%) of land is classified as Grade 1 and 2 land using the Agricultural Land Classification⁷, with the majority of the Grade 1 land located within the area south of the Wainfleet. Figure 8 provide an overview of Agricultural Land Classification within the catchment [36].

According to 2018 Corine Land Cover, 82% (4,835ha) of farmland is used for arable agriculture or horticulture (includes cereal crops, fruit and vegetables and other arable crops), including the limited production of some higher value crops in rotation such as potatoes and sugar beet. In 2021, this was estimated to be approximately 80% based on estimated land use, crop areas and livestock populations on commercial agricultural holdings [46].

Approximately 18% (1,060ha) is modified grassland used for the keeping of livestock, with a number of farms producing beef cattle and/or sheep - particularly in the central and northern parts of the catchment where land quality is lower due to poor drainage. In 2021, this was estimated to be approximately 16% based on estimated land use, crop areas and livestock populations on commercial agricultural landholdings.

It should be noted that since 2019 there has been significant change and market disruptions in agriculture and the food supply, notably the Agricultural Transition following UK exit from the EU. The Agricultural Transition includes the phase out of subsidies such as the Basic Payment Scheme, and the introduction of the Environmental Land Management Scheme and associated pilots which is influencing the type and extent of cropping across the catchment and beyond. Additionally, cropping regimes and agricultural output per hectare on an annual basis is also extremely variable from crop to crop and farm to farm, with impacts such as climate change and extreme weather also influencing agriculture in the catchment.

⁷ Grade 1 agricultural land is excellent quality agricultural land with no or very minor limitations for growing agricultural and horticultural crops.

Grade 2 agricultural land is very good quality land with minor limitations that affect crop yield, cultivations or harvesting.

Grade 3 agricultural land is good quality land capable of consistently producing moderate to high yields of a narrow range of arable crops, especially cereals, or moderate yields of other crops including oilseed rape, potatoes and sugar beet.




Figure 8: Overview of Agricultural Land Classification within the catchment.

4.2.1.3 Urban areas

Urban land is generally characterised by significant alteration from semi-natural habitats caused by humans. Nonetheless, urban land can contain a variety of natural and/or semi-natural natural capital assets in the form of blue or green urban infrastructure. Most commonly, these consist of parks, urban trees, gardens, and allotments. Certain types of green infrastructure such as parks are generally open to the general public and can be crucial in providing a variety of cultural ecosystem services, such as benefits for health and wellbeing, providing aesthetic value and cultural heritage.

The key urban centres in the catchment consisting of built-up areas and gardens are the villages of Wainfleet All Saints, Irby in the Marsh, Croft, Thorpe St Peter and Wainfleet St Mary.

There are a number of designated heritage assets located within the vicinity of urban areas, and two conservation areas within the catchment. For the historic environment, natural capital benefits can be attributed to different aspects of the environment resulting from the interaction between people and places through time, including all surviving physical remains of past human activity, whether visible, buried or submerged, and landscaped and planted or managed flora. Key benefits associated with historic environment assets include amenity, landscape and non-use values, with beneficiaries likely to include local communities and wider society that visit.

Additionally, a large number of non-designated heritage assets are present throughout the catchment including as yet undiscovered archaeological remains, which are likely to be particularly well preserved within peat soils.

4.2.1.4 Woodlands

Woodlands are essential for carbon sequestration, biodiversity, recreation, and economic value, significantly contributing to the country's natural capital. They also provide other vital ecosystem services such as soil formation, water regulation, and air purification. These services not only support environmental health but also enhance human well-being by improving air quality and reducing the impacts of climate change. Additionally, woodlands offer recreational opportunities that promote physical and mental health, and they support various economic activities, including timber production and tourism.

The catchment consists of approximately 60ha of woodland, predominantly made up of mixed deciduous trees (21ha) and commercial woodland that is actively managed (20ha) with a felling licence valid within the last 10 years [14]. There is also a small amount of coniferous woodland in the catchment (5ha). No ancient woodland is present.

4.2.1.5 Freshwaters and groundwaters

Freshwater habitats are crucial for supplying drinking water; managing flood risk; supporting a wide variety of activities that benefit health, wellbeing, and economic activity; and can be a significant carbon sink. The east of England is also particularly reliant on water for the irrigation of crops, particularly in the months between June and August; agriculture in this region accounts for approximately 10% of the total water demand [47] .

Freshwater habitats, including rivers and streams, comprise approximately 21ha within the catchment, however this is likely to be under representative due to the large number of drainage channels managed by land owners and IDBs that are not captured in the EA Main River, OS Open Rivers datasets or OpenStreetMap datasets.

The catchment is located within the Anglian River Basin District (RBD). The majority of the catchment falls within the lower reaches of the 'Lymn / Steeping' river catchment. This is

located within the Steeping and Eaus operational catchment, which in turn forms part of the Environment Agency's Witham management catchment. Some areas along the eastern extent of the catchment fall into the western edge of the neighbouring 'Cow Bank Drain' river water body catchment area, which is located within the same operational catchment. A portion of the far south-eastern extent of the catchment falls within 'Wash Outer' coastal water body extent, which is located within The Wash Transitional and Coastal (TraC) operational catchment, as part of the wider Anglian TRaC management catchment area. Section 4.4 of the Fens2100+ Steeping: Environment and Agriculture Appendix provides an overview of the Cycle 3 (2022) status classifications for surface water bodies within the catchment. The location and extent of these surface water bodies and catchments are shown in Figure 9.

The catchment is underlain by 4,782ha of groundwater bodies, namely the Steeping Long Eau Little Eau Chalk Unit, and the Steeping Great Eau Long Eau and Witham Spilsby Sandstone Unit. This is a crucial source of water for public use, agriculture and industry.




Figure 9: WER surface waterbodies in the catchment

4.2.1.6 Mountains, moorlands and heaths

Dwarf shrub communities typically support a range of bird species and can provide suitable habitat for all reptile species found in England. Coastal dwarf shrub heath communities provide crucial ecosystem services such as carbon sequestration, water regulation, soil protection, and support for biodiversity. They also offer cultural and recreational value and support a range of pollinators.

Approximately 16ha of heathland and shrub habitat comprising, dwarf shrub heath, hedgerows and dense scrub is present. Dwarf shrub heath is predominantly located in coastal areas, as transition communities on sand dunes, forming part of designated sites including Gibraltar Point SSSI, SPA, Ramsar site and National Nature Reserve (NNR).

4.2.2 Peatland in the catchment

Peatlands are wetland areas with a natural layer of peat soils, formed from carbon rich dead and decaying plants under waterlogged conditions, they provide a variety of crucial services to society [48]. Firstly, peatlands represent an important carbon sink globally and have the potential to regulate atmospheric carbon when in a healthy waterlogged condition. However, once peats are in a degraded condition they become carbon sources – this is the case for the majority of peatlands in the UK [49]. Peatlands also provide habitat for a diverse range of species, from *Sphagnum* mosses to butterworts *Pinguicula* to common crane *Grus grus* [48]. Healthy peatlands provide flood management by retaining water and slowing the flow, and provide important sources of high-quality water that can be easily treated for drinking water [48].

According to Natural England's dataset 'England Peat Status GHG and C storage' (2008), the catchment contains approximately 62ha of peatland. In contrast to Natural England's peatland data, Corine Land Cover data classifies areas with indicative peat presence as 'modified grassland' and 'arable and horticulture', which is indicative of peat degradation.

Furthermore, there is uncertainty and inconsistencies in data available on extent and condition across the wider Fens. Holman and Kechavarzi, estimate that less than 32,000ha of peatland (thick peat, peat at depth, or thin peat), was still extant in the Fens as a whole, compared to approximately 150,000ha identified as such by the peaty soils locations layer [50].

The Fens for the Future report [51] explains how peatland in the Fens has been degraded through drainage for arable cultivation. Once degraded, peatland emits CO₂ and methane; a report on peatland natural capital by the Office for National Statistics estimates that 7,600 kilotons of carbon dioxide equivalents per year (kt CO₂e/yr) are emitted as a result of arable practices on peatland [52].

The baseline condition of this peatland is categorised as 'drained' using the peatlands emission factor (tCO₂e/ha/yr), in accordance with the Peatland Code Field Protocol [36]. For peatland in a 'drained' condition the assumed emission factor is 4.54tCO₂e/ha/yr.

Natural England recently released the 2025 England Peat Map [39]. This provides a more accurate and detailed picture of the current state of peatland when compared to the 2008 England Peat Status dataset [15] that has been used for this Natural Capital Baseline. The new peat map has been developed using machine learning, satellite imagery, LiDAR, and updated field surveys to detect peat soils that were previously unmapped or misclassified. The new 2025 peat map shows both gains in mapped extent (due to better detection) and losses or degradation (due to environmental change).

The 2025 England Peat Map suggests that the peatland extents used in this baseline (based on the 2008 'England Peat Status GHG and C storage' dataset) are an

overestimation. The new peat map shows approximately 78,000ha of deep peaty soils across all catchments in the Fens2100+ study area, whereas the 2008 dataset shows approximately 155,000ha. There is approximately 70,000ha of intersecting peatland extents between the two datasets, highlighting that the new peat map has identified roughly 8,000ha of previously unmapped peatland across the Fens2100 study area. For climate regulation, this indicates the amount of carbon dioxide emitted as a result of actively eroding peatland has also been overestimated, so the value of climate regulation is expected to be greater than calculated for this baseline.

As the Natural Capital baseline assessment uses the 2008 'England Peat Status GHG and C storage' dataset, percentage coverages of peat are not directly comparable with the catchment summary report (which uses the 2025 England Peat Map).

4.2.3 Protected areas for biodiversity

In 2019, Natural England produced a natural capital account of all National Nature Reserves managed by Natural England, and highlighted the value of National Nature Reserves for biodiversity protection and landscape enhancement [1].

There are ten statutory internationally or nationally designated nature conservation sites within or partially within the catchment, some of which overlap with other designations. These comprise:

- **Gibraltar Point** – Special Protection Area (SPA), Ramsar site, National Nature Reserve (NNR) and Site of Special Scientific Interest (SSSI);
- **The Wash** – SPA, Ramsar site and SSSI;
- **Saltfleet by Theddlethorpe Dune & Gibraltar Point** – Special Area of Conservation (SAC), NNR and SSSI; and
- **Bratoft Meadows** – SSSI.

The location of each designated site is shown in Figure 10, and a summary of each site provided in sections 4.2.3.1 – 4.2.3.4.




Figure 10: Overview of designated sites for nature conservation within the catchment

4.2.3.1 Gibraltar Point (SPA, Ramsar, NNR and SSSI)

Gibraltar Point covers roughly 35ha of the catchment. The site is an actively accreting dune system with extensive sand dunes, salt marshes, freshwater pools, freshwater marshland and areas of grassland. The area is highly important for bird wildlife, supporting nationally significant breeding or wintering populations of a wide range of species, some examples include little tern; bar-tailed godwit; sanderling *Calidris alba*; grey plover and oystercatcher [53] [54]. It is also important for rare wetland invertebrate species including red data book species including endangered saltmarsh knot-horn *Gymnacyla canella* and a fungus gnat species, and eight species listed as rare including marsh moth and scarce pug [55].

4.2.3.2 The Wash (SPA, Ramsar and SSSI)

The Wash covers approximately 215ha of the catchment. The Wash SPA forms part of the larger Wash and North Norfolk Coast Special Area of Conservation which encompasses the largest embayment in the UK, as well as extensive intertidal sand and mudflats, subtidal sandbanks, biogenic and geogenic reef, saltmarsh and a barrier beach system unique in the UK. The Wash SPA supports a large number of non-breeding bird species such as: bar-tailed godwit; Bewick's swan; and black-tailed godwit [42] [43].

The Wash Ramsar site is Britain's largest estuarine system, fed by four rivers and has extensive saltmarshes, intertidal banks, and deep channels. This aspect of the site is crucial for wildfowl and wading birds, supports a shellfish fishery, and hosts large populations of common and grey seal [44] [55].

4.2.3.3 Saltfleet by Theddlethorpe Dune & Gibraltar Point (SAC, SSSI and NNR)

Saltfleet by Theddlethorpe Dune & Gibraltar Point covers approximately 33ha of the catchment. The site comprises two dune systems including shifting dunes and fixed dune features. Some of the specific features it is designated for include: shifting dunes along the shoreline with European beachgrass (white dunes); fixed coastal dunes with herbaceous vegetation (grey dunes) - Annex 1 priority habitat; dunes with sea buckthorn humid dune slacks and embryonic shifting dunes [56].

Saltfleetby-Theddlethorpe Dunes also supports natterjack toad with active management to support breeding, it is the most north-easterly breeding site for this species. The intertidal sands and muds provide feeding and roosting grounds for breeding and wintering wildfowl and waders, while the saltmarsh communities host species like glasswort *Salicornia* and sea lavender *Limonium* [55]

4.2.3.4 Bratoft Meadows (SSSI)

Bratoft Meadow covers approximately 2.1ha of the catchment. The site is one of the best examples of species-rich neutral grassland in Lincolnshire. The meadow is dominated by sweet vernal grass *Anthoxanthum odoratum*, red fescue *Festuca rubra*, meadow fescue *Festuca pratensis*, and creeping bent *Agrostis stolonifera*, with a rich sward featuring betony *Betonica officinalis*, dyer's greenweed *Genista tinctoria*, cowslip *Primula veris*, adder's tongue fern *Ophioglossum vulgatum*, and saw-wort *Serratula tinctoria* [55]. Wetter areas have cuckooflower *Cardamine pratensis* and ragged robin *Lychnis flos-cuculi*, and the southern field has abundant green-winged orchid *Anacamptis morio* [55]. A third field, separated by a narrow strip of improved pasture, includes yellow rattle *Rhinanthus minor*, great burnet *Sanguisorba officinalis*, and devil's-bit scabious *Succisa pratensis*, and is also

cut for hay [55]. The site attracts large numbers of butterflies and has 18 recorded species of terrestrial mollusc [55].

4.2.4 Risk Register

The NCRAT includes a Risk Register that highlights how drivers of environmental change may present risks to the quality and quantity of natural assets in the study area. This can be used to highlight which ecosystem services are under pressure and limited in their capacity to continue to provide future benefits.

The assessment of risk is qualitative and informed by the prevalence of the driver of environmental change, as well as the extent and condition of the natural asset in the study area. The drivers of change assessed in the NCRAT are described in Table 3.

Table 3: Drivers of environmental change considered in the Risk Register

Driver of change	Definition
Land-use change (incl. habitat loss)	This can be the result of a number of drivers including development and coastal erosion as well as hazards (e.g. forest fires). Land-use change is usually permanent but it could also be temporary. It is likely to affect the extent of natural capital asset rather than condition.
Climate change	This includes drivers on habitats from changing climatic conditions including temperature, precipitation, soil aridity and storminess. Assessments of the likely impact of climate change over this century are provided in UK Climate Projections ⁸
Resource extraction (incl. over-exploitation)	Over-exploitation of resources can impact natural assets through both direct physical damage and additional stress on the resources on which they rely (e.g. water availability).
Population growth	Driver which comes from the demands of a higher population on natural resources and the environment. Note that the impact of population growth may be through future land-use change, resource extraction, and pollution.
Pollution	Includes both diffuse sources (e.g. agriculture, urban run-off, atmospheric deposition) and point sources (e.g. sewer overflows, wastewater treatment, abandoned mines).
Invasive species/diseases/pathogens	Invasive species/diseases/pathogens can have significant impacts on ecosystems, threatening the abundance and health of native species.

⁸ UK Climate Projections are available at: <https://www.metoffice.gov.uk/research/approach/collaboration/ukcp/index>

For this catchment, the natural assets under most pressure are coastal margins; enclosed farmland; freshwaters; and mountains, moorlands and heaths. The primary drivers of this are land-use change (including habitat loss), climate change, resource extraction and pollution. Figure 11 shows which ecosystem services are most at risk due to strain on a natural asset.

Level of pressure categories	Description
High	High likelihood the pressure will affect natural asset(s) and high risk of loss of ecosystem service.
Medium	Medium likelihood the pressure will affect natural asset(s) and medium risk of loss of ecosystem service.
Low	Low likelihood the pressure will affect natural asset(s) and low risk of loss of ecosystem service.
None	The pressure is not expected to affect the natural asset(s) nor ecosystem service.
No data	No data available to assess likelihood of pressure and affect on natural asset(s) nor ecosystem service.
Not assessed	Risk rating not completed
	No viable benefit-asset attribution

Ecosystem Services Group	Natural capital assets (NEA Broad habitat type)							
	Coastal margins	Enclosed farmland	Freshwaters - Open waters, wetlands and floodplains	Marine	Mountains, moorlands and heaths	Semi-natural grasslands	Urban	Woodlands
Agriculture		High			High	None		
Fish and shellfish landings	High			None				
Water supply			High					
Timber								Medium
Renewable energy		High	High		High			
Climate regulation	High	High			High	None		Medium
Air quality	High	High	High		High	None		Medium
Hazard regulation								
Recreation	High	High	High		High	None	Medium	Medium
Physical Health	High	High	High		High	None	Medium	Medium
Education		High	High		High	None	Medium	Medium
Volunteering		High	High		High	None	Medium	Medium
Water quality		High	High		High	None	Medium	Medium

Figure 11: Natural Capital Risk Register

4.3 Ecosystem Services

The type and significance of ecosystem services in the catchment are shown in Figure 12. More information about each of these services is provided in the sections below.

As shown in Figure 12, the most significant ecosystem services in the catchment are water supply, arable agriculture, renewable energy, climate regulation and hazard regulation. The significance is based on a desk-top assessment of the relative provision of ecosystem service and the amount of people it will impact. The definition of each significance rating is shown in Table 4.




Figure 12: Rose diagram showing the significance of ecosystem services in the catchment.

Table 4: The definition of the ecosystem service significance ratings used in the NCRAT.

Significance rating	Definition
Highly significant	The benefit flow is (very) important to the place in relative terms* because it impacts the wellbeing of a significant proportion (and possibly a wide variety) of people and businesses and/or due to wider social and cultural aspects.
Significant	The benefit flow is of material importance in relative terms* because it impacts the wellbeing of a sizeable proportion of people and businesses.
Minor	The benefit flow is of minor importance in relative terms* because it does not impact most people and business. Note: it may, though, still be highly important to a small number of specific individuals or groups.
Not significant	There is very little or no provision of this benefit and it impacts only a very small number of people and businesses.

*Relative to the provision of other services/benefits within the catchment.

4.3.1 Provisioning Services

Food production

Agriculture plays an important role as an ecosystem service and land-use in the catchment. The majority of terrestrial habitat is comprised of arable fields and agricultural grasslands. Agricultural practices, such as arable farming and livestock rearing, contribute to national food security and support the local economy.

Water supply

The catchment is located above a nationally significant chalk aquifer spreading from the south-east of Yorkshire, southwards across the Humber and into Lincolnshire. Acting like a sponge, chalk aquifers hold rainwater which can emerge at ground levels from springs, facilitating the supply of drinking water to the local population.

The Steeping, Great Eau & Long Eau Abstraction Licensing Strategy (ALS) [57] covers waterbodies within the catchment. This shows that the whole catchment area at Q30 flows has available water⁹. At Q50 flows, the catchment has primarily restricted available water, with only the catchment area between Croft and Skegness having water available at this flow level.

Water is only available for abstraction from the Lower River Steeping on 138 days of the year. Notably, the total water available for abstraction per day across the entire catchment

⁹ The Environment Agency's Water Resources Geographical Information System provides a snapshot in time on water availability at four flow snapshots. From high to low, these are: Q30 (ie. the flow of a river which is exceeded on average for 30% of the time, therefore you would expect the river flow to be lower than Q30 on 256 days in an average year, i.e. a higher flow), Q50, Q70 and Q95. These are calculated for each integrated Water Framework Directive waterbody and looks at the balance between the flow in the river, the quantity needed to support the ecology and the water that can be licensed for abstraction.

is just 12,300m³, sufficient to provide irrigation of c.25mm across c.0.5km² of land. Consequently, irrigated crops can only be produced in this catchment reliably using water from the on-farm reservoirs which are in the area towards The Wash.

The catchment has a long history of flooding which impacts both properties and farming. Recent events include:

- On 12th June 2019 significant rainfall led to overtopping of embankments on the Wainfleet Relief Channel, and a series of breaches resulted in extensive flooding in the catchment. 85 houses were internally flooded, nearly 600 homes evacuated and 5.5km² of farmland flooded (circa 10% of the farmland in the catchment).
- Winter 2023/24 was at the centre of the wettest 18 months on record for Lincolnshire. On 23rd October 2023 Storm Babet caused widespread flooding in Lincolnshire with over topping in Wainfleet.

Flooding has major impacts on crops and to a lesser extent areas of land utilised for livestock grazing. Flood events can restrict the ability to establish new crops, reduce crop yields and quality, or can destroy crops once ready for harvest leading to direct agricultural damages and wider economic consequences for landowners.

Renewable energy

Many natural processes provide a renewable energy source that can be harnessed to produce electricity. Solar power (photovoltaics) and onshore wind generation are the main sources of onshore renewable energy generation in the catchment.

Timber

Woodlands play a fundamental role in providing timber and wood fuel in the UK. There is a small proportion of woodland cover in the catchment, with approximately 60ha of woodland. There are approximately 20ha of commercial woodland that is actively managed with a felling licence valid within the last 10 years [14].

4.3.2 Regulating Services

Climate regulation

Ecosystems regulate a wide range of atmospheric processes that control our climate, including the greenhouse effect, precipitation, the ozone layer, the albedo effect, and atmospheric chemical composition [58]. Locally, this can help reduce urban temperatures and remove air pollutants. Globally, the most significant example of climate regulation by ecosystems is the absorption of greenhouse gases [58].

Some of most significant habitats for carbon sequestration in the catchment are located along the coastline, such as saltmarshes and intertidal mudflats. A major source of carbon emissions identified in the NCRAT are habitats located on degraded peatland, such as agricultural fields on deep peat soils.

As outlined in section 3.1.7, the net carbon sequestration rate for arable land is based on Christie et al. 2011 at 0.107tCO₂e/ha/yr, resulting in a net sequestration of carbon [37]. More recently, it has been estimated that arable land may in fact emit 0.290tCO₂e/ha/yr [38]. Therefore, it is possible that the assessment of the carbon balance for the catchment is significantly different from the true figure.

Air quality

Most urban areas in England experience significant levels of air pollution which can contribute to adverse health impacts including premature mortality from cardiovascular and respiratory causes. Studies have shown that vegetation can play an important role in removing air pollutants and improving air quality. There are no AQMAs within the Catchment or within the East Lindsey District Council local authority area.

Air pollution removal occurs through dry deposition, a mechanism by which pollutants are accumulated on leaf surfaces. Gaseous pollutants can also be removed through leaf stomata uptake.

Hazard regulation

Many habitats can act as buffers to reduce the effect of natural hazards, such as flooding. Relative to bare soil or paved surfaces, natural habitats reduce flooding risk to downstream populations by buffering peak rainfall flows entering rivers. Likewise, coastlines with intact seagrass and kelp beds, coastal wetlands or mudflats are better protected from flooding and coastal erosion.

The coastal habitats present in the catchment dissipate wave energy and provide barriers reducing risk of damage to coastal defences, low lying land, and infrastructure [59]. If unimpeded by coastal development, natural intertidal habitats such as saltmarsh are likely to migrate with rising sea levels predicted under future climate scenarios, thereby providing resilience to future sea level rise.

Disease and pests

Pests can be responsible for agricultural losses and encourage the use of pesticides. Natural pest control, such as by bats and birds, can enhance crop production by reducing herbivorous insects that damage or use parts of crops. The provision of habitats, such as semi-natural grasslands and woodlands, can support species which prey on pests. Given the high cover of agricultural land in the catchment natural pest control of infestations is an important regulating service.

Pollination and seed dispersal

Semi-natural habitats such as grasslands, woodlands, hedgerows, and scrub support insect populations that pollinate crops growing on agricultural land. Seed dispersal allows plants to colonise new areas where the species may not already be present, ensuring genetic diversity and supporting other species by providing food, shelter, and habitat.

Soil retention and erosion

Soil retention is required for the maintenance of healthy ecosystems and delivery of other ecosystem services such as food and timber production. Soil resources in the catchment are degraded and eroded due to drainage and tillage of land for agriculture. Tree planting and increased diversity of vegetation structure and cover can help increase soil retention.

4.3.3 Cultural Services

Our cultures, knowledge systems, religious beliefs, social interactions and amenity has been shaped by our interaction with natural ecosystems. Cultural ecosystem services represent a suite of benefits related to amenity and culture, that can have a significant impact on society as a whole and are therefore often difficult to quantify or value.

Recreation

The catchment is comprised of a significant coastline, with a large proportion of protected and designated sites in the near vicinity of popular seaside towns and resorts, such as Skegness.

Based on datasets used for ORVal, the catchment contains a total of 11 recreational sites. The most common type of recreational site is grave yard (six sites), other sites include one recreation ground, one allotment, one cemetery, one nature park, one park, and one wood. ORVal data also shows that the catchment includes 12 paths.

These sites attract a range of visitors to the local area which contributes to the local economy.

Physical health

Approximately half of the visits made to the local area are likely to partake in activities that are beneficial for their physical health. Improvements to health improve the quality of life for individuals and reduce the pressure on healthcare systems

Education and volunteering

Visits to the catchment for scientific and educational purposes are difficult to quantify, but can create benefits for human society as a whole. The protected areas and heritage sites within the catchment have the potential to enable insights about past and future landscape changes in the area.

Designated sites provide an opportunity for public volunteering, which provides benefits to recipient organisations and their stakeholders, to wider society, and to the volunteers themselves [34].

Aesthetic / sense of place

The concept of sense of place in the context of natural capital refers to the emotional and symbolic meaning that the environment holds for individuals or communities. Intrinsically, it is difficult to measure, as it describes how people perceive, feel about, and interact with their environment.

In the context of this study, aesthetic / sense of place benefits of natural capital assets is based on the character of the landscape which includes rolling sand dunes, coastal towns/villages, agricultural fields, meandering watercourses and sandy beaches.

4.3.4 Bundled services

Bundled services are benefits provided by nature that are not easily attributable to specific ecosystem services as they provide multiple types of benefits. Some of the bundled services can also be classified as supporting services, they are crucial to the functioning of ecosystem processes and therefore support the provision of other ecosystem services. Bundled services are typically difficult to quantify as their primary benefit is the underpinning of all other ecosystem services, and therefore do not provide a direct measurable benefit.

Water quality

Ecosystems and vegetation can improve water quality in variety of ways, including through the absorption of excess nutrients; slowing the flow of water which increases water (and pollutant) retention in the soil, helping to prevent sewer overload; providing shade and reducing algae growth; and helping to stabilise river banks and beds with root systems, which reduces sediment runoff [60].

The bundled ecosystem service for water quality entails a range of benefits related to ecosystems functioning to remediate polluted water, the value of good-quality water for economic activities, recreation, education, and for healthy plant and animal populations.

As outlined in Section 4.2.1.5, of the three surface waterbodies in the catchment, all score 'moderate' for ecological status for the River Basin Management Plan Cycle 2 and Cycle 3.

Biodiversity

Biodiversity is defined by the Convention of Biological Diversity as “the variability among living organisms from all sources including, inter alia, terrestrial, marine and other aquatic ecosystems and the ecological complexes of which they are part; this includes diversity within species, between species and of ecosystems” [61]. Biodiversity is an integral characteristic of stable and resilient ecosystems, and is vital for the continued delivery of all other ecosystem services.

As described in Section 4.2.3, there are designated sites in the catchment that are internationally and nationally important for biodiversity. These areas include a variety of designated and rare habitats and species - with habitats ranging from salt marshes to meadows to deciduous woodland; and species ranging from oystercatcher to natterjack toad to the endangered saltmarsh knothorn.

4.4 Benefits

4.4.1 Natural capital benefits

Figure 13 shows the flow of ecosystem services from natural assets in the catchment. For each of the ecosystem services shown, data on the physical flow of benefits is used to estimate a monetary value. Sections 394.4.2 - 4.4.5 below describe the physical benefits from the ecosystem services in the catchment. It should be noted that the benefits of ecosystem services impact different groups of people differently, depending on the service and location. Often, ecosystem services will deliver direct benefits to local people, for example through improved health, and indirect benefits to the wider population, for example by reducing cost to health services. The monetary values shown have been calculated from the NCRAT.

Given the catchment context, and the overall value that the Fens contributes to the total agricultural production in the England (estimated to be more than 7% of England's total agricultural production, with a value of £1.23 billion in 2019 [62]), it is assumed that the majority of the benefits of this food production are to the national economy and farmers within the catchment. The majority of the recreational benefits and hazard regulation are assumed to be local to those that visit, live and work within the catchment.



Natural Assets



Ecosystem Services




Figure 13: Sankey diagram showing the flow of ecosystem services from natural assets in the catchment.

Some of the benefits that are not included in Figure 13 but are significant to the catchment include:

- Increased pollination and the control of pests and diseases, which promote healthy ecosystems and improve agricultural productivity;
- the cultural, historical and aesthetic value ecosystems provide to places; and
- the provision of all other ecosystem services through biodiversity and biosphere integrity.

These benefits are not assessed with the NCRAT, however a qualitative assessment of their benefit is provided below.

4.4.2 Provisioning Services

Food production

Agricultural practices, such as arable farming and livestock rearing, contribute significantly to national food security and support the local economy. The estimated annual yields for

arable and livestock (dairy and meat) are approximately 41,500 tonnes and 9,300 tonnes respectively.

Water supply

In 2018, it is estimated that over 10 billion m³ of water was abstracted from non-tidal surface waters and groundwater public supply, agriculture, industry and energy generation [63].

Overall, the volume of water abstracted from the catchment is approximated to be 830,000m³ per year.

Renewable energy

According to the Department for Energy Security and Net Zero, roughly 69 megawatts (MW) electricity was generated from onshore wind and 89MW from solar power in the East Lindsey District Council area in 2023 [64]. In comparison, in 2023 across England it was estimated that 3,094MW of electricity was generated from onshore wind, 43 MW from hydro generation and 13,976MW from photovoltaics. Since catchment-specific data is not available, the renewable energy generation value for the East Lindsey District Council renewable has been scaled to the catchment according to the relative size difference (the catchment is approximately 3.6% of the area of East Lindsey District Council). Using this method, it is estimated that 9,235MWh/year is generated in the catchment from renewable sources.

From desk-based research, notable renewable energy projects in the catchment include the following:

- The Hollies Solar Park (8.6MW) - a farm diversification project with solar PV and two wind turbines;
- Lincoln Farm Solar Park (11.4MW);
- Manor Farm Solar Park (in construction – with an expected capacity of 22MW); and
- White House Farm anaerobic digestion (AD) plant.

It is therefore considered that the 9,235MWh/year could be an underestimate of the overall contribution that renewable energy projects in the catchment provide to the wider East Lindsey District Council renewable energy provision.

Timber

Woodlands play a fundamental role in providing timber and wood fuel in the UK. Approximately half of the areas of woodland in the catchment have had a management plan or a felling license in the last 10 years or are owned by government organisations. While there are data limitations in relation to actual rates of timber extraction in woodlands within the catchment, national data can be applied to estimate a yearly volume of 261m³.

4.4.3 Regulating Services

Climate regulation

Data presented in the Climate Change Strategy for South Holland, Boston and East Lindsey District Councils suggests that land use, land use change and forestry (LULUCF) is the third largest source of greenhouse gas emissions in East Lindsey District Council at approximately 109,200tCO₂e/yr. For the catchment (which is 3.6% of East Lindsey District Council), the NCRAT presents net greenhouse gas fluxes as sequestering roughly

1,935tCO₂e/yr. This is calculated using the extent of different habitat types in the catchment and their average carbon sequestration rates.

The catchment is underlain by approximately 62ha of peatland, or 1% of the catchment area, which is in a 'drained' condition. The volume of carbon emitted by peatland in each of the above condition classes results in a central volume of net emission of 282tCO₂e/yr.

Enclosed farmland, saltmarsh and woodland in the catchment is calculated to sequester approximately 627tCO₂e/yr, 1166tCO₂e/yr and 405tCO₂e/yr, respectively.

It is important to note that the NCRAT assumes all agricultural land (i.e. arable and pastoral) is delivering net sequestration, which is likely incorrect (see limitations in section 3.1.7). At a national scale agriculture is a net emitter of greenhouse gases and it is likely this pattern is similar in the catchment; in 2021 approximately 48MtCO₂e were emitted from as a result of agricultural land management across the UK.

Air quality

The NCRAT considers the removal of four air pollutants, Particulate matter (PM_{2.5}), Sulphur Dioxide (SO₂), Nitrogen Dioxide (NO₂), and Ozone (O₃). Overall, approximately 340 tonnes of air pollutants are removed by vegetation each year. The removal of these air pollutants provides health benefits to the local population in the catchment. Given that the catchment is predominantly rural with a low population density, the health benefits of improved air quality are likely to be less significant than in more urban and densely populated areas.

It is important to note that air pollution is likely to be distributed unevenly across the catchment, however the NCRAT assumes an even distribution of air pollutant abatement across the whole catchment rather than modelling variable impact or value across the catchment. Therefore, both the impacts of poor air quality and the benefits of air pollutant removal may vary when measured on the ground.

Hazard regulation

The NCRAT only quantifies and values the cumulative yearly water storage capacity of woodlands in their canopy and soils. The annual volume of potential flood storage within areas of woodland in the catchment is approximately 9,990m³ of floodwater.

Disease and pests

The NCRAT does not quantify benefits relating to disease and pest control. It is estimated that globally 35–40% of potential crop yields are destroyed by pests [65]. Natural pest control, such as by bats, birds and insects, can enhance crop production by reducing herbivorous insects that damage or use parts of crops.

Pollination and seed dispersal

The NCRAT does not quantify benefits relating to pollination and seed dispersal. Pollination plays a crucial role in promoting healthy ecosystems and supporting the provision of other ecosystem services. Approximately 80% of British wildflower species and 37% of crops grown in the UK benefit from insect pollination, including beans and oilseed rape; additionally, a further 37% of crops grown in the UK, such as carrots and onions, require insect pollination to create viable seeds.

Within the catchment, the crop areas have been estimated based upon commercial agricultural holdings¹⁰ [46] as follows:

- Cereals – 2,550ha (approximately 52% of the catchment land use)
- Arable crops (excluding cereals) – 1,050ha
- Fruit and vegetables – 150ha

Many cereal and arable crops are wind-pollinated [66], whereas many fruit and vegetable crops (particularly oilseed rape, orchard fruit, soft fruit and field beans) make use of insects to transfer pollen. Without the pollinating insects these plants would likely have overall reduced quality and yields, resulting in a direct negative economic value impact. They also contribute to the diversity of wild plant species, habitats and wildlife in England, as well as its resilience and natural beauty. Pollination is therefore an essential ecosystem service which maintains biodiversity and supports other vital ecosystem functions, including soil protection, flood control and carbon sequestration.

The quantification of pollination benefits heavily relies on complex modelling, which is outside the scope of this assessment. Currently the direction and magnitude of changes in pollinator biodiversity, the value and functional relationship of pollinators to agriculture from farm to national scales and how this biodiversity and linked ecosystem service is only partly understood [67].

Soil retention and erosion

The NCRAT does not quantify benefits relating to soil erosion and retention. Soils provide many essential services including an environment for food production, filtering and absorbing water, reducing pollutants, regulating climate and providing habitat for organisms. It is estimated that over 2 million ha of soil are at risk of erosion in England and Wales and almost 4 million ha are at risk of soil compaction. This has implications for water resources, flooding and soil quality and food production [68].

4.4.4 Cultural Services

The NCRAT quantifies and values only four cultural ecosystem services: recreation, physical health, education, and volunteering. For this assessment, education and volunteering benefits were not possible to quantify due to lack of data availability.

Recreation

Access to open green spaces can provide a range of benefits to local economies. Based on data from the University of Exeter's ORVal Tool [33], approximately 171,800 adults (aged 16 or older) visit open green spaces in the catchment every year which supports the growth of local economies. A further 43,600 visits are made by children each year, but these are not included in the assessment of welfare value.

Physical health

Out of the 171,800 yearly visits to open green spaces within the catchment, roughly 88,500 are likely to partake in activities that are beneficial for their physical health.

¹⁰ Data based upon the most recent Local Authority data for East Lindsey District Council (2021) and apportioned to land within the Steeping catchment.

Education and volunteering

For this assessment, education and volunteering benefits were not possible to quantify due to lack of data availability. Although visits to the catchment for scientific and educational purposes are difficult to quantify, they do create benefits for human society as a whole. The protected areas and heritage sites within the catchment have the potential to enable insights about past and future landscape changes in the area.

Volunteering visits are similarly difficult to quantify, but can provide benefits to recipient organisations and their stakeholders, to wider society and to the volunteers themselves. Volunteering in nature and outdoor activities can have many positive benefits for the participants such as improving physical health, mental health and social connectedness. This not only provides a direct benefit to those participating in the activity but can have follow-on benefits by reducing national health care costs. Mental health improvements include reduced stress and anxiety and increased positive mood, self-esteem and resilience [69].

Aesthetic / sense of place

The aesthetic and sense of place benefits from nature are intrinsically difficult to measure, as it describes how people perceive, feel about, and interact with their environment. One means of measuring this benefit is through house prices, which tend to increase in areas where there is greater access to green and blue space due to improved local aesthetics.

4.4.5 Bundled services

Bundled services may not produce measurable benefits but are essential for ecological processes to function and facilitate the provision of all ecosystem services. Supporting services are typically not assessed on their own, as their importance underpins all individual ecosystem services discussed as part of this assessment.

Water quality

The bundled ecosystem service for water quality entails a range of benefits related to ecosystems functioning to remediate polluted water, the value of good-quality water for economic activities, recreation, education, and for healthy plant and animal populations.

The NCRAT applies non-market valuation approaches to quantify the value of good water quality within the catchment, mainly by applying data on people's willingness to pay for good water quality in combination with the ecological status of inland and coastal waterbodies and 2019 bathing water status. The catchment contains two WFD river waterbodies called Lymn / Steeping River and Cow Bank Drain, and one coastal waterbody called the Wash Outer - all three surface water bodies have a 'moderate' ecological classification status.

Biodiversity

The variety of living organisms in the catchment is important at a local level for increasing resilience to pests and diseases and supporting healthy ecosystems. At a global level, it is important for maintaining genetic diversity, which in turn is required for biosphere integrity, one of the nine planetary boundaries [70]. At both a local and a global level, Biodiversity enables the functioning of ecosystem services that are critical to all life of Earth.

While there are some ways to measure aspects or indicators of biodiversity (e.g. through Biodiversity Metrics, indices or indicator species), the NCRAT does not currently include this.

4.4.6 Beneficiaries Assessment

A qualitative Beneficiaries Assessment has been undertaken that explores who benefits from each ecosystem service, either directly or indirectly. The assessment considers who the beneficiaries of each ecosystem service are (i.e. the local population, wider society, public sector or private sector) and whether they are primary, secondary or indirect beneficiaries. The outputs of this are presented in Appendix A: Output - Benefits Statement.

The local population – including local communities and households (residents) – were identified as the primary beneficiary for most ecosystem services except provisioning services such as agriculture, water supply, timber and renewable energy. For these, as well as pollination, they have been identified as the secondary beneficiary.

The wider society – meaning those who live outside of the study area and include non-governmental organisations – has been identified as an indirect beneficiary for the majority of ecosystem services, specifically regulating and cultural services. For provisioning services and bundled services, the wider society is considered a secondary beneficiary. For climate regulation, the wider society is considered a primary beneficiary.

The public sector – referring to national and local government bodies – has been identified as an indirect beneficiary for the majority of ecosystem services, particularly provisioning services. The public sector is considered a secondary beneficiary of climate regulation, hazard regulation, disease and pest regulation, all bundled services and cultural services including recreation, health, volunteering and education. The public sector is considered the primary beneficiary of the mediation of wastes.

The private sector – including commercial enterprises, farming and industry – is considered a primary beneficiary for provisioning services such as agriculture, water supply, timber, and renewable energy as well as the regulation of disease and pests, pollination, soil condition and erosion, and mediation of wastes. The private sector is considered a secondary beneficiary of all bundled services, climate regulation and hazard regulation. For the remaining services, including cultural services and air quality regulation, the private sector is considered an indirect beneficiary.

4.5 Value

4.5.1 Natural capital value

The total natural capital value that has been estimated for the ecosystem services in the catchment is estimated to be £11.7 million per year, with a 100-year asset value of £360 million. Figure 14 shows how this figure can be broken down into the different benefits from ecosystem services. This value will be an underestimate of the real value of natural capital assets as some benefits have not been valued, including biodiversity which underpins the functioning of all other ecosystem services. More information on the value of ecosystem service benefits is provided in the subsequent sections. More information about the estimated values of ecosystem services is provided in Section 4.5.




Figure 14: Pie chart showing the natural capital value of different ecosystem services in the catchment (£ million).

Not all ecosystem services have been quantified into a monetary value and so show a value of £0 on Figure 14. Volunteering and education have not been included in the assessment because there is not enough data available to calculate a value. Since there is no marine habitat within the catchment, the provision of food from fish and shellfish landings has not been included as it is not appropriate to attribute these benefits to the study area. The NCRAT does not provide a monetary value for hazard regulation as the valuation methodology is not sufficiently robust. Other ecosystem services not included are those that are particularly hard to quantify, including biodiversity, pollination, regulation of disease and pests, regulation of soil quality and erosion, and cultural benefits relating to amenity, aesthetics and spirituality. Consequently, the total values of assets and services in a natural capital account will always be an underestimate.

4.5.2 Provisioning services

Food production

The estimated annual arable and livestock yields are over 41,500 tonnes and 9,300 tonnes respectively, which have a combined worth of over £5.9 million each year. The asset value over the next 100 years for agriculture is £176 million. As noted in section 3.1.7, the natural capital values do not include the potential costs from carbon emissions associated with agricultural activities (e.g. cost of carbon emissions from land management, fertiliser production, machinery etc.).

Water supply

Overall, the volume of water abstracted from the catchment is approximated to be 830,000m³ per year, equating to £4.2 million each year and £124 million over the next 100 years – these are the central values.

Renewable energy

The renewable energy generated for the catchment is estimated to be 9,235MWh per year, which has been valued at £130,000 a year and £3.8 million over the next 100 years.

Timber

The national value of timber and fuel extracted from woodlands is estimated at almost £15 billion. National data can be applied to the catchment to estimate a yearly timber production volume of 261m³, which is valued at approximately £8,740 per year. The 100-year asset value of this is £260,000.

A key consideration for this figure is that, in order to provide a constant benefit, woodlands need to be managed sustainably. As such, timber harvest rates should not negatively impact the health and therefore the wider ecosystem service provision within a woodland.

4.5.3 Regulating Services

Climate regulation

Based on the extent of habitat types in the catchment, the NCRAT presents the net greenhouse gas fluxes as sequestering roughly 1,935tCO₂e/yr for a central value, providing an annual value of £0.5 million - this has a 100-year asset value of over £30 million.

The NCRAT provides maximum and minimum values for climate regulation which should also be considered given the amber confidence interval for climate regulation (see Table 1 for confidence interval definitions) and uncertainty in the results. The minimum annual volume of carbon sequestered is 1,290tCO₂e/yr (£0.36 million per year) and the maximum is 2,590tCO₂e/yr (£0.73 million per year).

The estimated value for climate regulation is based on the extent of different habitat types in the catchment. The majority of the catchment is agricultural land, which has a relatively low carbon sequestration rate. The value for climate regulation would be higher if the area of woodland, wetlands or saltmarsh was increased, as these have a higher carbon sequestration rate. As explained in section 4.4.3, the peatland in the catchment is in a 'drained' condition and is a net emitter of carbon, which significantly reduces the value for climate regulation.

Air quality

The removal of 34 tonnes of air pollutants by vegetation each year leads to an estimated avoided cost of treatment and productivity loss of £60,000 per year and £1.8 million over the next 100 years.

Hazard regulation

In the catchment, an estimated 9,990m³ of floodwater is temporarily stored in areas of woodland. The NCRAT does not provide a monetary value for this benefit, however the cost of creating a flood storage reservoir to store a similar amount of water is estimated at £5,010 per year.

Disease and pests

The NCRAT does not provide a monetary value for the benefits associated with disease and pest control in the catchment. There is not sufficient information to calculate the precise economic impact of pests and disease. The spread of invasive and non-native species more broadly is estimated to cost the UK economy £1.8 billion each year [6].

Pollination

The NCRAT does not provide a monetary value for the benefits associated with pollination in the catchment. Pollination by insects and other animals has been estimated to be worth £500 million annually in the UK [6].

Soil erosion and retention.

The NCRAT does not provide a monetary value for the benefits associated with soil erosion and retention in the catchment. As peat is drained it shrinks; becoming lighter, friable and vulnerable to erosion. Production losses due to soil erosion are estimated to cost £40 million per year in England and Wales [6].

4.5.4 Cultural Services

Recreation

The value of approximately 171,800 visits by adults (aged 16 or older) to open green spaces in the catchment every year is worth approximately £693,000 each year and has a 100-year asset value of approximately £17 million. The welfare value of a further 43,600 visits made by children each year cannot be quantified but is likely to be significant.

Physical health

By applying a valuation approach to measuring the improvements in 'quality adjusted life years' (QUALYs), the NCRAT estimates that £164,630 in healthcare treatment costs can be avoided every year due to improved physical health as a result of 88,500 active visits to green spaces. The 100-year asset value of this is approximately £5 million.

Education and volunteering

Although education and volunteering have been scoped out of this assessment due to lack of data availability, they are valuable ecosystem services. The number of volunteer hours and educational visits can be used to calculate the value of education and volunteering in a catchment. It is likely that the protected areas and heritage sites within the catchment enable valuable volunteering and educational opportunities. All 141 National Nature Reserves managed by Natural England have a total volunteering value of £1.8 million (for 150,000 hours of work) and a total educational value of £123,000 [34] [69].

Aesthetic / sense of place

The NCRAT does not provide a monetary value for the benefits associated with aesthetics and sense of place. Recreation and aesthetic values are in part captured by house prices as there is value associated with proximity to green and blue spaces for recreation (free trips to nature) and the visual amenity (aesthetic) value provided by being able to view green and blue spaces. In 2021, the combined aesthetic and recreation value of these services was estimated to be worth £4 billion [71].

4.5.5 Bundled services

Water quality

Based on a willingness to pay approach, the river water quality is worth an estimated £0 per year as an alternative to having poorer water quality. This is because none of the surface waterbodies within the Fens East and West or Steeping and Eaus operational catchments meet a 'Good' or 'High' ecological status. The NCRAT only provides an annual value of good or high water quality per km.

The catchment contains two WFD watercourses and one WFD coastal waterbody. All have a 'moderate' ecological classification status.

Biodiversity

The NCRAT does not assess the value of biodiversity. However, the value of biodiversity to our economy and society should not be overlooked as it underpins the value of all other ecosystem services.

The Dasgupta Review provides a thorough assessment into the value of biodiversity to our economy and society [72]. The Report distinguishes six sources of biodiversity's value including human existence, human health, amenity, the provision of ecosystem services from Nature's goods and services, species existence, and the intrinsic value of nature itself [72].

4.5.6 Monetary value confidence intervals

Table 5 shows a RAG rating of confidence in the monetary values assigned to the annual and 100-year flow of benefits [34]. Definitions for confidence intervals are provided in section 3.1.4.

Table 5: The natural capital values calculated using the NCRAT colour coded according to the confidence interval.

Ecosystem Service	Annual value (central) £m	Asset value (PV; 100 yr) (low) £m	Asset value (PV; 100 yr) (high) £m	Asset value (PV; 100 yr) (central) £m
Agriculture - Arable	4.59	-	-	136.69
Agriculture - Livestock (dairy)	1.27	-	-	37.74
Agriculture - Livestock (meat)	0.08	-	-	2.46
Fish and shellfish landings	0.00	-	-	0.00
Water supply (public)	0.49	-	-	14.49
Water supply (energy generation)	-	-	-	-
Water supply (all other)	3.70	-	-	110.34
Timber	0.01	-	-	0.26
Renewable energy	0.13	-	-	3.81
Climate regulation	0.55	21.30	38.90	30.11
Air quality - PM2.5	0.01	-	-	0.33

Air quality - SO2	0.00	-	-	0.02
Air quality - NO2	0.03	-	-	0.24
Air quality - O3	0.02	-	-	1.22
Hazard regulation	-	-	-	-
Disease and pests				
Pollination				
Soil quality / erosion				
Noise mitigation				
Waste remediation				
Recreation (adults)	0.69	-	-	16.85
Recreation (children)	-	-	-	-
Physical Health	0.16	-	-	4.91
Education	0.00	0.00	0.00	0.00
Volunteering	0.00	-	-	0.00
Amenity				
Aesthetic / sense of place				
Spiritual				
Water quality - rivers	0.00	0.00	0.00	0.00
Water quality - transitional, coastal waters and lakes	0.00	0.00	0.00	0.00
Biodiversity				

5. Summary and next steps

5.1 Summary of the NCRAT results for the catchment

The flow of ecosystem services that are derived from natural assets within the catchment were estimated, which was used to calculate the value of the benefits those services provide. This process is summarised in Figure 15. The analysis is not exhaustive and does not include all assets, services and benefits, therefore the natural capital values shown are an underestimate. Figure 15 indicates the confidence in the valuation provided as High (3 coloured squares), Medium (2 coloured squares), or Low (1 coloured square).




Figure 15: Summary of the NCRAT results for the catchment

5.2 Strengths, weaknesses, opportunities and threats

5.2.1 Strengths

The key natural capital strengths in the catchment include:

- **Coastal priority habitat:** The catchment includes approximately 225ha of coastal saltmarsh providing important habitats sublittoral sandbanks, sandflats, mudflats, shallow inlets and reefs which attract a range of wildlife including wintering waterbirds; passage birds; waders; breeding birds; harbour seal; and shellfish [42] [43] [44]. These habitats provide crucial ecosystem services, including water quality improvement, carbon sequestration, biodiversity, recreation, tourism, and protection against erosion and flooding. Additionally, they serve as nurseries for many commercial fish species.

Coastal saltmarsh and intertidal mudflats are highly effective at capturing and sequestering carbon dioxide from the atmosphere; it is estimated that saltmarsh in the catchment sequesters 1,166 tonnes of CO₂ every year. This significant carbon storage capacity not only helps mitigate climate change by reducing greenhouse gas concentrations but also enhances the resilience of coastal ecosystems. By protecting and restoring these coastal habitats, the region can leverage their natural ability to sequester carbon, contributing to both local and global climate goals.

- **Statutory designations:** The coastal margins in the catchment all sit within a statutory designated site; namely Gibraltar Point (SPA, Ramsar, NNR and SSSI), The Wash (SPA, Ramsar and SSSI), or Saltfleet by Theddlethorpe Dune & Gibraltar Point (SAC, NNR and SSSI). Bratoft Meadows (SSSI) is the only designated site in the catchment located outside coastal margins. These designations provide legal protection to critical habitats - ensuring the conservation of valuable natural resources, supporting biodiversity, and protecting the continuous provision of ecosystem services.
- **Fertile soil:** The large majority (93%) of the catchment is enclosed farmland; the natural capital value of this land is estimated at £5.9 million annually through food production. Approximately 5,900ha of arable fields and agricultural grasslands support both arable farming and livestock rearing, producing over 42,000 tonnes of arable yields and 9,300 tonnes of livestock yields each year. These activities support food production, provide employment, and help maintain the landscape character of the Lincolnshire Coast and Marshes National Character Area.
- **Water environment:** The catchment contains important surface water and groundwater bodies that are crucial for human water supply and local ecosystem functioning. The main surface waterbodies are the Steeping River, Cowcroft Drain and River Lymn; the groundwater bodies underlying the catchment include the Steeping Long Eau Little Eau Chalk Unit, and the Steeping Great Eau Long Eau and Witham Spilsby Sandstone Unit. The natural capital value of water supply in the catchment is estimated at £4.3m annually.
- **Renewable energy:** The catchment contains two operational solar farms and two wind turbines, with an additional solar farm currently in construction. The renewable energy generated for the catchment is estimated to be 8,750MWh per year, which has been valued at £120,000 a year and £3.6 million over the next 100 years.

5.2.2 Weaknesses

The weaknesses of natural capital in the catchment are as follows:

- **Environmental degradation:** The vast majority of the catchment is made up of agricultural land, covering 93% of land area. Whilst this is important for food production, intensive agricultural practices can contribute to soil degradation, habitat fragmentation, and pollution. These activities can also lead to the loss of biodiversity, reduced soil fertility, and increased greenhouse gas emissions, impacting ecosystem health and resilience. Additionally, the use of chemical fertilizers and pesticides can contaminate water bodies and groundwater resources, further degrading the quality of natural habitats and affecting aquatic life. The catchment has a relatively low proportion of semi-natural and ancient woodlands, and the condition of peatland has been degraded through drainage. This is likely as a result of historic land use change driven by agricultural intensification.
- **Vulnerability to climate change:** The coastal habitats in the catchment are at risk from rising sea levels, increased frequency of extreme weather events, and changing precipitation patterns. These changes can lead to habitat loss, reduced biodiversity, and increased vulnerability to coastal erosion and flooding. In addition, it can impact food production through damage to crops and increased soil erosion during extreme weather events, and through unpredictable seasons. The impacts of climate change can also disrupt the balance of marine ecosystems, affecting fish populations and other marine life that are crucial for local fisheries and biodiversity.

In the catchment, agricultural land and ecologically valuable habitats - including designated sites - are at risk from retreating coastlines, saline intrusion, and coastal flooding. These threats can lead to the loss of valuable farmland, reduced agricultural productivity, and degradation of critical biodiversity areas, directly impacting both economy and ecology of the region, as well as indirectly through the erosion of ecosystem service provision due to habitat degradation.

5.2.3 Threats

Threats to the condition or quality of natural capital assets in the catchment have been identified, including:

- **Climate change:** The catchment is at risk of flooding, both from the sea given its low-lying coastal topography, as well as from the Steeping River which crosses the area. A significant proportion of the catchment area is considered to be at high risk of flooding from rivers or sea (Flood Zone 3) [73]. These changes can lead to habitat loss and reduced biodiversity (e.g. as a result of salination impacts), and increased vulnerability to coastal erosion and flooding.

Rising sea levels, increased frequency of extreme weather events, and changing precipitation patterns pose significant risks to food production and coastal habitats. The catchment has suffered from floods on multiple occasions e.g. On 12th June 2019 a series of breaches led to extensive flooding in the catchment, with over 100 houses and 550ha of farmland flooded (circa 10% of the farmland in the catchment). Again in 2023, Storm Babet caused widespread flooding in Lincolnshire with overtopping in Wainfleet.

Rising sea levels and an increased demand for groundwater abstraction increases the risk of saline intrusion to groundwater and soils. This can lead to a variety of issues such as habitat loss, reduced agricultural productivity and loss of aquifer water quality [74].

- **Agricultural practices:** Intensive farming practices, particularly on arable land, can lead to degradation of soil and peat; reduced soil fertility; and increased greenhouse gas emissions. The condition of any remaining peatland in the catchment is already poor due to decades of drainage and intensive agriculture.

The use of nitrogen and phosphorus-based fertilisers can also contribute to waterway pollution. The Environment Agency's reasons for not achieving Good (RNAG) dataset identifies diffuse pollution from agricultural and rural land management and physical modifications associated with flood defence structures, land drainage operations, and agricultural land use as the current RNAG's for the waterbodies within the catchment.

- **Water resource management:** Over-extraction of water from aquifers and rivers can lead to reduced water availability for ecosystems and human use, including irrigation for arable crops. This can exacerbate the impacts of droughts and reduce the resilience of freshwater habitats. Water is only available for abstraction from the Lower River Steeping and the Partney Bridge (River Lymn) on 138 days of the year. Notably, the total water available for abstraction per day across the entire catchment is just 12,300m³, sufficient to provide irrigation of c.25mm across approximately 50ha of land [75]. Consequently, irrigated crops can only be produced in this catchment reliably using water from the on-farm reservoirs which are in the area towards The Wash [65].

According to the Water Resources East, Lincolnshire is projected to have to water supply-demand deficit in the region of 15-40 megalitres per day by 2050 – demonstrating the importance of sustainable water management in the local area and the threat of severe water shortage in the future [47].

There are trade-offs between the benefits that are obtained from different ecosystem services, especially if natural assets are not managed sustainably. For example, abstraction of water for water supply to provide drinking water, grow crops and livestock may deplete water resource and have knock on negative implications for other ecosystems that rely on sufficient flows and volumes of water being available. This includes other provisioning services (such as timber) regulating services (such as climate regulation and soil quality), cultural services (such as recreation and amenity) and bundled services (such as water quality and biodiversity).

- **Pollution:** Air and water pollution from agriculture and urban areas can degrade natural habitats and reduce their ability to provide ecosystem services. This includes the accumulation of pollutants in vegetation and water bodies, impacting human health and biodiversity.
- **Invasive species:** The introduction and spread of invasive species can disrupt local ecosystems by outcompeting native species for resources, altering habitat structures, and introducing diseases. This can lead to a decline in native biodiversity and negatively affect ecosystem functions and services.
- **Urban development:** Expansion of urban areas and infrastructure can lead to habitat fragmentation, loss of green spaces, and increased pollution. This can negatively impact biodiversity and the provision of ecosystem services such as air quality regulation and recreational spaces.

5.2.4 Opportunities

There is opportunity to improve natural capital in the catchment through:

- **Enhancement of asset quality:** Investing in the restoration and enhancement of natural habitats can improve their condition and resilience. This includes reforestation, wetland restoration, peatland restoration (such as in the north western part of the catchment, south of Great Steeping), and the creation of green infrastructure in urban areas such as Wainfleet All Saints and Wainfleet Saint Mary.

The 'Site Improvement Plan Saltfleetby-Theddlethorpe Dunes & Gibraltar Point' indicates that investigation into interventions on hydrological; continuity may provide an opportunity for enhancements to this protected site.

Reducing pollution can also enhance asset quality, addressing sources of excess nutrients, plastics and other pollutants.

- **Habitat creation:** Creating new habitats and extension of existing habitats, such as artificial reefs, saltmarsh, grazing marsh, urban green spaces, and wildlife corridors, can enhance biodiversity and ecosystem service provision. Given the low topography and high flood risk of the catchment, creating wetland habitat and restoring lowland peatland will be especially important for the future of the catchment to manage flood risk and promote carbon sequestration. There is also an opportunity to manage coastal realignment in order to allow coastal habitats and the benefits they bring, to migrate landward.
- **Recreation and tourism:** Improving the recreational value of new or existing green space can significantly enhance community well-being, encourage outdoor activity, and promote environmental awareness. This has benefits for natural capital, such as improved human health and education. This can be achieved by improving accessibility to green space through well maintained and wheelchair-accessible paths, good public transport links and clear signage. Other opportunities to improve the recreational value of natural assets include improving amenities (e.g. benches, bins, lighting and toilet facilities); adding recreational facilities (e.g. playgrounds and sports areas); and by promoting nature engagement through educational signage and community gardens.
- **Use of NbS:** Implementing NbS, such as sustainable agriculture, green engineering, and ecosystem-based adaptation, can deliver multiple benefits. Natural flood management is a type of NbS encompassing a suite of measures which work with natural processes to reduce the risk of flooding. These processes protect, restore, and mimic the natural functions of catchments, floodplains and the coast to slow, spread and store water, and each intervention comes with its own set of multiple benefits. NBS solutions can enhance resilience to climate change, improve water quality, and support biodiversity while providing economic and social benefits.
- **Sustainable land and water management:** Adopting sustainable practices in agriculture and water management can reduce environmental impacts and enhance ecosystem services. This includes practices such as conservation agriculture, agroforestry and integrated water resource management. Planting hedgerows and grassland along the agricultural field margins in the south of the catchment provides an opportunity to improve biodiversity and connectivity, and will support other ecosystem services such as pollination, water quality, flood management and climate regulation.

- **Regenerative farming practices:** Implementing regenerative farming practices focuses on restoring soil health, increasing biodiversity, and improving water cycles. These practices can lead to more resilient and productive agricultural systems, enhancing ecosystem services and reducing environmental impacts, especially along key watercourses such as the Lymn/Steeping river and Cow Bank Drain. There is also opportunity to support appropriate management of salt marshes and grazing marsh, such as restoration of sustainable grazing.
- **Policy and governance:** Strengthening policies and governance frameworks to support natural capital conservation and sustainable development can create an enabling environment for positive change. This includes integrating natural capital into planning and decision-making processes and promoting stakeholder engagement and collaboration. One way of implementing this is through Local Nature Recovery Strategies (LNRS) or assessment and enhancement of natural capital through the local plan making process.

LNRS were introduced through the Environment Act 2021, as new systems of spatial strategies for nature in England. Each LNRS includes a local habitat map and a written statement of biodiversity priorities, outlining practical actions such as creating wetlands, restoring peatlands, planting trees, and managing existing woodlands more sustainably. Farmers and land managers will be able to use the LNRS to understand the potential of their land for strategic nature recovery actions, which in turn could be applicable for funding through Environmental Land Management Schemes and other funding mechanisms. LNRS should be used by plan-makers to inform the way they address the National Planning Policy Framework requirement for plans to protect and enhance biodiversity, which in turn will have positive outcomes for natural capital.

The Greater Lincolnshire LNRS, which includes the catchment, is in the early stages of project planning and organization, focusing on mapping existing habitats and identifying local priorities for nature recovery. It is due to be published in March 2026 following public consultation in Autumn 2025. Managed by Lincolnshire County Council in partnership with North Lincolnshire Council, North East Lincolnshire Council, and the Greater Lincolnshire Nature Partnership, the strategy includes projects like floodplain meadow restoration and aims for no net loss of priority habitats by 2025.

The Landscape Recovery Fund is a long-term, large-scale government project fund with a focus on addressing climate change and habitat creation. The Lincoln and Witham Valley Farming and Nature Network (LWFNN) is a Landscape Recovery Project in the Upper Witham Valley that aims to create 3,200ha of habitats. This includes river and riparian restoration, wetland habitat, mesic (damp) habitat, and dry scrub habitats. In addition, the project will work towards establishing a more regenerative food production system in the local area; reducing carbon emissions; increasing soil health and ecological connectivity; increasing access to green space; and new volunteering and learning opportunities in the community. Whilst this project is not within the study area, it provides a great example of the potential benefits of Landscape Recovery Projects in the area [76].

6. Bibliography

- [1] Natural Capital Coalition, “Natural Capital Protocol,” 2016.
- [2] Capitals Coalition, “Natural Capital Protocol,” Natural Capital Coalition, 2016. [Online]. Available: https://capitalscoalition.org/capitals-approach/natural-capital-protocol/?fwp_filter_tabs=guide_supplement. [Accessed 11 December 2024].
- [3] HM Treasury, “The Green Book,” 2022. [Online]. Available: https://assets.publishing.service.gov.uk/media/6645c709bd01f5ed32793cbc/Green_Book_2022__updated_links_.pdf. [Accessed 21 November 2024].
- [4] Department for Environment, Food and Rural Affairs, “Enabling a Natural Capital Approach guidance,” 19 July 2023. [Online]. Available: <https://www.gov.uk/government/publications/enabling-a-natural-capital-approach-enca-guidance/enabling-a-natural-capital-approach-guidance#introduction-to-natural-capital>. [Accessed 29 October 2024].
- [5] R. a. d. G. R. Leemans, “Millennium Ecosystem Assessment: Ecosystems and human well-being: a framework for assessment,” Island Press, Washington/Covelo/London, 2003.
- [6] J. Lusardi, P. Rice, J. Craven, C. Hinson, F. Bell, A. Morgan, K. Martin, M. Dobson, T. Sunderland and R. Waters, “State of Natural Capital Report for England 2024: Risks to nature and why it matters,” Natural England, 2024.
- [7] Environment Agency, “The Environment Agency’s Natural Capital Catchment Register and Account Tool and Scorecard,” Environment Agency, 2019.
- [8] R. Lenane, “The Natural Capital Register and Account Tool, Version 1.2 – Technical Report,” Environment Agency, Bristol, 2023.
- [9] Copernicus Land Monitoring Services, “Corine Land Cover,” 2018. [Online]. Available: <https://land.copernicus.eu/en/products/corine-land-cover/clc2018>. [Accessed 29 October 2024].
- [10] Natural England, “Provisional Agricultural Land Classification (ALC) (England),” 2019. [Online]. Available: <https://naturalengland-defra.opendata.arcgis.com/datasets/Defra::provisional-agricultural-land-classification-alc-england/about>. [Accessed 16 12 2024].
- [11] Environment Agency, “Statutory Main River Map,” 07 June 2024. [Online]. Available: <https://www.data.gov.uk/dataset/4ae8ba46-f9a4-47d0-8d93-0f93eb494540/statutory-main-river-map>. [Accessed 29 October 2024].
- [12] Ordnance Survey, “OS Open Rivers,” 2024. [Online]. Available: <https://www.ordnancesurvey.co.uk/products/os-open-rivers#get>. [Accessed 29 October 2024].

- [13] OpenStreetMap contributors, "OpenStreetMap database," OpenStreetMap Foundation, 2024. [Online]. Available: <https://www.geofabrik.de/data/download.html>. [Accessed 16 December 2024].
- [14] Forestry Commission, "National Forest Inventory England 2020," 30 June 2022. [Online]. Available: https://data-forestry.opendata.arcgis.com/datasets/30cf3ba3df444b6695cdafe250ed2172_0/about. [Accessed 29 October 2024].
- [15] Natural England, "England Peat Status GHG and C storage," 03 June 2024. [Online]. Available: <https://www.data.gov.uk/dataset/8ea9074e-bafc-4cc4-85dd-19cda1dfbfd5/england-peat-status-ghg-and-c-storage>. [Accessed 29 October 2024].
- [16] Natural England, "Ancient Woodland (England)," 03 June 2024. [Online]. Available: <https://www.data.gov.uk/dataset/9461f463-c363-4309-ae77-fdcd7e9df7d3/ancient-woodland-england>. [Accessed 29 October 2024].
- [17] Natural England, "Priority Habitats Inventory (England)," 25 September 2024. [Online]. Available: <https://www.data.gov.uk/dataset/4b6ddab7-6c0f-4407-946e-d6499f19fcde/priority-habitats-inventory-england>. [Accessed 29 October 2024].
- [18] Department for Transport, "Value for Money Framework," 2024. [Online]. Available: <https://assets.publishing.service.gov.uk/media/67459942b58081a2d9be968e/value-for-money-framework.pdf>. [Accessed 01 October 2024].
- [19] Department for Environment, Food and Rural Affairs, "Data Services Platform - WFD Groundwater Bodies Cycle 2 Classification 2019," 2020. [Online]. Available: <https://environment.data.gov.uk/dataset/6b5c81d9-1dc9-4635-b7bd-54d3d2488ccd>. [Accessed 29 October 2024].
- [20] Department for Environment, Food and Rural Affairs, "Data Services Platform - WFD Transitional and Coastal Waterbodies Cycle 2," 2013. [Online]. Available: <https://environment.data.gov.uk/dataset/78c2df61-d465-11e4-b839-f0def148f590>. [Accessed 29 October 2024].
- [21] Department for Environment, Food and Rural Affairs, "Data Services Platform - WFD Lake Waterbodies Cycle 2," 2014. [Online]. Available: <https://environment.data.gov.uk/dataset/76b02c00-d465-11e4-a772-f0def148f590>. [Accessed 29 October 2024].
- [22] Forestry Commission, "Felling Licence Applications England," 20 July 2021. [Online]. Available: https://data-forestry.opendata.arcgis.com/datasets/f0fe491790df46f8bb01d54e6875ef63_0/explore. [Accessed 29 October 2024].
- [23] Environment Agency, "Surface Water Availability for Water Resource Charging," 28 February 2021. [Online]. Available: <https://environment.data.gov.uk/dataset/3fb59850-ef87-4b0e-b429-1f42e9e101d9>. [Accessed 29 October 2024].
- [24] Environment Agency, "Groundwater Management Units coloured according to water resource availability colours," 12 April 2016. [Online]. Available: <https://environment.data.gov.uk/dataset/119e6ba7-b7fe-463a-961f-34232ef3e7cd>. [Accessed 29 October 2024].

- [25] Department for Environment, Food and Rural Affairs; and Environment Agency, "Swimfo: Find a bathing water," 2024. [Online]. Available: <https://environment.data.gov.uk/bwq/profiles/>. [Accessed 29 October 2024].
- [26] Natural England, "Sites of Special Scientific Interest (England)," 30 September 2024. [Online]. Available: <https://www.data.gov.uk/dataset/5b632bd7-9838-4ef2-9101-ea9384421b0d/sites-of-special-scientific-interest-england>. [Accessed 29 October 2024].
- [27] Natural England, "Special Protection Areas (England)," 03 July 2024. [Online]. Available: <https://www.data.gov.uk/dataset/174f4e23-acb6-4305-9365-1e33c8d0e455/special-protection-areas-england>. [Accessed 29 October 2024].
- [28] Natural England, "Special Areas of Conservation (England)," 03 July 2024. [Online]. Available: <https://www.data.gov.uk/dataset/a85e64d9-d0f1-4500-9080-b0e29b81fbc8/special-areas-of-conservation-england>. [Accessed 29 October 2024].
- [29] Natural England, "Ramsar (England)," 03 July 2024. [Online]. Available: <https://www.data.gov.uk/dataset/67b4ef48-d0b2-4b6f-b659-4efa33469889/ramsar-england>. [Accessed 29 October 2024].
- [30] Natural England, "National Nature Reserves (England)," 25 June 2024. [Online]. Available: <https://www.data.gov.uk/dataset/726484b0-d14e-44a3-9621-29e79fc47bfc/national-nature-reserves-england>. [Accessed 29 October 2024].
- [31] Natural England, "Local Nature Reserves (England)," 30 September 2024. [Online]. Available: <https://www.data.gov.uk/dataset/acdf4a9e-a115-41fb-bbe9-603c819aa7f7/local-nature-reserves-england>. [Accessed 29 October 2024].
- [32] Natural England, "Green Infrastructure Map," 2024. [Online]. Available: <https://designatedsites.naturalengland.org.uk/GreenInfrastructure/Map.aspx>. [Accessed 29 October 2024].
- [33] Land, Environment, Economics and Policy Institute at The University of Exeter, "Outdoor Recreation Valuation Tool (ORVal: Version 2.0)," No date. [Online]. Available: <https://www.leep.exeter.ac.uk/orval/>. [Accessed 07 October 2024].
- [34] Natural England, "Accounting for National Nature Reserves: A Natural Capital Account of the National Nature Reserves managed by Natural England (NERR078)," Natural England, 2019.
- [35] R. Lenane, "The Natural Capital Register and Account Tool version 1.2 – User Guide," Environment Agency, 2023.
- [36] International Union for Conservation of Nature and Natural Resources and Peatland Programme, "Peatland Code Field Protocol," 2022.
- [37] M. Christie, A. Hyde, R. Cooper, I. Fazey, P. Dennis, J. Warren, S. Colombo and N. Hanley, "Economic Valuation of the Benefits of Ecosystem Services delivered by the UK Biodiversity Action Plan (Defra Project SFFSD 0702)," 2011.
- [38] R. Gregg, J. Elias, I. Alonso, I. Crsoher, P. Muto and M. Morecraft, "Carbon Storage and Sequestration by Habitat 2021 (NERR094)," Natural England, 2021.
- [39] Natural England, "England Peat Map," Department for Environment, Food & Rural Affairs, 06 May 2025. [Online]. Available:

<https://environment.data.gov.uk/dataset/3c72669a-1331-4930-a12d-b4f8032aa5cd>.
[Accessed 06 June 2025].

- [40] J. Nix, "Pocketbook for Farm Management 52nd edition," [Online]. Available: <https://theandersonscentre.co.uk/shop/john-nix-pocketbook/>. [Accessed December 2024].
- [41] Office for National Statistics, "UK natural capital accounts methodology guide: 2024," 2024. [Online]. Available: <https://www.ons.gov.uk/economy/environmentalaccounts/methodologies/uknaturalcapitalaccountsmethodologyguide2024/pdf>. [Accessed 14 March 2025].
- [42] Joint Nature Conservation Committee, "The Wash and North Norfolk Coast," [Online]. Available: <https://sac.jncc.gov.uk/site/UK0017075>. [Accessed 29 October 2024].
- [43] Joint Nature Conservation Committee, "Greater Wash SPA," December 2020. [Online]. Available: <https://jncc.gov.uk/our-work/greater-wash-spa/>. [Accessed 29 October 2024].
- [44] Ramsar Sites Information Service, "The Wash," 01 January 2007. [Online]. Available: <https://rsis.ramsar.org/ris/395>. [Accessed 29 October 2024].
- [45] Department for Environment Food & Rural Affairs, "MAGIC Map," [Online]. Available: <https://magic.defra.gov.uk/MagicMap.aspx>. [Accessed 29 October 2024].
- [46] Collisons and Associates, "Key land use, crop types and livestock populations on commercial holdings by 2021," Arup, 2024.
- [47] Water Resources East, "Draft Regional Water Resources Management Plan for eastern," Water Resources East, 2022.
- [48] International Union for the Conservation of Nature, "Peatland Programme - about peatlands," [Online]. Available: <https://www.iucn-uk-peatlandprogramme.org/about-peatlands>. [Accessed 03 01 2025].
- [49] F. Worrall, P. Chapman, J. Holden, C. Evans, R. Artz, P. Smith and R. Grayson, "Peatlands and Climate Change," 2010.
- [50] I. a. K. Holman, "A revised estimate of peat reserves and loss in the East Anglian Fens," Fens for the Future, 2011. [Online]. Available: <https://www.fensforthefuture.org.uk/admin/resources/banners/cranfieldfenland-phase-2-peat-assessment-19-01-11-final.pdf>. [Accessed 22 November 2024].
- [51] Fens for the Future Partnership, "A Strategic Plan for Fenland : A Proposal for an Enhanced Ecological Network (Final Report)," Mere Oak Ecology, Shrewsbury, 2012.
- [52] Office for National Statistics, "UK natural capital: peatlands," Office for National Statistics, 2019.
- [53] UK Government, "Gibraltar Point SPA," No date.
- [54] Ramsar Sites Information Service, "Gibraltar Point," 01 January 1999. [Online]. Available: <https://rsis.ramsar.org/ris/589>. [Accessed 11 December 2024].
- [55] Ove Arup & Partners Limited, "Draft Steeping catchment biodiversity baseline Technical Note," 2024.

- [56] Joint Nature Conservation Committee, "Saltfleetby-Theddlethorpe Dunes and Gibraltar Point," No date. [Online]. Available: <https://sac.jncc.gov.uk/site/UK0030270>. [Accessed 11 December 2024].
- [57] Environment Agency, "Steeping, Great Eau & Long Eau Abstraction Licensing Strategy, A strategy to manage water resources sustainably," March 2020. [Online]. Available: <https://assets.publishing.service.gov.uk/media/5e74e07886650c296f6eda49/The-Steeping-Great-Eau-and-Long-Eau-abstraction-management-strategy.pdf>. [Accessed 05 October 2024].
- [58] M. Zari, "Utilizing relationships between ecosystem services, built environments, and building materials," *Materials for a Healthy, Ecological and Sustainable Built Environment*, pp. 3-27, 2017.
- [59] C. Earlie, G. Messelink and P. Russell, "The role of beach morphology on coastal cliff erosion under extreme waves," *EARTH SURFACE PROCESSES AND LANDFORMS*, vol. 43, pp. 1213-1228, 2018.
- [60] K. Vastila, P. Rowinski, J. Aberle, J. Jarvela and M. Kalinowska, "How vegetation can aid in coping with river management challenges: A brief review," *Ecohydrology & Hydrobiology*, vol. 18, no. 4, pp. 345-354, 2018.
- [61] Convention on Biological Diversity, "Sustaining Life on Earth," [Online]. Available: <https://www.cbd.int/convention/guide>. [Accessed 29 October 2024].
- [62] National Farmers Union, "Delivering for Britain," 2019. [Online]. Available: <https://www.nfuonline.com/media/uvhhtjio/delivering-for-britain-food-and-farming-in-the-fens.pdf>. [Accessed 23 January 2025].
- [63] Department for Environment, Food & Rural Affairs, "Water abstraction statistics: England, 2000 to 2018," 4 January 2023. [Online]. Available: <https://www.gov.uk/government/statistics/water-abstraction-estimates/water-abstraction-statistics-england-2000-to-2018>. [Accessed 25 February 2025].
- [64] Department for Energy Security & Net Zero (DESNZ), "Renewable electricity by local authority 2014-2022," 2022. [Online]. Available: https://view.officeapps.live.com/op/view.aspx?src=https%3A%2F%2Fassets.publishing.service.gov.uk%2Fmedia%2F6512b64ff6746b000da4b9f7%2FRenewable_electricity_by_local_authority_2014_2022.xlsx&wdOrigin=BROWSELINK. [Accessed 29 10 2024].
- [65] A. Clasen, M. Peters, S. Ferger, M. Helbig-Bonitz, J. Shmack, G. Maassen, M. Schleunig, E. Kalko, K. Bohning-Gaese and I. Steffan-Dewenter, "Complementary ecosystem services provided by pest predators and pollinators increase quantity and quality of coffee yields," *Proceedings of the Royal Society B*, vol. 281, no. 20133148, 2014.
- [66] Parliamentary office of Science and Technology, "Insect pollination POSTNOTE," 2010.
- [67] V. e. a. 2014, "Status and value of pollinators and pollination services," Defra, 2014.
- [68] Environment Agency, "The state of the environment: soil," Environment Agency, 2019.

- [69] T. Sunderland, "Accounting for National Nature Reserves: A natural capital account of the National Nature Reserves managed by Natural England. Natural England Research Report, Number 078," 2019.
- [70] K. R. e. al., "Earth beyond six of nine planetary boundaries," *Science Advances*, vol. 9, no. 37, 2023.
- [71] Office for National Statistics, "UK natural capital accounts: 2023," 2023. [Online]. Available: <https://www.ons.gov.uk/economy/environmentalaccounts/bulletins/uknaturalcapitalaccounts/2023>. [Accessed 29 10 2024].
- [72] P. Dasgupta, "The Economics of Biodiversity: The Dasgupta," HM Treasury, London, 2021.
- [73] Environment Agency, "Flood risk information for planning," 2024. [Online]. [Accessed 23 12 2024].
- [74] Environment Agency Engage, "Flood and Coastal Resilience Innovation Programme - Project Groundwater (Greater Lincolnshire)," 03 February 2025. [Online]. Available: <https://engageenvironmentagency.uk.engagementhq.com/lin011-groundwater>. [Accessed 08 August 2025].
- [75] F. 2. partnership, "Steeping River Appendix: Environmental and Agricultural baseline," Arup, 2025.
- [76] VisitLincolnshire, "Lincoln & Witham Landscape Recovery," 2024. [Online]. Available: <https://www.visitlincolnshire.com/lincoln-witham-landscape-recovery/>. [Accessed 28 April 2025].
- [77] Fens for the Future, "Conservation and Restoration," No date. [Online]. Available: <https://www.fensforthefuture.org.uk/creating-the-future/conservation-and-restoration>. [Accessed 28 April 2025].

Appendix A

Output - Benefits Statement

Natural Capital Benefits Statement

Data sensitivity: OFFICIAL
Place name: Steeping

Summary of physical and monetary flows

The Natural Capital Benefit Statement reports the provision (flow) of a range of benefits. It draws on the currently available evidence to quantify the flow of benefits in physical and monetary terms. The quantitative results are accompanied by the Significance Assessment that provides a broader qualitative account of benefits, including those not currently assessed in quantitative terms. Each row represents a discreet combination of physical measurement and monetary valuation methods, and therefore this sheet shows the minimum number of unique rows required to cover all the benefits analysed.

Benefits are categorised according to a modified ecosystem service classification that incorporates some abiotic elements: (i) provisioning services (the production of physical goods we consume such as food and water); (ii) regulating services (ecosystems processes related to climate, air quality, hazard regulation, etc.); (iii) cultural services (aspects of the natural environment that enrich our lives, such as recreational benefits); and (iv) 'bundled' services (that combine multiple benefits). The contribution of supporting services - the ecological processes and functions such as nutrient cycling that maintain the conditions for life on earth and underpin ecosystem service provision - to current benefits flows is embedded within the monetary benefits for provisioning, regulating, cultural and bundled services.

Instruction: The Benefits Statement is automatically populated from the input and process tabs.

	Ecosystem service	Significance	Benefit valued	Benefit type	Beneficiaries			Physical flows		Monetary values						
					Local population	Wider Society	Public Sector	Private Sector	Measure (unit)	Avg unit / year	Valuation basis	Annual value (central) £m	Asset value (PV; 100 yr) (low) £m	Asset value (PV; 100 yr) (high) £m	Asset value (PV; 100 yr) (central) £m	Representation
Provisioning	Agriculture - Arable	+++	Food	Market	●●●●●	●●●●●	●●●●●	●●●●●	Yield of arable production (tonnes/yr)	41,430	Gross margin	4.59	-	-	136.69	■
	Agriculture - Livestock (dairy)	++	Food	Market	●●●●●	●●●●●	●●●●●	●●●●●	Yield of livestock (dairy) production (kL/yr)	9,006	Gross margin	1.27	-	-	37.74	■
	Agriculture - Livestock (meat)	++	Food	Market	●●●●●	●●●●●	●●●●●	●●●●●	Yield of livestock (meat) production (tonnes/yr)	290	Gross margin	0.08	-	-	2.46	■
	Fish and shellfish landings	--	Food	Market	○●○●○	○●○●○	○●○●○	○●○●○	Volume of fish and shellfish landings (tonnes/yr)	0	Net profit	0.00	-	-	0.00	■
	Water supply (public)	+++	Water supply	Market	●●●●●	●●●●●	●●●●●	●●●●●	Abstracted raw water quantity (m3/yr)	406,166	Resource rent	0.49	-	-	14.49	■
	Water supply (energy generation)	--	Water supply	Market	○●○●○	○●○●○	○●○●○	○●○●○	Abstracted raw water quantity (m3/yr)	0	Not valued	-	-	-	-	■
	Water supply (all other)	+++	Water supply	Market	●●●●●	●●●●●	●●●●●	●●●●●	Abstracted raw water quantity (m3/yr)	425,673	Marginal value	3.70	-	-	110.34	■
	Timber	++	Timber	Market	○●○●○	○●○●○	○●○●○	○●○●○	Volume of timber removals (m3/yr)	261	Stumpage price	0.01	-	-	0.26	■
	Renewable energy	+++	Renewable energy	Market	●●●●●	●●●●●	●●●●●	●●●●●	Renewable energy generation (MWh/yr)	9,235	Resource rent	0.13	-	-	3.81	■
	Climate regulation	+++	Climate	Non-market	●●●●●	●●●●●	●●●●●	●●●●●	Net CO2eq sequestered (tonnes/yr)*	1,936	Abatement cost	0.55	21.30	38.90	30.11	■
Regulating	Air quality - PM2.5	++	Health	Non-market	●●●●●	●●●●●	●●●●●	●●●●●	PM2.5 removed (tonnes/yr)	0	Avoided cost (treatment and productivity) plus welfare value	0.01	-	-	0.33	■
	Air quality - SO2	++	Health	Non-market	●●●●●	●●●●●	●●●●●	●●●●●	SO2 removed (tonnes/yr)	10	Avoided cost (treatment and productivity) plus welfare value	0.00	-	-	0.02	■
	Air quality - NO2	++	Health	Non-market	●●●●●	●●●●●	●●●●●	●●●●●	NO2 removed (tonnes/yr)	8	Avoided cost (treatment and productivity) plus welfare value	0.03	-	-	0.24	■
	Air quality - O3	++	Health	Non-market	●●●●●	●●●●●	●●●●●	●●●●●	O3 removed (tonnes/yr)	323	Avoided cost (treatment and productivity) plus welfare value	0.02	-	-	1.22	■
	Hazard regulation	+++	Flood risk reduction	Non-market	●●●●●	●●●●●	●●●●●	●●●●●	Annual volume of potential flood storage by woodlands (m3/yr)	9,991	Not valued	-	-	-	-	■
	Disease and pests	++			●●●●●	●●●●●	●●●●●	●●●●●								■
	Pollination	++			●●●●●	●●●●●	●●●●●	●●●●●								■
	Soil quality / erosion	+++			●●●●●	●●●●●	●●●●●	●●●●●								■
	Noise mitigation	+			●●●●●	●●●●●	●●●●●	●●●●●								■
	Waste remediation	+			●●●●●	●●●●●	●●●●●	●●●●●								■
Cultural	Recreation (adults)	++	Recreation	Non-market	●●●●●	●●●●●	●●●●●	●●●●●	No. visits to open spaces (visits/yr)	171,791	Welfare value	0.69	-	-	16.85	■
	Recreation (children)	++	Recreation	Non-market	●●●●●	●●●●●	●●●●●	●●●●●	No. visits to open spaces (visits/yr)	43,574	Not valued	-	-	-	-	■
	Physical Health	++	Health	Non-market	●●●●●	●●●●●	●●●●●	●●●●●	No. active visits to open spaces (visits/yr)	88,472	Avoided treatment cost	0.16	-	-	4.91	■
	Education	++	Educational benefits	Non-market	●●●●●	●●●●●	●●●●●	●●●●●	No. educational visits (visits/yr)	0	Exchange value	0.00	0.00	0.00	0.00	■
	Volunteering	+	Volunteering	Non-market	●●●●●	●●●●●	●●●●●	●●●●●	No. volunteering days (days/yr)	0	Opportunity cost	0.00	-	-	0.00	■
	Amenity	++			●●●●●	●●●●●	●●●●●	●●●●●								■
	Aesthetic / sense of place	++			●●●●●	●●●●●	●●●●●	●●●●●								■
Bundled	Spiritual	+			●●●●●	●●●●●	●●●●●	●●●●●								■
	Water quality - rivers	++	Various	Non-market	●●●●●	●●●●●	●●●●●	●●●●●			Welfare value of good water quality	0.00	0.00	0.00	0.00	■
	Water quality - transitional, coastal waters and lakes	+	Various	Non-market	●●●●●	●●●●●	●●●●●	●●●●●			Welfare value of good water quality	0.00	0.00	0.00	0.00	■
	Biodiversity	+++			●●●●●	●●●●●	●●●●●	●●●●●								■
Total market benefit value (£m)												10.26			305.80	
Total non-market benefit value (£m)												1.46			53.67	
Total quantifiable value (£m)												11.72			359.48	

Notes:

* Net carbon dioxide equivalent sequestration is estimated taking into account both carbon sequestration by habitats and green house gas sequestration and emissions from peatland

- Annual value is estimated flow at base year (Year 0)

- Asset value calculated as discounted flow of benefits (present value terms; 100 years). Future flows are discounted in accordance with HM Treasury Green Book guidance. This is an estimate of the gross asset value since it does include (net) costs of maintaining asset condition.




- Where available low - high ranges are reported for (gross) asset value, otherwise the cell has been left blank (-)

Significance key

Highly significant (+ + +)	The benefit flow is (very) important to the place in <i>relative terms</i> , because it impacts the wellbeing of a significant proportion (and possibly a wide variety) of people and businesses and/or due to wider social and cultural aspects.
Significant (+ +)	The benefit flow is of material importance in <i>relative terms</i> , because it impacts the wellbeing of a sizeable proportion of people and businesses.
Minor (+)	The benefit flow is of minor importance in <i>relative terms</i> , because it does not impact most people and business. Note: it may, though, still be highly important to a small number of specific individuals or groups.
Not significant (- -)	There is very little or no provision of this benefit and it impacts only a very small number of people and businesses.
Not assessed (-)	Significance rating not completed.

See "Input - Significance Assessment" tab for more details

Beneficiaries key




	Direct (primary and secondary) beneficiary from the ecosystem service provision.
	Indirect beneficiary from the ecosystem service provision.
	Beneficiary does not benefit from the ecosystem service provision.
Not assessed (-)	Beneficiaries rating not completed.

See "Input - Beneficiaries" tab for more details

Monetary valuation confidence key

Green	High confidence in results. Input data and assumptions are based on statistical reports, peer reviewed values or industry standard methodologies
Amber	Moderate confidence in results. Input data and/or assumptions from single source/not peer reviewed, or based on sources that are not specifically tailored to this context. We have used some assumptions or estimation and some of these may be open to question. Accuracy is better than +/- 50%
Red	Low confidence in results. Input data and/or parameters from single source and low level of transferability from original context. We are confident that the number is in the right order of magnitude. Order of magnitude implies that for an estimate of 5 that we are confident that the real figure is within the range 0.5 to 50
No value (-)	Not valued or it is not possible to provide a result that can be judged to be in the right order of magnitude. This is due to unquantifiable uncertainty in the science, valuation or the relationship between them. Currently limited to qualitative assessment only.

Representation key

	The benefit flow is substantially represented in the monetary valuation process, with only minor elements not included.
	The benefit flow is partially represented in the monetary valuation process.
	The benefit flow is not represented in the monetary valuation process.
Not assessed (-)	Representation rating not completed.

See "Input - Significance Assessment" tab for more details