



Public health recommendations for asymptomatic contacts of high consequence infectious disease (HCID) mammarenaviruses in UK settings

This document should be used for exposures to high consequence infectious disease (HCID) mammarenaviruses that have occurred in the UK.

This includes:

- Lassa virus
- Lujo virus
- Argentine haemorrhagic fever (AHF) caused by Junin virus
- Bolivian haemorrhagic fever (BHF) caused by Machupo virus
- Chapare virus (CHAPV)

In this guidance, 'case' refers to a confirmed symptomatic case, unless otherwise specified. 'Body fluids' can include blood, vomit, faeces, urine, saliva and semen.

For PPE requirements for the management of HCID mammarenavirus cases, please see the [ACDP guidance on the management of patients with viral haemorrhagic fever](#).

Any individual who has not had contact with the case within the last 21 days is not deemed to be a contact. For those individuals identified as potential contacts on a plane, please see the European Centre for Disease Prevention and Control (ECDC) [RAGIDA guidance](#) for contact classification.

The UKHSA Imported Fever Service or UKHSA Emerging Infections and Zoonoses team can provide assistance with contact categorisation on request.

Matrix: exposures in the UK

Category	Example scenarios	Actions
<p>Category 5 (very high risk of exposure)</p> <p>Description Unprotected high risk or protracted contact with a case or infectious material</p>	<p>Household or overnight contact of a case while they were symptomatic.</p> <p>Very significant exposure to material from a case such as:</p> <ul style="list-style-type: none"> • needlestick injury • confirmed history of splashes on mucous membranes or broken skin 	<ul style="list-style-type: none"> • Inform about risk (very high risk of exposure) • Give information sheet for Category 5 contacts <p>For 21 days following last exposure:</p> <ul style="list-style-type: none"> • self-isolate in private accommodation only, unless an alternative has been agreed with UKHSA • self-monitor and record temperature and symptoms twice daily (temperature to be taken before taking any anti-fever medication such as aspirin, paracetamol or ibuprofen) and report to UKHSA designated contact. If there is inability to take temperature, alternative arrangements for consultation with a clinician will be made with the relevant health services • if feeling ill in between reporting to the designated contact, take temperature, and if 37.5°C or higher, or have other symptoms suggestive of an HCID mammarenavirus (as outlined in the relevant contact information sheet) phone designated contact immediately • if any delay in making contact with the UKHSA designated contact phone NHS 111, and state that they have been exposed to an HCID mammarenavirus, giving details of which one. If seriously ill, dial 999, again reporting contact with an HCID mammarenavirus, giving details of which one <p>Additionally, for 21 days following the last exposure:</p> <ul style="list-style-type: none"> • avoid contact with other household members • do not share towels, bed linen, toothbrushes, razors or eating and drinking utensils such as cutlery or cups • avoid sexual contact <p>For category 5 contacts of Lassa virus and AHF caused by Junin virus, post-exposure prophylaxis can be considered and should be discussed on a case-by-case basis with the Imported Fever Service.</p>

Category	Example scenarios	Actions
<p>Category 4 (high risk of exposure)</p> <p>Description</p> <p>Unprotected close (within 2 metres) or direct contact with a case or infectious material</p>	<p>Direct contact (excluding direct contact with mucous membranes or broken skin) with a case or their body fluids (without appropriate PPE, or where there has been a breach).</p> <p>Unprotected sexual contact with a recovered case of an HCID mammarenavirus within 3 months of their illness.</p> <p>Direct exposure of skin or mucous membranes to an environment contaminated with blood or body fluids of a case without appropriate PPE, including on clothing or bedding (for example in a room after the case has left but before decontamination).</p> <p>Unprotected handling of clinical or laboratory specimens from a case, for example through transporting samples or work in a laboratory without appropriate PPE or laboratory processes.</p> <p>Passengers sat directly next to a case on a plane, and any crew, passengers or cleaning staff known to have had direct contact with the case, the case's body fluids, or items contaminated by their body fluids.</p>	<ul style="list-style-type: none"> • Inform about risk (high risk of exposure) • Give information sheet for Category 4 contacts <p>For 21 days following last exposure:</p> <ul style="list-style-type: none"> • self-monitor and record temperature and symptoms twice daily (temperature to be taken before taking any anti-fever medication such as aspirin, paracetamol or ibuprofen) and report to UKHSA designated contact. If there is inability to take temperature, alternative arrangements for consultation with a clinician will be made with the relevant health services • if feeling ill in between reporting to the designated person, take temperature, and if 37.5°C or higher, or there are other symptoms suggestive of an HCID mammarenavirus (as outlined in the relevant contact information sheet) phone designated contact immediately • if any delay in making contact with the UKHSA designated contact, phone NHS 111, and state that they have been exposed an HCID mammarenavirus, giving details of which one. If seriously ill, dial 999, again reporting possible contact with an HCID mammarenavirus, giving details of which one. <p>If the contact does not have any symptoms, then they can carry out normal activities during the 21 days after the last exposure, except for:</p> <ul style="list-style-type: none"> • if a healthcare worker, no patient contact for 21 days after last exposure • avoid contact with immunosuppressed people [note 1], pregnant women, and children aged under 18 years of age for 21 days from last exposure. <ul style="list-style-type: none"> ○ Following a risk assessment redeploy or exclude from work for 21 days if work involves direct contact with individuals who are immunosuppressed, pregnant or children aged under 18 years of age. <p>If they do not have any symptoms, the following apply focusing on reducing contact with others where-ever reasonably possible:</p> <ul style="list-style-type: none"> • advise all contacts to speak to line manager about options to reduce amount of contact with others at work, for example, remote working • any length of trip by private car within the UK is permitted • travel on local commuter or public transport is permitted, principally for work. Such trips should be as short as possible and avoided if possible • no travel internationally or via air within the UK for 21 days after exposure • non-essential medical or dental treatment should be postponed. For essential treatment, the healthcare provider must be informed prior to the procedure. • do not share towels, bed linen, toothbrushes, razors or utensils • avoid sexual contact • avoid situations where self-isolation would be impossible in the event of becoming ill (such as crowded public places)

Category	Example scenarios	Actions
Category 3 (medium risk of exposure) Description Unprotected close contact (in close proximity with a case but excluding direct contact)	<p>Someone who has been in the same room as and within 2 metres of the symptomatic case, without direct contact with the case, the case's body fluids, or items contaminated by their body fluids.</p> <p>On a plane:</p> <ul style="list-style-type: none"> passengers seated one seat away from the case in any direction (including diagonally) passengers in the 3 nearest seats across the aisle if the case was seated in an aisle seat (the passenger directly across the aisle, the passenger across the aisle one row in front and the passenger across the aisle one row behind) cabin crew who served the section of the aircraft where the case was seated, unless it is known that they had direct contact with the case 	<ul style="list-style-type: none"> Reassure (medium risk of exposure) Give information sheet for category 3 contacts <p>For 21 days following last exposure:</p> <ul style="list-style-type: none"> self-monitor temperature twice daily (temperature to be taken before taking any anti-fever medication such as aspirin, paracetamol or ibuprofen) if temperature is 37.5°C or higher, or if there are other symptoms suggestive of an HCID mammarenavirus (as outlined in the relevant contact information sheet), report to designated contact. If seriously ill, dial 999, reporting possible contact with an HCID mammarenavirus, giving details of which one. <p>If the contact does not have any symptoms, then they can carry out normal activities during the 21 days after the last exposure, except for:</p> <ul style="list-style-type: none"> non-essential medical or dental treatment should be postponed. For essential treatment, the healthcare provider must be informed of the exposure history prior to the procedure do not share towels, bed linen, toothbrushes or razors use barrier contraception or avoid unprotected sexual contact <p>There are no restrictions on work, travel or social activities as long as the contact has no symptoms. However, healthcare workers should inform their line manager that they have been identified as a contact of a mammarenavirus but are able to work with no restrictions as long as they are asymptomatic.</p>
Category 2 (low risk of exposure) Description Protected direct or close (within 2 metres) contact with a case or infectious material outside of a controlled environment	<p>Direct or close (within 2 metres) contact with a case or their body fluids outside of a controlled environment (for example a Trexler, designated HCID unit), but wore appropriate PPE correctly with no known breaches.</p> <p>Exposure to an environment contaminated with blood or body fluids of a case while wearing appropriate PPE correctly with no known breaches (for example in a room after the case has left but before decontamination).</p> <p>Laboratory staff in non-containment level 4 facilities handling specimens from a case who have had no breach of laboratory practice/PPE.</p>	<ul style="list-style-type: none"> Reassure (low risk of exposure) Give information sheet for Category 2 contacts <p>For 21 days following last exposure:</p> <ul style="list-style-type: none"> self-monitor temperature twice daily (temperature to be taken before taking any anti-fever medication such as aspirin, paracetamol or ibuprofen) if temperature is 37.5°C or higher, or if there are other symptoms suggestive of an HCID mammarenavirus (as outlined in the relevant contact information sheet), report to designated contact. If seriously ill, dial 999, reporting possible contact with an HCID mammarenavirus, giving details of which one. <p>If the contact does not have any symptoms, then they can carry out normal activities during the 21 days after the last exposure, except for:</p> <ul style="list-style-type: none"> non-essential medical or dental treatment should be postponed. For essential treatment, the healthcare provider must be informed of the exposure history prior to the procedure do not share towels, bed linen, toothbrushes or razors use barrier contraception or avoid unprotected sexual contact <p>There are no restrictions on work, travel or social activities as long as the contact has no symptoms.</p>

Category	Example scenarios	Actions
Category 1 (very low risk of exposure) Description Protected contact with a case or infectious material within a controlled environment Or: Unprotected household contact with a case in the 21 days prior to symptom onset	Household contact of a case while the case was asymptomatic in the 21 days before symptoms developed, but not after they became symptomatic. Clinical staff caring for a case in a controlled environment (for example, Trexler, designated HCID Unit) wearing appropriate PPE with no known breaches. Laboratory staff in a containment level 4 diagnostic laboratory or within a controlled environment handling specimens from a case who have had no breach of laboratory practice/PPE	<ul style="list-style-type: none"> • Reassure (very low risk of exposure) • Give information sheet for Category 1 contacts • Normal activities (no restrictions) Key clinical advice for contact: <ul style="list-style-type: none"> • if feeling ill (within 21 days of last exposure), take temperature (before taking any anti-fever medication such as aspirin, paracetamol, ibuprofen) • if temperature is 37.5°C or higher, or if there are other symptoms suggestive of an HCID mammarenavirus (as outlined in the relevant contact information sheet), call NHS 111 or, if very unwell, 999, and state recent contact with an HCID mammarenavirus, giving details of which one

Note 1: Immunosuppressed patients, as per the [Green Book definition](#), includes those with primary or acquired immunodeficiency, or individuals on immunosuppressive therapy, and includes those with: solid organ cancer; haematological disease and/or stem cell transplant; Child's-Pugh class B or C liver cirrhosis; stage 4 or 5 chronic kidney disease; immune mediated inflammatory disorders (including neurological and rheumatological conditions) treated with B-cell depleting therapy within 12 months; uncontrolled HIV; solid organ transplant recipients.

Produced by UKHSA's EIZ, RIPL and Guidance teams March 2026.