




Common cuttlefish

(*Sepia officinalis*)

Identifying features

Tentacular club	• Five to six rows of suckers that vary in size. • Five to six suckers in the middle are two times bigger than the others.
Body	• Broadly oval. • Up to 450mm mantle length.
Colour	• Variable: blackish-brown. • Striped or mottled.
Quota or non-quota	Non-quota.



© Photo by A.M.Arias via ictiorm.es

© Illustrations from Food and Agriculture Organization of the United Nations. Reproduced with permission.



Elegant cuttlefish

(*Sepia elegans*)

Identifying features

Tentacular club

- Short and oval.
- Six to eight rows of suckers and three to four larger suckers in the middle.

Body


- Elongated oval.
- Typically up to around 75mm mantle length.

Colour

- Red-brown, more yellow than common cuttlefish.

Quota or non-quota

Non-quota.



© Illustrations from Food and Agriculture Organization of the United Nations. Reproduced with permission.




Pink cuttlefish

(*Sepia orbignyana*)

Identifying features

Tentacular club	• Short and oval. • Five to six rows of suckers. • Three large suckers in the middle with one slightly smaller on each side of them.
Body	• Elongated oval. • Up to around 120mm mantle length. • Hard spine sticks out the back of the body.
Colour	• Red-orange, more red than common cuttlefish.
Quota or non-quota	Non-quota.



© Photo by A.M.Arias via ictioterm

© Illustrations from Food and Agriculture Organization of the United Nations. Reproduced with permission.